A Clone Resource of *Magnaporthe oryzae* Effectors That Share Sequence and Structural Similarities Across Host-Specific Lineages

Yohann Petit-Houdenot,^{1,2} Thorsten Langner,¹ Adeline Harant,¹ Joe Win,^{1,†} and Sophien Kamoun^{1,†}

¹ The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, NR4 7UK, U.K.

² UMR BIOGER, INRA, AgroParisTech, Université Paris-Saclay, Avenue Lucien Brétignières, BP 01, F-78850 Thiverval-Grignon, France

Abstract

The blast fungus *Magnaporthe oryzae* (syn. *Pyricularia oryzae*) is a destructive plant pathogen that can infect about 50 species of both wild and cultivated grasses, including important crops such as rice and wheat. *M. oryzae* is composed of genetically differentiated lineages that tend to infect specific host genera. To date, most studies of *M. oryzae* effectors have focused on the rice-infecting lineage. We describe a clone resource of 195 effectors of *Magnaporthe* species predicted from all the major host-specific lineages. These clones are freely available as Golden Gate-compatible entry plasmids. Our aim is to provide the community with an open source effector clone library to be used in a variety of functional studies. We hope that this resource will encourage studies of *M. oryzae* effectors on diverse host species.

Funding

This research was funded by the Gatsby Charitable Foundation, the Biotechnology and Biological Sciences Research Council, and European Research Council BLASTOFF project. Y. Petit-Houdenot was supported by a Young Scientist Contract from INRA.

Keywords

fungal effectors, genomics, proteomics

Resource Announcement

Plant pathogens secrete effectors that play central roles in subjugating plants for colonization. Fungal effectors typically have signal peptides and occasionally carry conserved folds and motifs (Franceschetti et al. 2017; Lo Presti et al. 2015). *Magnaporthe oryzae* (syn. *Pyricularia oryzae*) is an important plant pathogen that is able to infect around 50 species of both wild and cultivated grasses, including important cereals of the Poaceae family. *M. oryzae* is mostly known to cause rice blast but can also cause disease on other crops, such as barley, wheat, foxtail millet, and finger millet. The global population of *M. oryzae* is composed of genetically differentiated lineages that, in some cases, still exhibit a measurable degree of gene flow (Gladioux et al. 2018; Langner et al. 2018). Fungal isolates from each of those lineages show a preference for a specific host and also encode distinct repertoires of effector genes (Yoshida et al. 2016).

The first genomic sequence of *M. oryzae* was released in 2005 for the lab strain 70-15 and allowed researchers to predict a large set of secreted proteins, such as enzymes involved in secondary metabolism, and virulence-associated factors, including putative effectors (Dean et al. 2005). Recently an increasing number of genome sequences of isolates from

[†]Corresponding authors: J. Win; joe.win@tsl.ac.uk and S. Kamoun; sophien.kamoun@tsl.ac.uk

*The *e*-Xtra logo stands for “electronic extra” and indicates that a supplementary figure and a supplementary file are published online.

The author(s) declare no conflict of interest.

Accepted for publication 27 April 2020.

different lineages have become available, allowing the research community to perform comparative genomic studies (Chiapello et al. 2015; Yoshida et al. 2016). One of these lineages of *M. oryzae* causes the wheat blast disease and has recently emerged as a serious threat to food security in South East Asia (Islam et al. 2016, 2019). As part of the response to the wheat blast epidemic, we have committed to generating and sharing open source data and resources (Kamoun et al. 2019). This project is part of the wider OpenWheatBlast initiative.

Many of the validated effectors of *M. oryzae* are known as the MAX (*Magnaporthe* avirulence [AVR] and ToxB like) effectors. These effectors, while showing little primary sequence similarity, share a conserved structural fold composed of six β -sheets alternating in an antiparallel manner (de Guillen et al. 2015). The MAX family has expanded in *M. oryzae*, as MAX effectors account for 5 to 10% of the repertoire of secreted proteins and for approximately 50% of the already cloned effectors of *M. oryzae* (de Guillen et al. 2015). Indeed, the identification of structural motifs has enabled more sensitive predictions of effectors from pathogen genomes compared with sequence similarity searches (Franceschetti et al. 2017).

The aim of this project was to computationally identify a set of *M. oryzae* effectors from the main host-specific lineages and to develop an open access clone resource for functional analyses. To develop this resource, we initially analyzed secretomes of 13 *M. oryzae* isolates infecting different hosts, such as rice, wheat, finger millet, foxtail millet, and oat, and of two isolates of *M. grisea*—infecting species of the *Digitaria* genus. The secretomes used in this study have been obtained from the genome resources associated with the studies of Chiapello et al. (2015) and Yoshida et al. (2016). We searched for effector candidates in those secretomes using the following computational pipeline. First, we identified proteins showing sequence similarity to known *M. oryzae* effectors with an AVR activity (i.e., detected by plant immune receptors), using BLAST and a library of 21 previously characterized effectors. Those 21 effectors are AVR-PikD, AVR-Pia and AVR-Pii (Yoshida et al. 2009), AVR1-CO39 (Cesari et al. 2013), AVR-Piz-t (Li et al. 2009), AVR-Pita1 (Orbach et al. 2000), AVR-Pita3 (Khang et al. 2008), AVR-Pib (Zhang et al. 2015), MGG_17227-t26_1 (Zhong et al. 2018), AVR-Pi9 (Wu et al. 2015), AVR-Rmg8 (Anh et al. 2018), PWT3 and PWT4 (Inoue et al. 2017), Pwl2 (Sweigard et al. 1995), Pwl1, Pwl3, and Pwl4 (Kang et al. 1995), APikL1, APikL2, APikL3, and ms119 (*unpublished*). The BLAST search was performed with a blosum 62 substitution matrix, a gap opening penalty of 11, and a cut-off E-value of 10. Second, we performed a search for remote relationships using a hidden Markov model (HMM) trained with multiple, structure-based alignments of *M. oryzae* MAX effectors (de Guillen et al. 2015), using the HMMER program (Finn et al. 2011) and the HMM profile generated by de Guillen et al. (2015), with a cut-off E-value of 1e-4. We performed HMMsearch analyses with a cut-off E-value of 10. Next, we grouped candidates obtained by both approaches to form a nonredundant list of putative effectors and functionally annotated them using two methods. The first is a classical gene ontology (GO) search using the Blast2GO program (Conesa et al. 2005). The second is a BLASTP similarity search against a custom-made database (the Darwin database) that contains more than 2,600,000 predicted proteins of 137 different eukaryotic organisms. We removed candidates with functional domain annotations, as most *M. oryzae* effectors are proteins with no predicted function. Finally, we manually removed candidates with poor gene model predictions and low sequence similarity to *M. oryzae* effectors. The combination of these approaches resulted in 194 effector candidates that we selected for gene synthesis as a clone resource. We also added a homolog of the well-studied *M. oryzae* effector AVR-Pik from the *Lolium perenne* isolate PGKY to this library, bringing the total number to 195 predicted *M. oryzae* effectors.

The Golden Gate cloning system enables rapid and high throughput assembly of multiple sequence modules, such as promotor, terminator, or tags into a common vector (Patron et al. 2015). This cloning strategy is ideal for wide dissemination of cloning material by using a universal code for cloning. We synthesized the 195 *Magnaporthe* species effectors in a Golden Gate-compatible fashion to enable the transfer of these genes into a variety of vectors for various applications, such as yeast two-hybrid assays, heterologous protein expression, and fungal transformation (for example, the fungal transformation vectors described by Pennington et al. [2017, 2018]).

We synthesized the coding sequences corresponding to the mature proteins (without the signal peptide) of the 195 candidates in the Golden Gate-compatible vector pUC57-Kan. Each coding sequence was flanked by *Bsa*I restriction enzyme sites and relevant overhang

Table 1. Features of the 195 *Magnaporthe oryzae* effectors^a

Hosts	Number with similarity to		Total
	AVR effectors	MAX effectors	
Rice	23	27	50
Wheat	6	5	11
<i>Setaria</i> spp.	3	3	6
<i>Eleusine</i> spp.	6	2	8
<i>Avena</i> spp.	3	0	3
<i>Digitaria</i> spp.	20	19	39
<i>Lolium</i> spp.	1	0	1
Multiple hosts ^b	38	39	77
	100	95	195

^a Candidate effectors were selected based on their sequence identity to a known effector of *M. oryzae* or structural similarity with MAX (*Magnaporthe* avirulence [AVR] and ToxB like) effectors according to the hidden Markov model search.

^b Those isolates that infect more than one host among the seven listed hosts.

sequences for Golden Gate cloning reactions. The coding sequences were manually codon-optimized for expression in *Saccharomyces cerevisiae*, *Nicotiana benthamiana*, and *Escherichia coli* (when it was not possible to optimize for all three organisms, priority was given to *S. cerevisiae* and *N. benthamiana*). Table 1 summarizes the different candidates identified, their host specificity, and whether they were selected based on sequence or structural similarities. Supplementary File 1 describes the 195 candidates and includes information such as sequence similarity to *M. oryzae* effectors, results of the HMM search, SignalP scores, functional annotation, and host specificity of the strain in which the effector candidate was identified. An overview of the Golden Gate system and a generic schematic representation of the pUC57-Kan vector insertion site is shown in the Supplementary Figure. The protein sequences and codon-optimized sequences of the selected genes as well as additional information including the plasmid maps of the Level 0 constructs are provided through the data repository Zenodo.

Level 0 modules are available via Addgene. Catalog numbers are provided in Supplementary File S1.

Author-Recommended Internet Resources

Addgene: https://www.addgene.org/Sophien_Kamoun

OpenWheatBlast: <http://wheatblast.net>

Zenodo repository, Kamoun lab Darwin database: <http://doi.org/10.5281/zenodo.3699564>; and for sequences of candidate effectors: <http://doi.org/10.5281/zenodo.3268775>

Literature Cited

- Anh, V. L., Inoue, Y., Asuke, S., Vy, T. T. P., Anh, N. T., Wang, S., Chuma, I., and Tosa, Y. 2018. Rmg8 and Rmg7, wheat genes for resistance to the wheat blast fungus, recognize the same avirulence gene *AVR-Rmg8*. *Mol. Plant Pathol.* 19: 1252-1256.
- Cesari, S., Thilliez, G., Ribot, C., Chalvon, V., Michel, C., Jauneau, A., Rivas, S., Alaux, L., Kanzaki, H., Okuyama, Y., Morel, J.-B., Fournier, E., Tharreau, D., Terauchi, R., and Kroj, T. 2013. The rice resistance protein pair RGA4/RGA5 recognizes the *Magnaporthe oryzae* effectors AVR-Pia and AVR1-CO39 by direct binding. *Plant Cell* 25:1463-1481.
- Chiapello, H., Mallet, L., Guérin, C., Aguilera, G., Amselem, J., Kroj, T., Ortega-Abboud, E., Lebrun, M.-H., Henrissat, B., Gendrait, A., Rodolphe, F., Tharreau, D., and Fournier, E. 2015. Deciphering genome content and evolutionary relationships of isolates from the fungus *Magnaporthe oryzae* attacking different host plants. *Genome Biol. Evol.* 7:2896-2912.
- Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M., and Robles, M. 2005. Blast2GO: A universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21:3674-3676.
- de Guillen, K., Ortiz-Vallejo, D., Gracy, J., Fournier, E., Kroj, T., and Padilla, A. 2015. Structure analysis uncovers a highly diverse but structurally conserved effector family in phytopathogenic fungi. *PLoS Pathog.* 11:e1005228.
- Dean, R. A., Talbot, N. J., Ebbole, D. J., Farman, M. L., Mitchell, T. K., Orbach, M. J., Thon, M., Kulkarni, R., Xu, J.-R., Pan, H., Read, N. D., Lee, Y.-H., Carbone, I., Brown, D., Oh, Y. Y., Donofrio, N., Jeong, J. S., Soanes, D. M., Djonovic, S., Kolomiets, E., Rehmeier, C., Li, W., Harding, M., Kim, S., Lebrun, M.-H., Bohnert, H., Coughlan, S., Butler, J., Calvo, S., Ma, L.-J., Nicol, R., Purcell, S., Nusbaum, C., Galagan, J. E., and Birren, B. W. 2005. The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature* 434:980-986.
- Finn, R. D., Clements, J., and Eddy, S. R. 2011. HMMER web server: Interactive sequence similarity searching. *Nucleic Acids Res.* 39:W29-W37.
- Franceschetti, M., Maqbool, A., Jiménez-Dalmaroni, M. J., Pennington, H. G., Kamoun, S., and Banfield, M. J. 2017. Effectors of filamentous plant pathogens: Commonalities amid diversity. *Microbiol. Mol. Biol. Rev.* 81:81.
- Gladieux, P., Ravel, S., Rieux, A., Cros-Arteil, S., Adreit, H., Milazzo, J., Thierry, M., Fournier, E., Terauchi, R., and Tharreau, D. 2018. Coexistence of multiple endemic and pandemic lineages of the rice blast pathogen. *MBio* 9:e01806-e01817.

- Inoue, Y., Vy, T. T. P., Yoshida, K., Asano, H., Mitsuoka, C., Asuke, S., Anh, V. L., Cumagun, C. J. R., Chuma, I., Terauchi, R., Kato, K., Mitchell, T., Valent, B., Farman, M., and Tosa, Y. 2017. Evolution of the wheat blast fungus through functional losses in a host specificity determinant. *Science* 357:80-83.
- Islam, M. T., Croll, D., Gladioux, P., Soanes, D. M., Persoons, A., Bhattacharjee, P., Hossain, M. S., Gupta, D. R., Rahman, M. M., Mahboob, M. G., Cook, N., Salam, M. U., Surovy, M. Z., Sancho, V. B., Maciel, J. L., Nhanijúnior, A., Castroagudín, V. L., Reges, J. T., Ceresini, P. C., Ravel, S., Kellner, R., Fournier, E., Tharreau, D., Lebrun, M. H., McDonald, B. A., Stitt, T., Swan, D., Talbot, N. J., Saunders, D. G., Win, J., and Kamoun, S. 2016. Emergence of wheat blast in Bangladesh was caused by a South American lineage of *Magnaporthe oryzae*. *BMC Biol.* 14:84.
- Islam, M. T., Kim, K. H., and Choi, J. 2019. Wheat blast in Bangladesh: The current situation and future impacts. *Plant Pathol. J.* 35:1-10.
- Kamoun, S., Talbot, N. J., and Islam, M. T. 2019. Plant health emergencies demand open science: Tackling a cereal killer on the run. *PLoS Biol.* 17: e3000302.
- Kang, S., Sweigard, J. A., and Valent, B. 1995. The *PWL* host specificity gene family in the blast fungus *Magnaporthe grisea*. *Mol. Plant-Microbe Interact.* 8:939-948.
- Khang, C. H., Park, S.-Y., Lee, Y.-H., Valent, B., and Kang, S. 2008. Genome organization and evolution of the *AVR-Pita* avirulence gene family in the *Magnaporthe grisea* species complex. *Mol. Plant-Microbe Interact.* 21:658-670.
- Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E. L. 2001. Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes. *J. Mol. Biol.* 305:567-580.
- Langner, T., Bialas, A., and Kamoun, S. 2018. The blast fungus decoded: Genomes in flux. *MBio* 9:e00571-18.
- Li, W., Wang, B., Wu, J., Lu, G., Hu, Y., Zhang, X., Zhang, Z., Zhao, Q., Feng, Q., Zhang, H., Wang, Z., Wang, G., Han, B., Wang, Z., and Zhou, B. 2009. The *Magnaporthe oryzae* avirulence gene *AvrPiz-t* encodes a predicted secreted protein that triggers the immunity in rice mediated by the blast resistance gene *Piz-t*. *Mol. Plant-Microbe Interact.* 22:411-420.
- Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A., Reissmann, S., and Kahmann, R. 2015. Fungal effectors and plant susceptibility. *Annu. Rev. Plant Biol.* 66:513-545.
- Orbach, M. J., Farrall, L., Sweigard, J. A., Chumley, F. G., and Valent, B. 2000. A telomeric avirulence gene determines efficacy for the rice blast resistance gene *Pi-ta*. *Plant Cell* 12:2019-2032.
- Patron, N. J., Orzaez, D., Marillonnet, S., Warzecha, H., Matthewman, C., Youles, M., Raitskin, O., Leveau, A., Farré, G., Rogers, C., Smith, A., Hibberd, J., Webb, A. A. R., Locke, J., Schormack, S., Ajioka, J., Baulcombe, D. C., Zipfel, C., Kamoun, S., Jones, J. D. G., Kuhn, H., Robatzek, S., Van Esse, H. P., Sanders, D., Oldroyd, G., Martin, C., Field, R., O'Connor, S., Fox, S., Wulff, B., Miller, B., Breakspear, A., Radhakrishnan, G., Delaux, P.-M., Loqué, D., Granell, A., Tissier, A., Shih, P., Brutnell, T. P., Quick, W. P., Rischer, H., Fraser, P. D., Aharoni, A., Raines, C., South, P. F., Ané, J.-M., Hamberger, B. R., Langdale, J., Stougaard, J., Bouwmeester, H., Udvardi, M., Murray, J. A. H., Ntoukakis, V., Schäfer, P., Denby, K., Edwards, K. J., Osbourn, A., and Haseloff, J. 2015. Standards for plant synthetic biology: A common syntax for exchange of DNA parts. *New Phytol.* 208:13-19.
- Pennington, H. G., Youles, M., and Kamoun, S. 2017. Golden-Gate compatible *Magnaporthe oryzae* protoplast transformation vectors. Figshare 10.6084/m9.figshare.5375428.v1. Published online.
- Pennington, H. G., Youles, M., and Kamoun, S. 2018. Golden-Gate compatible *Magnaporthe oryzae* agrobacterium transformation vectors. Figshare 10.6084/m9.figshare.5808348.v1. Published online.
- Sweigard, J. A., Carroll, A. M., Kang, S., Farrall, L., Chumley, F. G., and Valent, B. 1995. Identification, cloning, and characterization of *PWL2*, a gene for host species specificity in the rice blast fungus. *Plant Cell* 7:1221-1233.
- Wu, J., Kou, Y., Bao, J., Li, Y., Tang, M., Zhu, X., Ponaya, A., Xiao, G., Li, J., Li, C., Song, M. Y., Cumagun, C. J., Deng, Q., Lu, G., Jeon, J. S., Naqvi, N. I., and Zhou, B. 2015. Comparative genomics identifies the *Magnaporthe oryzae* avirulence effector *AvrPi9* that triggers *Pi9*-mediated blast resistance in rice. *New Phytol.* 206:1463-1475.
- Yoshida, K., Saitoh, H., Fujisawa, S., Kanzaki, H., Matsumura, H., Yoshida, K., Tosa, Y., Chuma, I., Takano, Y., Win, J., Kamoun, S., and Terauchi, R. 2009. Association genetics reveals three novel avirulence genes from the rice blast fungal pathogen *Magnaporthe oryzae*. *Plant Cell* 21:1573-1591.
- Yoshida, K., Saunders, D. G. O., Mitsuoka, C., Natsume, S., Kosugi, S., Saitoh, H., Inoue, Y., Chuma, I., Tosa, Y., Cano, L. M., Kamoun, S., and Terauchi, R. 2016. Host specialization of the blast fungus *Magnaporthe oryzae* is associated with dynamic gain and loss of genes linked to transposable elements. *BMC Genomics* 17:370.
- Zhang, S., Wang, L., Wu, W., He, L., Yang, X., and Pan, Q. 2015. Function and evolution of *Magnaporthe oryzae* avirulence gene *AvrPib* responding to the rice blast resistance gene *Pib*. *Sci. Rep.* 5:11642.
- Zhong, Z., Chen, M., Lin, L., Han, Y., Bao, J., Tang, W., Lin, L., Lin, Y., Somai, R., Lu, L., Zhang, W., Chen, J., Hong, Y., Chen, X., Wang, B., Shen, W. C., Lu, G., Norvienyeku, J., Ebbola, D. J., and Wang, Z. 2018. Population genomic analysis of the rice blast fungus reveals specific events associated with expansion of three main clades. *ISME J.* 12:1867-1878.