New Linear Lipopeptides Produced by *Pseudomonas cichorii* SF1-54 Are Involved in Virulence, Swarming Motility, and Biofilm Formation

Ellen Pauwelyn,^{1,2} Chien-Jui Huang,¹ Marc Ongena,³ Valérie Leclère,⁴ Philippe Jacques,⁴ Peter Bleyaert,² Herbert Budzikiewicz,⁵ Mathias Schäfer,⁵ and Monica Höfte¹

¹Department of Crop Protection, Laboratory of Phytopathology, Ghent University, Coupure Links 653, 9000 Ghent, Belgium;

²Inagro vzw, Ieperseweg 87, 8800 Rumbek, Belgium; ³Walloon Centre for Industrial Biology, University of Liège–Gembloux Agro-Bio Tech, Passage des Déportés 2, 5030 Gembloux, Belgium; ⁴Laboratoire de Procédés Biologiques, Génie

Enzymatique et Microbien (ProBioGEM), Université de Lille Sciences et Technologies, F-59655 Villeneuve d'Ascq Cedex,

France; ⁵Institut für Organische Chemie, Universität zu Köln, Greinstr. 4, 50939 Köln, Germany

Submitted 2 November 2012. Accepted 5 February 2013.

Pseudomonas cichorii is the causal agent of lettuce midrib rot, characterized by a dark-brown to green-black discoloration of the midrib. Formation of necrotic lesions by several plant-pathogenic pseudomonads is associated with production of phytotoxic lipopeptides, which contribute to virulence. Therefore, the ability of *P. cichorii* SF1-54 to produce lipopeptides was investigated. A cell-free culture filtrate of SF1-54 showed surfactant, antimicrobial, and phytotoxic activities which are typical for lipopeptides. High-performance liquid chromatography analysis of *P. cichorii* SF1-54 culture filtrate revealed the presence of seven compounds with lipopeptide characteristics. Two related lipopeptides, named cichofactin A and B, were studied in more detail: they are linear lipopeptides with a decanoic and dodecanoic lipid chain, respectively, connected to the N-terminus of an eight-amino-acid peptide moiety. Both cichofactins are new members of the syringafactin lipopeptide family. Furthermore, two nonribosomal peptide synthetase-encoding genes, *cifA* and *cifB*, were identified as responsible for cichofactin biosynthesis. A *cifAB* deletion mutant no longer produced cichofactins and was impaired in swarming motility but showed enhanced biofilm formation. Upon spray inoculation on lettuce, the cichofactin-deficient mutant caused significantly less rotten midribs than the wild type, indicating that cichofactins are involved in pathogenicity of *P. cichorii* SF1-54. Further analysis revealed that *P. cichorii* isolates vary greatly in swarming motility and cichofactin production.

Many plant-pathogenic bacteria are known to produce non-host-specific phytotoxins that contribute to virulence (Kunkel and Zhongying 2006). Well-characterized phytotoxins are those produced by *Pseudomonas syringae* and include chlorosis-inducing toxins (e.g., coronatine, phaseolotoxin, and tabtoxin) and necrosis-inducing toxins (e.g., syringomycin and syringopeptin) (Bender et al. 1999). The latter are cyclic lipopeptides

(CLP), which are amphipathic molecules composed of a fatty acid tail linked to a cyclic oligopeptide head. Due to this amphipathic nature, they can insert into the plasma membrane of their host cells, which is their primary target, and form pores. This results in transmembrane fluxes of H⁺, Ca²⁺, and K⁺ and leads to a collapse of the pH gradient across the plasma membrane, consequently resulting in cell death. This process is highly efficient, because only a few nanomoles of toxin suffice to induce pore formation (Bender and Scholz-Schroeder 2004; Bender et al. 1999). In addition to phytotoxicity, these CLP toxins also exhibit hemolytic and antimicrobial activity (Hutchison and Gross 1997; Hutchison et al. 1995; Lavermicocca et al. 1997). This latter feature of lipopeptides has been exploited by researchers to develop bioassays that have expedited detection of lipopeptide production, purification, and genetic studies on these cyclic lipopeptides.

However, lipopeptides are not always phytotoxic. Depending on their structure, they may have diverse other roles and functions and can be involved in swarming motility, biofilm development, and colonization, or display zoosporicidal or antimicrobial activity (Raaijmakers et al. 2006, 2010). These functions are extensively described for antagonistic *Pseudomonas* spp. but to a lesser extent for plant-pathogenic pseudomonads. The only available examples to date are the syringafactins, which are essential for swarming motility of *P. syringae* pv. *tomato* DC3000 and *P. syringae* pv. *syringae* B728a (Berti et al. 2007; Burch et al. 2010) and viscosin, which is required for *P. fluorescens* 5064 to colonize the surface of broccoli florets (Hildebrand et al. 1998).

P. cichorii is an important plant-pathogenic bacterium, with a wide host range and a worldwide distribution. *P. cichorii* causes necrotic leaf and stem lesions on vegetables, ornamentals, cereals, and woody plants (Bradbury 1986). Economically important hosts include lettuce, celery, chrysanthemum, tomato, coffee, and soybean (Bradbury 1986; Jagger 1914; Jones et al. 1984; Wilk and Dye 1974; Yu and Lee 2012). In Belgian greenhouse-grown butterhead lettuce, *P. cichorii* causes a dark-brown to green-black discoloration of the midrib (Cottyn et al. 2009, 2011; Pauwelyn et al. 2011). Symptom development on lettuce leaves is strongly associated with apoptosis-like programmed cell death (Kiba et al. 2006, 2009). Intriguingly, the type III secretion system is essential for pathogenicity of *P. cichorii* on eggplant but not on lettuce (Hojo et al. 2008; Kajihara et al. 2012). Symptom development may be associ-

Corresponding author: Monica Höfte; E-mail: monica.hofte@ugent.be

*The e-Xtra logo stands for “electronic extra” and indicates that four supplementary figures and three supplementary tables are published online.

ated with the production of necrosis-inducing cyclic lipopeptide toxins by this bacterium. The production of antimicrobial and phytotoxic substances by *P. cichorii* has already been described (Hikichi et al. 1998; Hu et al. 1998; Lazzaroni et al. 2003) but no bioactive compound has been characterized to date.

Therefore, the aim of this study was to examine the ability of *P. cichorii* to produce phytotoxic and antimicrobial compounds using bioassays. Seven bioactive compounds produced in culture by the highly virulent *P. cichorii* SF1-54 were isolated and partly characterized by reverse-phase high-performance liquid chromatography (RP-HPLC) and liquid chromatography electrospray ionization mass spectrometry (LC-ESI-MS). The structures of two of these lipopeptides produced by *P. cichorii* SF1-54 were elucidated using LC-ESI-MS and high-resolution (HR)-ESI tandem mass spectrometry (MS-MS). They appear to be linear lipopeptides, consisting of eight amino acids, and are novel members of the syringafactin family of lipopeptides. In addition, genome mining of *P. cichorii* SF1-54 revealed two nonribosomal peptide synthetase (NRPS) genes, *cifA* and *cifB*, that encode the lipopeptide synthetases responsible for biosynthesis of cichofactins. Based on the *cifA* and *cifB* gene sequences, a cichofactin-deficient mutant was constructed to investigate the function of these compounds for *P. cichorii* SF1-54.

RESULTS

Biological activity of strain SF1-54 and its secreted metabolites.

Injection of a *P. cichorii* SF1-54 bacterial suspension (1×10^8 CFU/ml) into lettuce or chicory leaves resulted in brown lesions in the treated area within 24 h postinoculation (Fig. 1A). Injection of a crude culture filtrate of *P. cichorii* SF1-54 caused identical symptoms (Fig. 1B), suggesting that a phytotoxic compound is produced and secreted into the SRM_{AF} medium (Gross 1985). No symptom was visible by injection of noninoculated medium in chicory. Although minor rotting symptoms were often observed on the lettuce leaves of the noninoculated treatments, they looked clearly different from the symptoms caused by *P. cichorii* SF1-54 and its crude culture filtrate, indicating that they are possibly due to wounding, contamination, or senescence (Fig. 1C).

Symptom development is similar in lettuce and chicory leaves; therefore, only chicory leaves were used for further phytotoxicity tests, because they are less prone to contamination and easier to handle for reproducible infection.

P. cichorii SF1-54 inhibited growth of *Bacillus megaterium* (a bacterium), *Geotrichum candidum* (a filamentous fungus), and *Rhodotorula mucilaginosa* (a yeast). The indicator microorganisms were selected according to the spectrum of antimicrobial activities of *P. syringae* pv. *syringae* toxins (Lavermicocca et al. 1997). In direct antagonistic assays

against these microorganisms, a *P. cichorii* SF1-54 colony produced inhibition zones of 25 to 41 mm against *B. megaterium*, 38 to 53 mm against *G. candidum*, and 50 to 56 mm against *R. mucilaginosa* under our experimental conditions. The crude culture filtrate and semipurified culture extract (solid-phase extraction on C18 cartridge) retained these inhibitory capacities against *B. megaterium*, *G. candidum*, and *R. mucilaginosa* (Fig. 2B).

The active molecules produced by strain SF1-54 in SRM_{AF} medium were isolated from the semipurified culture extracts by repeated injections in semipreparative RP-HPLC. Repeated injections were performed on a protein and peptide C-18 column. The eluates were monitored spectrophotometrically at 220 nm. All eluates absorbing at 220 nm were independently collected in 23 fractions, corresponding to the observed peaks in the RP-HPLC chromatogram (Fig. 2A). All fractions were first tested for surfactant activity and seven compounds (Fig. 2A, letters A to G) caused drop collapse, which is characteristic of biosurfactant metabolites.

Antimicrobial activity and phytotoxicity of these purified molecules were then assayed (Fig. 2B). Compound A inhibited the growth of *B. megaterium*. Compound B had antimicrobial activity against *G. candidum* and *R. mucilaginosa*, while compound C only inhibited *R. mucilaginosa*. Compounds D and E were both inhibitory to *R. mucilaginosa* and *B. megaterium*. Compounds F and G were both phytotoxic and inhibited *B. megaterium*. Thus, the culture filtrate of *P. cichorii* SF1-54 is a complex mixture of several bioactive metabolites.

Chemical characterization of compounds D and E (cichofactins).

The concentration of compounds D and E in the culture filtrate obtained from SRM_{AF} medium was 29 ± 15 mg/liter based on calculated data from five different cultures. Several media other than SRM_{AF} and various incubation periods and temperatures were tested for production of all the bioactive compounds by *P. cichorii* SF1-54 but only compounds D and E were consistently produced in high amounts (data not shown), indicating that these metabolites could be the main compounds produced by *P. cichorii* SF1-54. In addition to consistent production, compounds D and E exhibited not only surfactant but also inhibitory activities to *R. mucilaginosa* and *B. megaterium*. Thus, we first focused on identification and characterization of these two metabolites.

HPLC-purified samples of compounds D and E were submitted to acid hydrolysis, and further analysis of the released amino acids revealed that both molecules contained Leu, Val, and Glx (representing either Gln or Glu) residues in similar molar ratios of 1.91:1.05:5.12 and 2.02:1.03:5.01, respectively. These results support the peptidic nature of compounds D and E and they were further characterized by HPLC-ESI-MS using parameters to optimize in-source fragmentation. Identification of b and y'' ions revealed very similar fragmentation patterns

Fig. 1. Necrosis induction in chicory and lettuce leaves at 24 h after inoculation with **A**, *Pseudomonas cichorii* SF1-54 (1×10^8 CFU/ml); **B**, crude culture filtrate of *P. cichorii* SF1-54; and **C**, noninoculated SRM_{AF} medium (the noninoculated control treatment). Each treatment was repeated three times, and a picture representative of each of the treatments is shown.

of the $[M+H]^+$ ions (m/z 1,109 and 1,137, respectively) corresponding to compounds D and E (Table 1; Supplementary Fig. S1). However, m/z values of b-ion fragments were 28 Da larger for compound E (Table 1), suggesting that its fatty acid chain contains two extra methylene groups. The presence of a 3-hydroxydecanoyl fatty acid in compound D and a 3-hydroxydecanoyl fatty acid in compound E is fully supported by the detection of ions at m/z 284 and 312, respectively, in accordance to the [FA-Leu]-fragments.

Based on the fragmentation pattern, both lipopeptides could have the peptide sequence Leu-Leu-Gln-Leu-Gln-Val-Leu-Leu.

HR-ESI-MS-MS analyses.

In-depth structural characterization was obtained using HR-ESI-MS-MS performed on the $[M+Na]^+$ form of compounds D and E. The exact m/z of $[M+Na]^+$ of compound D is 1,131.732 and of compound E is 1,159.762 (calculated masses are 1,131.737 and 1,159.768 for $C_{55}H_{100}N_{10}O_{13}Na$ and $C_{57}H_{104}N_{10}O_{13}Na$, respectively). These masses suggest that the compounds are linear because their mass would be 18 Da smaller (loss of one water molecule for cyclization) if they were cyclic. Moreover, the first loss of Leu-COOH (m/z 1,018 and 1,046, respectively) is an additional evidence for a linear structure.

Fig. 2. A, Reverse-phase high-performance liquid chromatography (RP-HPLC) chromatogram of the semipurified extract of *Pseudomonas cichorii* SF1-54. Each peak was checked for surfactant activity, antimicrobial activity, and phytotoxic activity in chicory leaves. All (bio)active peaks are indicated with a letter. B, Antimicrobial and phytotoxic activity of each bioactive fraction separated by RP-HPLC. Antimicrobial activity against *Bacillus megaterium* (Bm), *Geotrichum candidum* (Gc), and *Rhodotorula mucilaginosa* (Rm) and phytotoxicity to chicory (PT) of each fraction was assayed. CF: culture filtrate of *P. cichorii* SF1-54.

All pertinent fragments from the HR-ESI-MS-MS spectra of the $[M+Na]^+$ form of compounds D and E are shown in Table 2. The b-type ions are formed upon collision activation (CA) of the $[M+Na]^+$ species by rearrangement processes (Fuchs and Budzikiewicz 2001) (Supplementary Fig. S2). The degradation of the peptide chain by elimination of the C-

Table 1. Listing of b and y'' ions obtained from the $[M+H]^+$ species of compounds D and E formed in the liquid chromatography electrospray ionization mass spectrometry analysis^a

Peptide	b ions		y'' ions	
	Compound D	Compound E	Compound D	Compound E
FA	(171)	(199)	...	...
Leu	284	312	(939)	(939)
Leu	397	425	(826)	(826)
Gln	525	553	713	713
Leu	638	666	585	585
Gln	766	794	472	472
Val	865	893	344	344
Leu	978	1,006	245	245
Leu	1,091	1,119	(132)	(132)
$[M+H]^+$	1,109	1,137	...	...

^a Numbers in parentheses are calculated values not observed in the mass spectra.

Table 2. Listing of fragment ions and of y'' ions obtained from the $[M+Na]^+$ species of compounds D and E formed in the high-resolution electrospray ionization tandem mass spectrometry analysis^a

Peptide	Fragment ions				y'' ions	
	Compound D		Compound E		Compound D	Compound E
FA	(211)	(165)	(239)	(193)	...	...
Leu	324	(278)	(352)	(306)	(961)	(961)
Leu	437	391	465	419	848	848
Gln	565	519	593	547	735	735
Leu	678	632	706	660	607	607
Gln	806	760	834	788	494	494
Val	905	859	933	887	(366)	(366)
Leu	1,018	972	1,046	1,000	(267)	(267)
Leu ($[M+H]^+$)	1,131	...	1,159	...	(154)	(154)

^a Numbers in parentheses are calculated values not observed in the mass spectra.

terminal amino acid with back transfer of the hydroxyl group can be repeated several times (sequential loss of $NH = CHR$ units). Thus, the HR-ESI-MS-MS fragmentation pattern supported the LC-ESI-MS data completely: a 3-hydroxydecanooyl fatty acid in compound D and a 3-hydroxydodecanooyl fatty acid in compound E coupled to a peptide moiety containing Leu-Leu-Gln-Leu-Gln-Val-Leu-Leu for both compounds.

Comparison of compounds D and E with previously described lipopeptides produced by *Pseudomonas* spp. revealed that these molecules are new members of the syringafactin family. The only difference is the replacement of Thr present at position five in syringafactins A and D by Gln in the compounds produced by *P. cichorii* SF1-54. Because of the high structural resemblance of these compounds to the syringafactins A and D, we renamed compounds D and E cichofactins A and B, respectively.

Identification and bioinformatic analysis of putative cichofactin synthetase-encoding genes.

Because cichofactins show high structural similarities with the syringafactins of *P. syringae* pv. *tomato* DC3000, the genome of *P. cichorii* SF1-54 (unpublished) was compared with the genome sequence of *P. syringae* pv. *tomato* DC3000 using

Fig. 3. Schematic representation of the cichofactin biosynthetic gene clusters and the flanking open reading frames in the *Pseudomonas cichorii* SF1-54 genome. Module and domain organization are indicated below *cifA* and *cifB* genes. Predicted amino acid specificity is represented below each module and all amino acids are identified by standard three-letter biochemical notation. The region between arrowheads was deleted in the cichofactin-deficient mutant.

the MAUVE multiple genome alignment system to identify the gene cluster which encodes the NRPS for cichofactins synthesis.

In the genome of *P. cichorii* SF1-54, a gene cluster was identified containing two large genes, *cifA* and *cifB*, that are 9,498 and 17,734 bp in size and encode for three and five NRPS modules, respectively, with specific condensation (C), adenylation (A), and thiolation (T) domains (Fig. 3). Both genes exhibit two typical features of NRPS genes involved in lipopeptide synthesis: the presence of an N-terminal C domain in the initiating module and two tandem thioesterase domains at the end of the NRPS cluster. Furthermore, *cifA* and *cifB* genes of *P. cichorii* SF1-54 showed 76 and 75% identity at the amino acid level to syringafactin-encoding genes *syfA* and *syfB*, respectively, of *P. syringae* pv. *tomato* DC3000.

To predict the peptide sequence of cichofactins from genome sequencing data, bioinformatic analysis of the A domain from each NRPS module was performed. The putative peptide sequence of cichofactins is Leu-Leu-Gln/Asp-Leu-Gln/Asp-Val-Leu-Leu (Supplementary Table S1). The prediction with NRPSpredictor2 for the amino acids at third and fifth position is based on the specificity codes which, in both cases, show 70% identity with the specificity code of Gln and Asp. According to the results of structural analysis of cichofactins, both the third and fifth amino acids are Gln and the prediction is in full agreement with the peptide sequence of cichofactins determined by MS.

Furthermore, the D/L configuration of amino acid residues of cichofactins was investigated by phylogenetic analysis of the peptide sequences of C domains. In *Pseudomonas* spp., dual condensation/epimerization (C/E) domains are responsible for generating D-residues into lipopeptides (Balibar et al. 2005; Rausch et al. 2007). According to the phylogenetic analysis, C domains of modules 2, 3, 4, 6, and 8 could be dual C/E domains; therefore, the peptidic structure of cichofactins could be D-Leu-D-Leu-D-Gln-L-Leu-D-Gln-L-Val-D-Leu-L-Leu. The locations of dual C/E domains of cichofactins are identical to those of syringafactins (Berti et al. 2007).

LuxR-type regulators and transporter genes are often positioned up- and downstream of the lipopeptide biosynthesis genes in various pseudomonads (Berti et al. 2007; de Bruijn and Raaijmakers 2009; de Bruijn et al. 2008; Dubern et al. 2008; Kang and Gross 2005; Quigley et al. 1993; Roongsawang et al. 2003; Vallet-Gely et al. 2010). Therefore, we examined the flanking regions of the cichofactin biosynthesis cluster in the *P. cichorii* SF1-54 genome. Two open reading frames (ORF) downstream of the cichofactin biosynthesis cluster showed sequence similarities to subunits of a multidrug efflux pump, MacA (77% protein identity with *P. entomophila* L48) and MacB (89% protein identity with *P. syringae* pv. *tomato* DC3000) (Fig. 3). On each side of the cichofactin biosynthesis cluster, two ORF were identified showing sequence similarity with the LuxR family of transcriptional regulators (66 and 69% protein identity with *P. syringae* pv. *tabaci* ATCC 11528 and *P. syringae* pv. *maculicola* ES4326, respectively).

Biological role of cichofactins produced by *P. cichorii* SF1-54.

To investigate the biological role of the cichofactins and their importance in pathogenicity for *P. cichorii* SF1-54, we constructed a cichofactin-deficient mutant, SF1-54- Δ *cifAB*, by deleting a 9,715-bp fragment within the cichofactin-encoding genes *cifA* and *cifB* (Fig. 3). LC-ESI-MS analysis confirmed that this mutant was completely abolished in production of both cichofactins (Fig. 4). In addition, the production of the phytotoxic compounds F and G was approximately 10 times higher in the mutant compared with the wild type. Deletion of the cichofactin biosynthesis genes did not influence the production of compound A, while the amounts of compounds B and C were too low to be detected in the semipurified culture filtrate of the wild type and the cichofactin mutant by LC-ESI-MS analysis.

When *P. cichorii* SF1-54 cells were suspended into a droplet of water on parafilm, the water droplet collapsed immediately, but a cell suspension of *P. cichorii* SF1-54- Δ *cifAB* did not

Fig. 4. Liquid chromatography electrospray ionization mass spectrometry analysis of culture supernatants of *Pseudomonas cichorii* SF1-54 and its cichofactin-deficient mutant.

cause spreading of the water on the parafilm surface (data not shown). Because of the biological activity of the remaining compounds, the cichofactin-deficient mutant retained the ability to inhibit growth of *B. megaterium*, *G. candidum*, and *R. mucilaginosa* in direct antagonistic assays. However, *P. cichorii* SF1-54- Δ cifAB produced remarkably larger inhibition zones than the wild type against *B. megaterium*, probably due to the overproduction of compounds F and G. The mean inhibition

zone of *P. cichorii* SF1-54- Δ cifAB was 47 mm compared with 35 mm for *P. cichorii* SF1-54.

The cichofactin-deficient mutant still induced discoloration and necrosis of chicory leaves, which is not surprising because the mutant overproduces the phytotoxic compounds F and G (Fig. 4).

Syringafactins produced by *P. syringae* pv. *tomato* DC3000 and *P. syringae* pv. *syringae* B748a function as biosurfactants

Fig. 5. Biological roles of cichofactins in *Pseudomonas cichorii* SF1-54. **A**, Swarming behavior of *P. cichorii* SF1-54 and its cichofactin-deficient mutant on Luria-Bertani soft agar (0.5% [wt/vol]) plates. *P. cichorii* SF1-54 covered the entire plate within 48 h after inoculation (left panel), whereas its cichofactin-deficient mutant was impaired in motility (right panel). **B**, Biofilm formation of *P. cichorii* SF1-54Nal^R and its cichofactin-deficient mutant. Spectrophotometric quantification of the biofilm formed by *P. cichorii* SF1-54Nal^R and its cichofactin-deficient mutant (*P. cichorii* SF1-54- Δ cifAB) 24 h after inoculation; the quantity of crystal-violet-stained biofilm was determined by measuring optical density at 595 nm. The experiment (seven repetitions per treatment) was carried out three times and similar results were obtained each time. Representative data from one experiment are presented. Bars indicated with the same letter are not statistically different based on nonparametric Kruskal-Wallis and Mann-Whitney comparisons ($P < 0.05$). **C**, Virulence of *P. cichorii* SF1-54Nal^R and its cichofactin-deficient mutant (SF1-54- Δ cifAB) on greenhouse-grown butterhead lettuce. Plants were inoculated with suspensions of *P. cichorii* strains (1×10^6 CFU/ml) after head formation of lettuce. Virulence is represented as disease severity (left panel) and mean number of diseased midribs (right panel). Tap water was used as the noninoculated control. The experiment was carried out twice with similar results. Data from one experiment are shown. Bars indicated with the same letter are not statistically different based on nonparametric Kruskal-Wallis and Mann-Whitney comparisons ($P < 0.05$).

and are involved in swarming motility (Berti et al. 2007; Burch et al. 2010). *P. cichorii* SF1-54 swarmed outward from the inoculation point to cover the entire plate containing soft Luria-Bertani (LB) medium (0.5% agar) within 48 h, whereas swarming motility was completely abolished in the cichofactin-deficient mutant (Fig. 5A).

Because lipopeptides produced by plant-associated *Pseudomonas* spp. are often described to influence biofilm formation (de Bruijn et al. 2007, 2008; Kuiper et al. 2004; Roongsawang et al. 2003), we tested the influence of cichofactins on biofilm production of *P. cichorii* SF1-54. *P. cichorii* SF1-54 formed biofilms in polystyrene 96-well microtiter plates, and the biofilm formation of the cichofactin-deficient mutant was significantly improved compared with the wild type (Fig. 5B).

To determine the effect of cichofactins on the virulence of *P. cichorii* SF1-54, the wild type or its cichofactin-deficient mutant were spray-inoculated on butterhead lettuce in greenhouse experiments. Lettuce plants inoculated with the cichofactin-deficient mutant exhibited significantly less rotten midribs than lettuce plants inoculated with the wild type, which is also reflected in the significantly lower disease severity score obtained with the mutant ($P < 0.05$) (Fig. 5C). However, when the wild type or the *cifAB*-deletion mutant were injected in lettuce midribs, both strains grew vigorously and reached similar population densities in the inoculated section at 1, 2, and 3 days postinoculation (Supplementary Fig. S3). Furthermore, the effect of cichofactins on spread of *P. cichorii* in lettuce midribs was investigated (Supplementary Table S2). Both the wild-type strain and the cichofactin-deficient mutant moved to the adjacent midrib section (Supplementary Fig. S4) 3 days postinoculation but the mutant moved at a significantly slower rate than the wild type.

A successful attachment is the first step in colonization and pathogenesis (Danhorn and Fuqua 2007; Turnbull et al. 2001).

Hence, we wanted to understand whether cichofactins are involved in attachment of *P. cichorii* on three different regions of leaf surface. The wild-type strain and the cichofactin-deficient mutant attached equally well on the leaf surface but, for the petiole and center regions, very small but significant differences in bacterial attachment were observed.

Lettuce tissue inoculated with *P. cichorii* SF1-54 was analyzed using LC-ESI-MS to determine lipopeptide production of *P. cichorii* SF1-54 in planta. Cichofactins were not detected the first day after strain injection but the concentration of cichofactins gradually increased over the next 3 days to reach amounts of 1.2 to 8.1 $\mu\text{g/g}$ of leaf fresh weight in two different assays. No peaks corresponding to possible lipopeptide compounds were observed in uninfected lettuce tissue. In addition, no trace of the other lipopeptides could be detected (Fig. 6).

***P. cichorii* strains and isolates differ in swarming motility and cichofactin production.**

Belgian *P. cichorii* isolates from lettuce can be categorized into three subgroups based on colony morphology and BOX-polymerase chain reaction (PCR) patterns (Cottyn et al. 2009). Isolates belonging to the C3 group are significantly less virulent on lettuce than isolates of the C1 and C2 groups (Pauwelyn et al. 2011). The swarming ability of two *P. cichorii* isolates of each subgroup next to SF1-54 (C1 group)—the Japanese tar spot strain SPC9018 (C2 group), the Californian varnish spot strain 9D42 (related to the C3 group), the chrysanthemum strain NCPPB 907 (not related with any of the three groups), and the type strain NCPPB 943^T (not related with any of the three groups)—was investigated. *P. cichorii* isolates belonging to different groups differ in their swarming motility, with isolates from groups C1 and C2 swarming more rapidly than isolates of the C3 group (Fig. 7A). Surprisingly, the varnish spot

Fig. 6. Liquid chromatography electrospray ionization mass spectrometry detection of *Pseudomonas cichorii* SF1-54 lipopeptides in **A**, semipurified culture filtrate of *P. cichorii* SF1-54; **B**, extract of lettuce tissue infected with *P. cichorii* SF1-54; and **C**, extract of noninoculated lettuce tissue. Lipopeptides were extracted and analyzed 5 days postinoculation.

Fig. 7. A, Swarming behavior of *Pseudomonas cichorii* isolates of different groups as defined by Cottyn and associates (2009) on Luria-Bertani (LB) soft agar (0.5% [wt/vol]) plates. Plates were photographed 24 h after incubation. *Strain 9D42 is related to group C3. **B**, Quantification of cichofactin production by *P. cichorii* isolates of different groups on LB soft agar (0.5% [wt/vol]) plates. After 24 h of incubation, the bacterial cells were scraped from the swarming plates and suspended in 100% acetonitrile, and the optical density at 620 nm (OD₆₂₀) of each sample was measured as the indicator of biomass. Cichofactin production of each sample was measured by high-performance liquid chromatography electrospray ionization mass spectrometry by using single-ion recording integration of peaks corresponding to the exact mass of A and B homologues. Data are expressed as relative production based on the peak area per OD₆₂₀. Vertical bars indicate standard deviations. For each isolate, two plates were analyzed. No production of cichofactins was detected in the type strain of *P. cichorii* NCPPB 943^T.

strain 9D42, the type strain NCPPB 943^T and chrysanthemum strain NCPPB 907 did not show swarming motility. Interestingly, these three strains are weakly virulent or nonvirulent on lettuce (Pauwelyn et al. 2011) (Supplementary Table S3).

In addition, the level of cichofactin production in the various *P. cichorii* isolates was measured by HPLC-ESI-MS (Fig. 7B). Isolates of the C2 group secreted much more cichofactin than isolates of the C1 and C3 group. Very low amounts of cichofactins were produced by strains 9D42 and NCPPB 907, while no cichofactins could be detected in the type strain NCPPB 943^T. However, the presence of the *cif* genes could be detected in all strains with the primers (Table 3) used to construct the deletion plasmid pMQ30- Δ *cifAB* (data not shown). Our results show a correlation between cichofactin production and swarming and suggest, indirectly, an impact on virulence in lettuce.

DISCUSSION

Several plant-pathogenic *Pseudomonas* spp., such as *P. syringae* pv. *syringae*, *P. fuscovaginae*, *P. corrugata*, and *P. fluorescens* 5064, are known to produce (cyclic) lipopeptides (Bender et al. 1999; Catara 2007; Flamand et al. 1996; Hildebrand et al. 1998). Because these lipopeptides are surfactants, often exhibiting antimicrobial or phytotoxic activity, bioactivity assays are commonly used to assess their presence (Bender et al. 1999; Raaijmakers et al. 2010).

In this study, by means of bioassays, we showed that *P. cichorii* SF1-54 has surfactant, antimicrobial, and phytotoxic activity. These results confirm and extend the findings of Hu and associates (1998), who showed the production of an antibiotic compound by *P. cichorii*, and of Lazzaroni and associates (2003), mentioning the production of both phytotoxic and antibiotic compounds by *P. cichorii*. However, these compounds were not further characterized. Because the culture filtrate of *P. cichorii* SF1-54 showed characteristics similar to those of *P. cichorii* SF1-54 cells, bioactivity was due to compounds released by bacteria in the culture filtrate. Because tar symptoms caused by *P. cichorii* on crisphead lettuce in Japan are partly

due to apoptosis-like programmed cell death (Kiba et al. 2006, 2009), we suggest that phytotoxic compounds or virulence factors in the culture filtrate may be involved in the induction of apoptosis in lettuce.

An HPLC chromatogram of the semipurified extract from the culture filtrate of *P. cichorii* SF1-54 revealed that *P. cichorii* SF1-54 produces a complex mixture of several metabolites (Fig. 2). The HPLC elution pattern of the semipurified extract was similar to the elution pattern of the lipopeptide metabolites of *P. syringae* pv. *syringae* (Ballio et al. 1991) and *P. fuscovaginae* (Flamand et al. 1996), which is characterized by the presence of two groups of compounds with different polarity; more specifically, the lipodepsinonapeptides (such as syringomycins) and the more hydrophobic lipopeptides, such as syringopeptins and fuscopeptins. Preliminary data suggest that compounds B and C are pseudomycin like (Ballio et al. 1994), whereas compounds F and G are structurally related to corpeptin, a phytotoxic lipopeptide produced by *P. corrugata* (Emanuele et al. 1998), but this has to be confirmed by further chemical analysis.

HPLC fractionation of the culture filtrate of *P. cichorii* SF1-54 showed that the bioactivity of the strain could be ascribed to seven different compounds (Fig. 2). These compounds all showed surfactant activity, which is the primary characteristic of lipopeptide metabolites. However, bioactivities of these compounds are clearly different. These tests showed that two of them (D and E) were particularly interesting and their chemical structure was further investigated. Because of the close resemblance of these compounds to syringafactins (Berti et al. 2007), they were renamed cichofactins A and B. Their molecular masses are 1,109 and 1,137 Da, respectively. Their similar bioactivity and their molecular weight difference of 28 Da that corresponds to two methylene groups show that these compounds are homologs, only differing in the length of their fatty acid moieties. This relationship has been found with several other *Pseudomonas* lipopeptide homologs such as several syringopeptins (Ballio et al. 1991; Grgurina et al. 2002, 2005; Isogai et al. 1995). Up till now, cichofactins and syringafactins

Table 3. Strains, plasmids, and primers used in this study

Strain, plasmid, or primer	Genotype or description ^a	Source or reference
Strains		
<i>Pseudomonas cichorii</i>		
SF1-54, SF0068-02, SF0075-01, SF0057-3A1, AF0089-01, LMG8401, SF0125-01	Natural isolates from infected greenhouse butterhead lettuce in Belgium	Cottyn et al. 2011
9D42	Isolate from lettuce with varnish spot disease in California	Grogan et al. 1977
SPC9018	Isolate from lettuce with tar disease in Japan	Kiba et al. 2006
NCPPB 943 ^T	Type strain isolated from <i>Cichorium endivia</i> in Germany (=LMG2162 ^T)	Cottyn et al. 2009
NCPPB 907	Isolate from <i>Chrysanthemum manifoldum</i> in USA	Yamamoto et al. 2000
SF1-54NaIR	Spontaneous nalidixic acid-resistant mutant of SF1-54	This study
SF1-54- Δ <i>cifAB</i>	Cichofactin deficient-mutant of SF1-54 NaIR	This study
<i>Escherichia coli</i> S17-1	<i>thi pro hsdR recA::RP4-2- Tc^r::Mu Km^r::Tn7(λpir), Sm^r</i>	Simon et al. 1983
<i>Bacillus megaterium</i> LMG 7127	An indicator strain for lipopeptide	...
<i>Rhodolotula mucilaginosa</i> MUCL 30397	An indicator strain for lipopeptide	...
<i>Geotrichum candidum</i> MUCL 28959	An indicator strain for lipopeptide	...
<i>Saccharomyces cerevisiae</i> InvSc1	<i>MATa/MATa leu2/leu2 trp1-289/trp1-289 ura3-52/ura3-52 his3-Δ1/his3-Δ1</i>	Invitrogen
Plasmids		
pMQ30	7.6 kb mobilizable suicide vector used for gene replacements in <i>Pseudomonas</i> spp.: <i>SacB</i> , <i>URA3</i> , <i>Gm^r</i>	Shanks et al. 2006
pMQ30- <i>cifAB</i>	pMQ30 containing two ca. 1 kb fragments of the cichofactin biosynthesis genes to obtain a 9.7-kb deletion in these genes	This study
Primers (5'→3')		
NRPS14mut-F1	<i>GGAATTGTGAGCGGATAACAATTCACACAGGAAACAGCTGAGGTGCTG</i>	This study
NRPS14mut-R1	<i>GTGGACTTCAAT</i>	This study
NRPS14mut-F2	<i>ATCGAACTGACAGGCGTCT GTACCTGACCCGAGGTAGCA</i>	This study
NRPS14mut-R2	<i>TGCTACCTCGGGTCAGGTAC AGACGCCTGTCAAGTTTCGAT</i>	This study
	<i>CCAGGCAAAATCTGTTTTATCAGACCGCTTCTGCGTTCTGATTGTGTTTC</i>	This study
	<i>TGTCTCGCCTTG</i>	

^a Primer extensions are in italic font. Tc^r, Km^r, Sm^r, and Gm^r indicate resistant to tetracycline, kanamycin, streptomycin, and gentamycin, respectively.

(Berti et al. 2007) are the only members of the syringafactin family of linear *Pseudomonas*-produced lipopeptides. It has been suggested that the syringafactin NRPS system in *P. syringae* evolved from the arthrofactin (a cyclic lipoundecapeptide produced by *Pseudomonas* sp. MIS38) system, after which three modules of the arthrofactin NRPS were deleted, including the module that incorporates the threonyl residue that forms the ester linkage involved in cyclization. This may explain why syringafactins (and cichofactins) are linear (Berti et al. 2007; Roongsawang et al. 2011).

The genes encoding two nonribosomal peptide synthetases, *cifA* and *cifB*, were identified in the genome of *P. cichorii* SF1-54 as responsible for cichofactin biosynthesis. The putative peptide sequence of cichofactins predicted by *in silico* analysis of the A domain correlates perfectly with the peptide sequence determined by MS. In the flanking regions of cichofactin biosynthesis clusters, genes coding for ABC transporter systems and LuxR-type regulators have been found (Fig. 3) as previously identified in the lipopeptide biosynthesis genes of many *Pseudomonas* strains (Berti et al. 2007; de Bruijn and Raaijmakers, 2009; de Bruijn et al. 2008; Dubern et al. 2008; Kang and Gross 2005; Quigley et al. 1993; Roongsawang et al. 2003; Vallet-Gely et al. 2010). In *P. syringae* pv. *tomato* DC3000, only the LuxR-type transcriptional regulator upstream of the syringafactin biosynthesis genes appeared to be essential, whereas a mutation of the LuxR-type regulator located downstream did not affect the production of that compound (Berti et al. 2007).

Several biological roles for lipopeptides have been proposed, including their function in motility, antimicrobial activity, biofilm formation, and pathogenicity (D'aes et al. 2010; Raaijmakers et al. 2006, 2010). Little is known about functions of the lipopeptides belonging to the syringafactin family except for the involvement in swarming motility (Berti et al. 2007; Burch et al. 2010, 2012). To clarify the functions of cichofactins in *P. cichorii* SF1-54, a cichofactin-deficient mutant was constructed. This mutant does not produce cichofactins anymore but, surprisingly, an enhanced production of the phytotoxic compounds G and F was observed (Fig. 4). A similar phenomenon has been observed in *B. subtilis* 168 derivative strains which are able to synthesize lipopeptides after integration of a functional *sfp* gene (Coutte et al. 2010). Disruption of the plipastatin operon enhanced surfactin production. The redirection of common precursors, which needs ATP and cofactors, from the disrupted biosynthetic pathway to the other intact biosynthetic pathway of a lipopeptide may explain this phenomenon (Coutte et al. 2010).

Cichofactins were shown to be essential for surface motility in all *P. cichorii* isolates tested. This role has been established for many surfactants produced by pseudomonads such as syringafactin, entolysin, massetolide, viscosin, putisolvin, and arthrofactin (Berti et al. 2007; de Bruijn et al. 2007, 2008; Kruijt et al. 2009; Kuiper et al. 2004; Roongsawang et al. 2003; Vallet-Gely et al. 2010). Furthermore, microtiter plate assays showed that the cichofactin-deficient mutant formed significantly more biofilm than the wild-type strain *P. cichorii* SF1-54 (Fig. 5B). Several lipopeptides of *Pseudomonas* have been described to positively or negatively influence biofilm formation. For example, viscosin, massetolide, and CLP1 (sessilin) produced by *P. fluorescens* strains SBW25 and SS101 and *Pseudomonas* CMR12a, respectively (D'aes et al. 2010; de Bruijn et al. 2007, 2008), have a positive influence on biofilm formation, whereas putisolvins, arthrofactin, and CLP2 (orfamide) produced by *P. putida*, *Pseudomonas* sp. MIS38, and *Pseudomonas* CMR12a, respectively (D'aes et al. 2010; Kuiper et al. 2004; Roongsawang et al. 2003), negatively affect biofilm formation.

Cichofactins are also involved in pathogenicity of *P. cichorii* SF1-54 on lettuce but are not phytotoxic per se. Our data suggest that cichofactins are important for the in planta spread of *P. cichorii* because the cichofactin mutant strain spread in lettuce midribs at a significantly slower rate than the wild type. Both cichofactins A and B were detected in infected plant tissue (Fig. 6), indicating their importance in infection or pathogenesis of lettuce leaves by *P. cichorii*. In contrast, the production of the other compounds in infected leaf tissues by *P. cichorii* SF1-54 remains unclear. They may be produced in very low amounts compared with cichofactins and, thus, in insufficient amounts for adequate detection using HPLC-ESI-MS. This latter hypothesis should not be surprising because syringomycins were detected in infected leaf tissue only in minute amounts by repeated fractionations using HPLC (Grgurina et al. 2002). Another possibility is that they are produced at earlier time points and degrade to low amounts or interact with the plasma membrane or other components of the plant cell (Fogliano et al. 1999; Henry et al. 2011), which may thwart chemical detection.

Interestingly, *P. cichorii* isolates that have low virulence on lettuce produce less cichofactin than more pathogenic isolates (Fig. 7B). It remains to be investigated whether these isolates also differ in their production of the phytotoxic compounds F and G.

In conclusion, this study showed that *P. cichorii* SF1-54 produces seven bioactive compounds with lipopeptide characteristics, and two new lipopeptides which we focused on were identified as isomers and named cichofactins. Our results provide evidence that cichofactins are responsible for swarming motility and are involved in virulence of *P. cichorii* SF1-54 on lettuce. We are currently investigating the role of compounds F and G in virulence and bioactivity of *P. cichorii*.

MATERIALS AND METHODS

Microorganisms and culture conditions.

Microorganisms used in this study are listed in Table 3. *P. cichorii* isolates and mutants and *Escherichia coli* strains were kept at -80°C in LB broth (Sambrook and Russel 2001) with 20% glycerol for long-term storage. *R. mucilaginosa* MUCL 30397, *G. candidum* MUCL 28959, and *B. megaterium* LMG 7127 were used as indicator organisms for bioassays of antimicrobial activities. *Bacillus* and *Rhodotorula* strains were maintained at -80°C in LB broth with 20% glycerol, and *G. candidum* on potato dextrose agar (PDA; Difco, Erembodegem, Belgium) slants. *Pseudomonas* strains were routinely grown on Pseudomonas Agar F (PAF; Difco) or SRM_{AF} medium (Gross 1985). *B. megaterium* and *E. coli* strains were grown on LB medium at 28 and 37°C , respectively. *R. mucilaginosa* and *G. candidum* were cultured on PDA at 28°C . *Saccharomyces cerevisiae* was grown on yeast-extract-peptone-dextrose medium at 30°C (Shanks et al. 2006). Antibiotics were added to the growth media (if required) at the following concentrations: gentamycin at 100 $\mu\text{g/ml}$ and nalidixic acid at 10 $\mu\text{g/ml}$.

Biological activity assays.

Biosurfactant production was tested using the drop collapse assay, as described by Kuiper and associates (2004). Briefly, cells from a single colony, previously grown on PAF-medium for 24 h, were transferred by a toothpick and suspended into a droplet of 50 μl of sterile water on a hydrophobic background (more specifically, parafilm 'M' laboratory film). When crude or semipurified culture filtrate or purified compound was tested, 5 μl of the material was added to a droplet of 50 μl of H_2O and placed onto parafilm. The reduction in surface tension was visualized by observation of the spread of the droplet over the parafilm surface.

Biological activity was assayed with *P. cichorii* SF1-54 suspension (1.0×10^8 CFU/ml in 50 mM phosphate buffer [PB], pH 7.0), crude culture filtrate, semipurified culture filtrate, and purified compounds.

Phytotoxic activity was tested on chicory or lettuce leaves by injection of 300 μ l of bacterial suspension or culture filtrate with a syringe through the abaxial surface. The infiltrated leaves were incubated at 100% humidity and 25°C for 24 h and appearance of necrotic lesions was scored at 24 h after injection. PB, SRM_{AF}, or methanol were used as negative control treatments.

Antimicrobial activity was tested against *B. megaterium*, *G. candidum*, and *R. mucilaginosa* using the protocol described by Schaad and associates (2001), with minor modifications. Briefly, a bacterial suspension of *P. cichorii* SF1-54 (1×10^8 CFU/ml) was spotted on PDA plates in triplicate, allowing growth at 25°C for 5 days. The areas of colony growth were marked and the colonies removed with a sterile swab. The plates were exposed to chloroform vapors for 20 min to kill the remaining bacterial cells, followed by dissipation of the vapors for an additional 40 min. When culture filtrate or purified compound was tested, 50 or 10 μ l was spotted onto PDA plates (for *G. candidum* and *R. mucilaginosa*) or LB plates (for *B. megaterium*), respectively, and followed by drying of the drops. The plate was then sprayed with a spore/cell suspension containing approximately 10^6 cells or spores per milliliter of the indicator organism. After 24 to 48 h of incubation at 28°C, inhibition of the indicator organism was scored.

Lipopeptide production, extraction, and purification.

P. cichorii SF1-54 was grown in 100-ml Erlenmeyer flasks containing 30 ml of SRM_{AF} medium. After incubation for 6 days at 25°C without shaking, the *P. cichorii* SF1-54 culture was centrifuged ($10,000 \times g$, 10 min) (Sigma 4K15; Sartorius, Göttingen, Germany) and the supernatant was filter sterilized through a 0.2- μ m filter (Millipore; Sartorius) to obtain the crude culture filtrate. Before use, 100 μ l of this crude culture filtrate was plated on PAF medium in three replications to confirm its sterility. The crude culture filtrate was loaded on C₁₈ solid-phase extraction cartridges (C18 SPE Maxi-Clean cartridges; Alltech, Grace, Lokeren, Belgium) and washed with 5% methanol, and lipopeptides were desorbed with 100 ml of 100% methanol to obtain the semipurified culture filtrate.

To purify the *P. cichorii* SF1-54 metabolites, 200 ml of semipurified culture filtrate was further concentrated using rotavaporation (Rotavapor R-2000; Büchi (Essen, Germany) Flawil, Switzerland) and vacuum speed (Savant Speed-Vac Plus SC110A; Thermo Fisher Scientific, Bremen, Germany) and separated with an Agilent 1100 series HPLC (Hewlett Packard, Diegem, Belgium) on a protein and peptide Vydac C18 column (5 μ m, 4.6 by 250 mm; Grace, Lokeren, Belgium). Metabolites of *P. cichorii* SF1-54 were eluted with a gradient of acetonitrile acidified with 0.1% formic acid in water acidified with 0.1% formic acid at a constant flow rate of 1 ml min⁻¹ and at room temperature. The UV signal was monitored at 220 nm. All different fractions were collected, concentrated using rotavaporation and vacuum speed, and used for bioactivity assays.

Chemical analysis of lipopeptides produced by *P. cichorii* SF1-54.

Amino acid analysis. For the amino acid analysis, HPLC-purified compounds were collected in glass tubes and dried using vacuum speed (Savant Speed-Vac Plus SC110A). The samples were hydrolyzed at 115°C for 6 h with 1.5 ml of 6 N HCl. After drying in a Speedvac (Savant Speed-Vac Plus SC110A) and dissolving in 100 μ l of 0.1 N HCl, the samples were analyzed on a ZORBAX Eclipse AAA column (Agilent,

Diegem, Belgium) using an Agilent 1100 series HPLC (Hewlett Packard) by strictly following the instructions provided by the Agilent manufacturer. Briefly, the hydrolyzed samples and amino acid standard solutions were automatically derivatized in the autosampler with o-phthalaldehyde (OPA) for most amino acids and 9-fluorenylmethyl-chloroformate (FMOC) for lysine, hydroxyproline, and proline residues. After derivatization, 0.5 μ l of each sample was injected with a flow rate 2 ml/min and detected at 338 nm (OPA) and 262 nm (FMOC). Amino acids in the lipopeptide hydrolysis products were identified and their relative amounts in the samples were estimated based on the amino acid standards analyzed simultaneously.

LC-ESI-MS. Analysis of lipopeptides produced by *P. cichorii* was carried out with LC-ESI-MS (RP-HPLC Waters Alliance 2695/diode array detector, in tandem with a Single Quad Waters SQD Mass spectrometer; Waters, Zellik, Belgium) on an XTerra MS C18 column (3.5 μ m, 4.6 by 150 mm) (Waters). Putative lipopeptides of *P. cichorii* SF1-54 were eluted with a gradient of acetonitrile acidified with 0.1% formic acid in water acidified with 0.1% formic acid at a constant flow rate 0.5 ml/min at 40°C. Compounds were monitored in the positive ion mode and in-source settings in the SQD were as follows: source temperature, 130°C; desolvation temperature, 280°C; nitrogen flow, 600 liters/h; cone voltage, 100 V. A similar setting was used for lipopeptide fragmentation, except the cone voltage was 75 V instead of 100 V. Both instruments were controlled with the MassLynx software (Waters).

HR-ESI-MS-MS. All measurements were conducted at the MS lab of the Institute of Organic Chemistry at the University of Cologne on a LTQ-Orbitrap XL instrument (Thermo Fisher Scientific), with a conventional ESI ion source (ESI-voltage, 3.5 kV; capillary temperature, 275°C; and flow rate, 5 μ l/min). Acidified methanol/water solutions were used in micromolar concentrations. MSⁿ-product ion spectra were produced in a linear ion trap (LTQ part of the Orbitrap) or in an octopole collision cell. The precursor ions were selected and activated until extensive fragmentation occurred.

Identification and bioinformatic analysis of the NRPS genes.

Whole-genome sequencing of *P. cichorii* SF1-54 was performed at Genoscreen, France, with a Roche 455 Gs-FLX sequencer. The draft genome was automatically annotated using the RAST server (Aziz et al. 2008). Putative NRPS encoding regions were found by comparison of the genome sequence of *P. cichorii* SF1-54 with the genomes of *P. syringae* pv. *tomato* DC3000 and *P. syringae* pv. *syringae* B748a (Buell et al. 2003; Feil et al. 2005) using the program MAUVE (Darling et al. 2010). Gap filling in the putative NRPS-encoding gene regions was performed by way of PCR and Sanger sequencing of the PCR product (Agowa Sequencing, Berlin). ORF were identified using the program Glimmer (Delcher et al. 1999), available on the National Center for Biotechnology Information website. The catalytic domains present in the NRPS genes were identified using the NRPS/PKS analysis website (Bachmann and Ravel 2009) and the NRPS-PKS website (Ansari et al. 2004). The specificity prediction of the A domain was conducted by using the web-based software NRPSpredictor2 (Rausch et al. 2005; Rottig et al. 2011). The D/L configurations of amino acid residues of nonribosomal peptide synthetases were investigated by phylogenetic analysis of the C domains via the Phylogeny.fr website (Dereeper et al. 2008). Genes adjacent to the putative NRPS genes were compared with the sequences available in GenBank using the program BlastX. Multiple alignments and phylogeny tree design were performed using the ClustalW2 tool available at the EBI site.

Construction of the cichofactin-deficient

P. cichorii SF1-54 mutant.

A cichofactin-deficient mutant of *P. cichorii* SF1-54NaI^R was constructed by means of an in vivo cloning technique with the yeast *S. cerevisiae* InvSc1 (Hoang et al. 1998; Shanks et al. 2006). To construct deletion plasmid pMQ30- Δ cifAB, primers NRPS14mut-F1 and NRPS14mut-R1 were used to amplify the upstream fragment and primers NRPS14mut-F2 and NRPS14mut-R2 were used for the downstream fragment (Table 3). The cichofactin-deficient mutant contained a deletion of 9,715 bp in the cichofactin-encoding genes *cifA* and *cifB*. Deletion of the NRPS fragment was confirmed by PCR and the mutant was characterized phenotypically.

Biological roles of cichofactin in *P. cichorii*.

Surface motility of *P. cichorii* isolates was assayed on soft LB plates (0.5% agar). Bacteria of a fully grown colony on PAF-medium were applied in the center of the soft agar plates with a sterile toothpick and the plates were subsequently incubated at 28°C up to 72 h after inoculation.

Biofilm formation was tested in polystyrene 96-well plates based on protocols described by Merrit and associates (2005) and Maddula and associates (2006), with minor modifications. Briefly, strains were grown overnight in LB broth. The 96-well flat-bottom polystyrene microtiter plates (Greiner bio-one, Wemmel, Belgium) containing 120 μ l of King's B broth (King et al. 1954) were inoculated using 100-fold dilutions of these culture, previously adjusted to an optical density at 620 nm (OD₆₂₀) [defined] of 0.8. Plates were incubated at 21°C for 24 h without shaking and final cell densities were determined (OD₆₂₀). Subsequently, the medium was removed from the microtiter plates, rinsed with water, filled with 150 μ l of 0.1% crystal violet, and incubated for 10 min at room temperature. Excess of stain was removed by inversion of the plate followed by two washings with tap water. When the plates were air dried, adherent cells were decolorized with 95% ethanol solution for 10 min and transferred to new microtiter plates in order to measure the absorbance at 595 nm. The experiment was carried out three times, with seven repetitions per treatment. Statistical analysis was performed in SPSS 17.0 (SPSS Inc., Chicago) for Windows using nonparametric Mann-Whitney U tests ($P = 0.05$).

Pathogenicity of the cichofactin-deficient mutant and of the *P. cichorii* strains NCPB 907 and NCPB 943^T was tested at Inagro, Rumbeke, Belgium, according to the methods described in a previous study (Pauwelyn et al. 2011). The strains were inoculated on butterhead lettuce 'Flandria' plants (Rijk Zwaan, De Lier, The Netherlands) at head formation. Tap water without bacteria was used as the noninoculated treatment. Each treatment was assayed in three replicates with 24 plants per replicate and the experiment was carried out twice. At harvest, disease severity was assessed by giving each plant a score of 0 to 4, where 0 = healthy plant; 1 = little black spots on the leaf periphery of the inner crop leaves; 2 = infection of small side ribs or black spots or stripes on the midrib; 3 = one, two, or three rotten midribs; and 4 = four or more rotten midribs. In addition, the number of rotten midribs on each plant was scored. Data were analyzed using nonparametric Kruskal-Wallis and Mann-Whitney comparisons ($P = 0.05$).

To test the ability of *P. cichorii* SF1-54- Δ cifAB to grow in lettuce midribs, the midrib of the inner lettuce leaves was injected with 0.2 ml of bacterial suspension (1×10^5 CFU/ml). The injected leaves were incubated at 100% humidity at 25°C. The inoculated sections (2 cm²) were excised from the midribs of three lettuce leaves at 0, 1, 2, and 3 days postinoculation and macerated in 2 ml of sterile 0.05 M potassium PB (pH 7.0). Tenfold serial dilutions from the macerate were prepared

with PB and 100- μ l aliquots of each dilution were spread on three PAF plates with naladixic acid at 50 μ g/ml. Colonies were counted 2 days after incubation at 28°C to estimate population densities. For assessment of bacterial spread in midribs, the adjacent sections (2 cm²) were also excised from the midribs 3 days postinoculation to estimate population densities as described above. The experiment was repeated twice using three leaves from three different plants per treatment.

Assays for bacterial attachment on lettuce leaves were performed according to the method of Kroupitski and associates (2011), with slight modification. Five leaf discs from different regions (near petiole, center, and near blade) from five different butterhead lettuce leaves were cut using an aseptic cork borer (1.4 cm in diameter) and the entire experiment was repeated once. The leaf discs were immersed in 10-ml of a bacterial suspension in a 50-ml falcon tube and incubated at 25°C for 2 h without shaking. The leaf discs were washed twice for 1 min in 10 ml of sterile water and macerated in 1 ml of sterile 0.05 M potassium PB (pH 7.0) for estimation of bacterial population sizes as described before.

All data were statistically analyzed using the software package SPSS 17.0 for Windows. Because the data did not meet the conditions of normality and homogeneity of variance, nonparametric Kruskal-Wallis and Mann-Whitney comparisons ($P = 0.05$) were performed.

In planta detection

of lipopeptides produced by *P. cichorii* SF1-54.

A bacterial suspension of *P. cichorii* SF1-54 (1×10^6 CFU/ml) was inoculated in the midrib of lettuce leaves. The lettuce leaves were placed on upside-down petri dishes into plastic infection trays filled with wetted sterile cotton. The infection trays were covered with a plastic lid to obtain high relative humidity and incubated at room temperature for 5 days. At 1, 2, 3, 4, and 5 days postinoculation, symptomatic tissue was excised from the lettuce leaves and ground in liquid nitrogen. After thawing, 10 ml of ground symptomatic leaf tissue was suspended in 40 ml of 95% acetonitrile acidified with 0.1% formic acid, shaken for 1 h at room temperature, and then centrifuged (5 min at 10,000 rpm; Beckman avanti J-25i). The pellet was dissolved again in 10 ml of 95% of acetonitrile acidified with 0.1% formic acid and centrifuged. Supernatants of both extractions were mixed, concentrated five times using rotavaporation, and analyzed with LC-ESI-MS.

ACKNOWLEDGMENTS

M. Ongena is Research Associate at the F.R.S.-FNRS in Belgium. This work was financed by the "Fonds voor Wetenschappelijk Onderzoek-Vlaanderen" (grant number G.0002.10N) and by the INTERREG IV program France-Wallonie-Vlaanderen (Phytobio project). We thank L. Franzil (Gembloux Agro-Bio Tech) and I. Delaere (Ghent University) for technical assistance during the experimental work.

LITERATURE CITED

- Ansari, M. Z., Yadav, G., Gokhale, R. S., and Mohanty, D. 2004. NRPS-PKS: a knowledge-based resource for analysis of NRPS/PKS megasynthases. *Nucleic Acids Res.* 32:405-413.
- Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., Formsma, K., Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman, A. L., Overbeek, R. A., McNeil, L. K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G. D., Reich, C., Stevens, R., Vassieva, O., Vonstein, V., Wilke, A., and Zagnitko, O. 2008. The RAST server: Rapid annotations using subsystems technology. *BMC Genomics* 9:75.
- Bachmann, B. O., and Ravel, J. 2009. Methods for *In silico* prediction of microbial secondary metabolic pathways from DNA sequence data. *Methods Enzymol.* 458:181-217.
- Balibar, C. J., Vaillancourt, F. H., and Walsh, C. T. 2005. Generation of D

- amino acid residues in assembly of arthrofactors by dual condensation/epimerization domains. *Chem. Biol.* 12:1189-1200.
- Ballio, A., Barra, D., Bossa, F., Collina, A., Grgurina, I., Marino, G., Monetti, G., Paci, M., Pucci, P., Segre, A., and Simmaco, M. 1991. Syringopeptins, new phytotoxic lipodepsipeptides of *Pseudomonas syringae* pv. *syringae*. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 291:109-112.
- Ballio, A., Bossa, F., Di Giorgio, D., Ferranti, P., Paci, M., Pucci, P., Scaloni, A., Segre, A., and Strobel, G.A. 1994. Novel bioactive lipodepsipeptides from *Pseudomonas syringae*—The pseudomycins. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 355:96-100.
- Bender, C. L., and Scholz-Schroeder, B. K. 2004. New insights into the biosynthesis, mode of action, and regulation of syringomycin, syringopeptin and coronatine. Kluwer Academic Press, Dordrecht, The Netherlands.
- Bender, C. L., Alarcon-Chaidez, F., and Gross, D. C. 1999. *Pseudomonas syringae* phytotoxins: Mode of action, regulation, and biosynthesis by peptide and polyketide synthetases. *Microbiol. Mol. Biol. Rev.* 63:266-292.
- Berti, A. D., Greve, N. J., Christensen, Q. H., and Thomas, M. G. 2007. Identification of a biosynthetic gene cluster and the six associated lipopeptides involved in swarming motility of *Pseudomonas syringae* pv. *tomato* DC3000. *J. Bacteriol.* 189:6312-6323.
- Bradbury, J. F. 1986. Guide to Plant Pathogenic Bacteria. Oxford University Press, New York.
- Buell, C. R., Joardar, V., Lindeberg, M., Selengut, J., Paulsen, I. T., Gwinn, M. L., Dodson, R. J., Deboy, R. T., Durkin, A. S., Kolonay, J. F., Madupu, R., Daugherty, S., Brinkac, L., Beanan, M. J., Haft, D. H., Nelson, W. C., Davidsen, T., Zafar, N., Zhou, L. W., Liu, J., Yuan, Q. P., Khouri, H., Fedorova, N., Tran, B., Russell, D., Berry, K., Utterback, T., Van Aken, S. E., Feldblyum, T. V., D'Alessandro, M., Deng, W. L., Ramos, A. R., Alfano, J. R., Cartinhour, S., Chatterjee, A. K., Delaney, T. P., Lazarowitz, S. G., Martin, G. B., Schneider, D. J., Tang, X. Y., Bender, C. L., White, O., Fraser, C. M., and Collmer, A. 2003. The complete genome sequence of the Arabidopsis and tomato pathogen *Pseudomonas syringae* pv. *tomato* DC3000. *Proc. Natl. Acad. Sci. U.S.A.* 100:10181-10186.
- Burch, A. Y., Shimada, B. K., Browne, P. J., and Lindow, S. E. 2010. Novel high-throughput detection method to assess bacterial surfactant production. *Appl. Environ. Microbiol.* 76:5363-5372.
- Burch, A. Y., Shimada, B. K., Mullin, S. W. A., Dunlap, C. A., Bowman, M. J., and Lindow, S. E. 2012. *Pseudomonas syringae* coordinates production of a motility-enabling surfactant with flagellar assembly. *J. Bacteriol.* 194:1287-1298.
- Catara, V. 2007. *Pseudomonas corrugata*: Plant pathogen and/or biological resource? *Mol. Plant Pathol.* 8:233-244.
- Challis, G. L., Ravel, J., and Townsend, C. A. 2000. Predictive, structure-based model of amino acid recognition by nonribosomal peptide synthetase adenylation domains. *Chem. Biol.* 7:211-224.
- Cottyn, B., Heylen, K., Heyrman, J., Vanhouteghem, K., Pauwelyn, E., Bleyaert, P., Van Vaerenbergh, J., Höfte, M., De Vos, P., and Maes, M. 2009. *Pseudomonas cichorii* as the causal agent of midrib rot, an emerging disease of greenhouse-grown butterhead lettuce in Flanders. *Syst. Appl. Microbiol.* 32:211-225.
- Cottyn, B., Baeyen, S., Pauwelyn, E., Verbaednert, I., De Vos, P., Bleyaert, P., Höfte, M., and Maes, M. 2011. Development of a real-time PCR assay for *Pseudomonas cichorii*, the causal agent of midrib rot in greenhouse-grown lettuce, and its detection in irrigating water. *Plant Pathol.* 60:453-461.
- Coutte, F., Leclère, V., Bechet, M., Guez, J. S., Lecouturier, D., Chollet-Imbert, M., Dhulster, P., and Jacques, P. 2010. Effect of *pps* disruption and constitutive expression of *srfA* on surfactin productivity, spreading and antagonistic properties of *Bacillus subtilis* 168 derivatives. *J. Appl. Microbiol.* 109:480-491.
- D'aes, J., De Maeyer, K., Pauwelyn, E., and Höfte, M. 2010. Biosurfactants in plant-*Pseudomonas* interactions and their importance to biocontrol. *Environ. Microbiol. Rep.* 2:359-372.
- Danhorn, T., and Fuqua, C. 2007. Biofilm formation by plant-associated bacteria. *Annu. Rev. Microbiol.* 61:401-422.
- Darling, A. E., Mau, B., and Perna, N. T. 2010. progressiveMauve: Multiple genome alignment with gene gain, loss and rearrangement. *PloS One* 5:e11147. Published online.
- de Bruijn, I., and Raaijmakers, J. M. 2009. Diversity and functional analysis of LuxR-type transcriptional regulators of cyclic lipopeptide biosynthesis in *Pseudomonas fluorescens*. *Appl. Environ. Microbiol.* 75:4753-4761.
- de Bruijn, I., de Kock, M. J. D., Yang, M., de Waard, P., van Beek, T. A., and Raaijmakers, J. M. 2007. Genome-based discovery, structure prediction and functional analysis of cyclic lipopeptide antibiotics in *Pseudomonas* species. *Mol. Microbiol.* 63:417-428.
- de Bruijn, I., de Kock, M. J. D., de Waard, P., van Beek, T. A., and Raaijmakers, J. M. 2008. Massetolide a biosynthesis in *Pseudomonas fluorescens*. *J. Bacteriol.* 190:2777-2789.
- Delcher, A. L., Harmon, D., Kasif, S., White, O., and Salzberg, S. L. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res.* 27:4636-4641.
- Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J. F., Guindon, S., Lefort, V., Lescot, M., Claverie, J. M., and Gascuel, O. 2008. Phylogeny.fr: Robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res.* 36 (Suppl. 2):W465-W469.
- Dubert, J. F., Coppoolse, E. R., Stiekema, W. J., and Bloemberg, G. V. 2008. Genetic and functional characterization of the gene cluster directing the biosynthesis of putisolvin I and II in *Pseudomonas putida* strain PCL1445. *Microbiology* 154:2070-2083.
- Emanuele, M. C., Scaloni, A., Lavermicocca, P., Jacobellis, N. S., Camoni, L., Di Giorgio, D., Pucci, P., Paci, M., Segre, A., and Ballio, A. 1998. Corpeptins, new bioactive lipodepsipeptides from cultures of *Pseudomonas corrugata*. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 433:317-320.
- Feil, H., Feil, W. S., Chain, P., Larimer, F., DiBartolo, G., Copeland, A., Lykidis, A., Trong, S., Nolan, M., Goltsman, E., Thiel, J., Malfatti, S., Loper, J. E., Lapidus, A., Detter, J. C., Land, M., Richardson, P. M., Kyrpides, N. C., Ivanova, N., and Lindow, S. E. 2005. Comparison of the complete genome sequences of *Pseudomonas syringae* pv. *syringae* B728a and pv. *tomato* DC3000. *Proc. Natl. Acad. Sci. U.S.A.* 102:11064-11069.
- Flamand, M. C., Pelsser, S., Ewbank, E., and Maraite, H. 1996. Production of syringotoxin and other bioactive peptides by *Pseudomonas fuscovaginae*. *Physiol. Mol. Plant Pathol.* 48:217-231.
- Fogliano, V., Gallo, M., Vinale, F., Ritieni, A., Randazzo, G., Greco, M., Lops, R., and Graniti, A. 1999. Immunological detection of syringopeptins produced by *Pseudomonas syringae* pv. *lachrymans*. *Physiol. Mol. Plant Pathol.* 55:255-261.
- Fuchs, R., and Buzikiewicz, H. 2001. Rearrangement reactions in the electrospray ionization mass spectrometry of pyoverdins. *Int. J. Mass Spectrom.* 210/211:603-612.
- Grgurina, I., Mariotti, F., Fogliano, V., Gallo, M., Scaloni, A., Iacobellis, N. S., Lo Cantore, P., Mannina, L., Castelli, V. V., Greco, M. L., and Graniti, A. 2002. A new syringopeptin produced by bean strains of *Pseudomonas syringae* pv. *syringae*. *Biochim. Biophys. Acta* 1597:81-89.
- Grgurina, I., Bensaci, M., Pocsfalvi, G., Mannina, L., Cruciani, O., Fiore, A., Fogliano, V., Sorensen, K. N., and Takemoto, J. Y. 2005. Novel cyclic lipodepsipeptide from *Pseudomonas syringae* pv. *lachrymans* strain 508 and syringopeptin antimicrobial activities. *Antimicrob. Agents Chemother.* 49:5037-5045.
- Grogan, R. G., Misaghi, I. J., Kimble, K. A., Greathead, A. S., Ririe, D., and Bardin, R. 1977. Varnish spot, destructive disease of lettuce in California caused by *Pseudomonas cichorii*. *Phytopathology* 67:957-960.
- Gross, D. C. 1985. Regulation of syringomycin synthesis in *Pseudomonas syringae* pv. *syringae* and defined conditions for its production. *J. Appl. Bacteriol.* 58:167-174.
- Henry, G., Deleu, M., Jourdan, E., Thonart, P., and Ongena, M. 2011. The bacterial lipopeptide surfactin targets the lipid fraction of the plant plasma membrane to trigger immune-related defense responses. *Cell. Microbiol.* 13:1824-1837.
- Hikichi, Y., Suzuki, K., Toyoda, K., Horikoshi, M., Hirooka, T., and Okuno, T. 1998. Successive observation of growth and movement of genetically lux-marked *Pseudomonas cichorii* and the response of host tissues in the same lettuce leaf. *Ann. Phytopathol. Soc. Jpn.* 64:519-525.
- Hildebrand, P. D., Braun, P. G., McRae, K. B., and Lu, X. 1998. Role of the biosurfactant viscosin in broccoli head rot caused by a pectolytic strain of *Pseudomonas fluorescens*. *Can. J. Plant Pathol.* 20:296-303.
- Hoang, T. T., Karkhoff-Schwiezer, R. R., Kutchna, A. J., and Schweizer, H. P. 1998. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: Application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* 212:77-86.
- Hojo, H., Koyanagi, M., Tanaka, M., Kajihara, S., Ohnishi, K., Kiba, A., and Hikichi, Y. 2008. The *hrp* genes of *Pseudomonas cichorii* are essential for pathogenicity on eggplant but not on lettuce. *Microbiology* 154:2920-2928.
- Hu, F. P., Young, J. M., and Fletcher, M. J. 1998. Preliminary description of biocidal (syringomycin) activity in fluorescent plant pathogenic *Pseudomonas* species. *J. Appl. Microbiol.* 85:365-371.
- Hutchison, M. L., and Gross, D. C. 1997. Lipopeptide phytotoxins produced by *Pseudomonas syringae* pv. *syringae*: Comparison of the biosurfactant and ion channel-forming activities of syringopeptin and syringomycin. *Mol. Plant-Microbe Interact.* 10:347-354.
- Hutchison, M. L., Tester, M. A., and Gross, D. C. 1995. Role of biosurfactant and ion channel-forming activities of syringomycin in transmem-

- brane ion flux—A model for the mechanism of action in the plant-pathogen interaction. *Mol. Plant-Microbe Interact.* 8:610-620.
- Isogai, A., Iguchi, H., Nakayama, J., Kusai, A., Takemoto, J. Y., and Suzuki, A. 1995. Structural analysis of new syringopeptins by tandem mass-spectrometry. *Biosci. Biotechnol. Biochem.* 59:1374-1376.
- Jagger, I. C. 1914. A bacterial leaf spot disease of celery. (Abstr.) *Phytopathology* 4:395.
- Jones, J. B., Raju, B. C., and Engelhard, A. W. 1984. Effects of temperature and leaf wetness on development of bacterial spot of geranium and chrysanthemum incited by *Pseudomonas cichorii*. *Plant Dis.* 68:248-251.
- Kajihara, S., Hojo, H., Koyanagi, M., Tanaka, M., Mizumoto, H., Ohnishi, K., Kiba, A., and Hikichi, Y. 2012. Implication of *hrpW* in virulence of *Pseudomonas cichorii*. *Plant Pathol.* 61:355-363.
- Kang, H. J., and Gross, D. C. 2005. Characterization of a resistance-nodulation-cell division transporter system associated with the *syd-syp* genomic island of *Pseudomonas syringae* pv. *syringae*. *Appl. Environ. Microbiol.* 71:5056-5065.
- Kiba, A., Sangawa, Y., Ohnishi, K., Yao, N., Park, P., Nakayashiki, H., Tosa, Y., Mayama, S., and Hikichi, Y. 2006. Induction of apoptotic cell death leads to the development of bacterial rot caused by *Pseudomonas cichorii*. *Mol. Plant-Microbe Interact.* 19:112-122.
- Kiba, A., Lee, K. Y., Ohnishi, K., Park, P., Nakayashiki, H., Tosa, Y., Mayama, S., and Hikichi, Y. 2009. Induction of reactive oxygen generation and functional changes in mitochondria and their involvement in the development of bacterial rot in lettuce caused by *Pseudomonas cichorii*. *Physiol. Mol. Plant Pathol.* 74:45-54.
- King, E. O., Ward, M. K., and Raney, D. E. 1954. Two simple media for the demonstration of pyocyanin and fluorescein. *J. Lab Clin. Med.* 44:301-307.
- Kroupitski, Y., Pinto, R., Belausov, E., and Sela, S. 2011. Distribution of *Salmonella typhimurium* in romaine lettuce leaves. *Food Microbiol.* 28:990-997.
- Kruijt, M., Tran, H., and Raaijmakers, J. M. 2009. Functional, genetic and chemical characterization of biosurfactants produced by plant growth-promoting *Pseudomonas putida* 267. *J. Appl. Microbiol.* 107:546-556.
- Kuiper, I., Lagendijk, E. L., Pickford, R., Derrick, J. P., Lamers, G. E. M., Thomas-Oates, J. E., Lugtenberg, B. J. J., and Bloemberg, G. V. 2004. Characterization of two *Pseudomonas putida* lipopeptide biosurfactants, putisolvin I and II, which inhibit biofilm formation and break down existing biofilms. *Mol. Microbiol.* 51:97-113.
- Kunkel, B. N., and Zhongying, C. 2006. *Virulence of Plant Pathogenic Bacteria*. Springer, New York.
- Lavermicocca, P., Iacobellis, N. S., Simmaco, M., and Graniti, A. 1997. Biological properties and spectrum of activity of *Pseudomonas syringae* pv. *syringae* toxins. *Physiol. Mol. Plant Pathol.* 50:129-140.
- Lazzaroni, S., Evidente, A., and Surico, G. 2003. *Toxic metabolites and lipopolysaccharides from Pseudomonas cichorii*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Maddula, V., Zhang, Z., Pierson, E. A., and Pierson, L. S. 2006. Quorum sensing and phenazines are involved in biofilm formation by *Pseudomonas chlororaphis* (aureofaciens) strain 30-84. *Microb. Ecol.* 52:289-301.
- Merritt, J. H., Kadouri, D. E., and O'Toole, G. A. 2005. Growing and analyzing static biofilms. Pages 1B.1.1-1B.1.17 in: *Current Protocols in Microbiology*. R. Coico, T. Kowalik, J. Quarles, B. Stevenson, and R. Taylor, eds. John Wiley & Sons, Chichester, U.K.
- Pauwelyn, E., Vanhouteghem, K., Cottyn, B., De Vos, P., Maes, M., Bleyaert, P., and Höfte, M. 2011. Epidemiology of *Pseudomonas cichorii*, the cause of lettuce midrib rot. *J. Phytopathol.* 159:298-305.
- Quigley, N. B., Mo, Y. Y., and Gross, D. C. 1993. *SyrD* is required for syringomycin production by *Pseudomonas syringae* pathovar *syringae* and is related to a family of ATP-binding secretion proteins. *Mol. Microbiol.* 9:787-801.
- Raaijmakers, J. M., de Bruijn, I., and de Kock, M. J. D. 2006. Cyclic lipopeptide production by plant-associated *Pseudomonas* spp.: Diversity, activity, biosynthesis, and regulation. *Mol. Plant-Microbe Interact.* 19:699-710.
- Raaijmakers, J. M., de Bruijn, I., Nybroe, O., and Ongena, M. 2010. Natural functions of lipopeptides from *Bacillus* and *Pseudomonas*: More than surfactants and antibiotics. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Rev.* 34:1037-1062.
- Rausch, C., Weber, T., Kohlbacher, O., Wohlleben, W., and Huson, D. H. 2005. Specificity prediction of adenylation domains in nonribosomal peptide synthetases (NRPS) using transductive support vector machines (TSVM). *Nucleic Acids Res.* 33:5799-5808.
- Rausch, C., Hoof, I., Weber, T., Wohlleben, W., and Huson, D. H. 2007. Phylogenetic analysis of condensation domains in NRPS sheds light on their functional evolution. *BMC Evol. Biol.* 7:78.
- Roongsawang, N., Hase, K., Haruki, M., Imanaka, T., Morikawa, M., and Kanaya, S. 2003. Cloning and characterization of the gene cluster encoding arthrofactin synthetase from *Pseudomonas* sp. MIS38. *Chem. Biol.* 10:869-880.
- Roongsawang, N., Washio, K., and Morikawa, M. 2011. Diversity of non-ribosomal peptide synthetases involved in the biosynthesis of lipopeptide biosurfactants. *Int. J. Mol. Sci.* 12:141-172.
- Rottig, M., Medema, M. H., Blin, K., Weber, T., Rausch, C., and Kohlbacher, O. 2011. NRPSpredictor2-a web server for predicting NRPS adenylation domain specificity. *Nucleic Acids Res.* 39:W362-W367.
- Sambrook, J., and Russel, D. W. 2001. *Molecular Cloning: A Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, U.S.A.
- Schaad, N. W., Jones, J. B., and Chun, W. 2001. *Laboratory Guide for Identification of Plant Pathogenic Bacteria*. American Phytopathological Society Press, St. Paul, MN, U.S.A.
- Shanks, R. M. Q., Caiazza, N. C., Hinsä, S. M., Toutain, C. M., and O'Toole, G. A. 2006. *Saccharomyces cerevisiae*-based molecular tool kit for manipulation of genes from gram-negative bacteria. *Appl. Environ. Microbiol.* 72:5027-5036.
- Simon, R., Priefer, U., and Puhler, A. 1983. A broad host range mobilization system for in vivo genetic engineering: Transposon mutagenesis in gram-negative bacteria. *Nat. Biotechnol.* 1:784-791.
- Turnbull, G. A., Morgan, J. A., Whipps, J. M., and Saunders, J. R. 2001. The role of bacterial motility in the survival and spread of *Pseudomonas fluorescens* in soil and in the attachment and colonisation of wheat roots. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Ecol.* 36:21-31.
- Vallet-Gely, I., Novikov, A., Augusto, L., Liehl, P., Bolbach, G., Pechy-Tarr, M., Cosson, P., Keel, C., Caroff, M., and Lemaitre, B. 2010. Association of hemolytic activity of *Pseudomonas entomophila*, a versatile soil bacterium, with cyclic lipopeptide production. *Appl. Environ. Microbiol.* 76:910-921.
- Wilk, J. D., and Dye, D. W. 1974. *Pseudomonas cichorii* causing tomato and celery diseases in New Zealand. *N. Z. J. Agric. Res.* 17:123-130.
- Yamamoto, S., Kasai, H., Arnold, D. L., Jackson, R. W., Vivian, A., and Harayama, S. 2000. Phylogeny of the genus *Pseudomonas*: Intrageneric structure reconstructed from the nucleotide sequence of *gyrB* and *rpoD* genes. *Microbiology* 146:2385-2394.
- Yu, S.-M., and Lee, Y. H. 2012. First report of *Pseudomonas cichorii* associated with leaf spot on soybean in South Korea. *Plant Dis.* 96:142.

AUTHOR-RECOMMENDED INTERNET RESOURCES

NRPS/PKS analysis website: nrps.igs.umaryland.edu/nrps
 European Bioinformatics Institute (EBI) website: www.ebi.ac.uk
 NRPS-PKS (Nonribosomal Peptide Synthetases and Polyketide Synthases) website: www.nii.res.in/nrps-pks.html
 NRPSpredictor2 database: nrps.informatik.uni-tuebingen.de/
 University of Wisconsin Madison MAUVE genome alignment software: gel.ahabs.wisc.edu/mauve