G Protein α Subunit Genes Control Growth, Development, and Pathogenicity of *Magnaporthe grisea*

Shaohua Liu and Ralph A. Dean

Department of Plant Pathology and Physiology and Clemson Genome Laboratory, Clemson University, Clemson, SC 29634-0377, U.S.A.

Accepted 3 September 1997.

Three G protein α subunit genes have been cloned and characterized from *Magnaporthe grisea*: *magA* is very similar to *CPG-2* of *Cryphonectria parasitica*; *magB* is virtually identical to *CPG-1* of *Cryphonectria parasitica*, to *gna1* of *Neurospora crassa*, and to *fadA* of *Emericella nidulans*; and *magC* is most similar to *gna2* of *Neurospora crassa*. Homologous recombination resulting in targeted deletion of *magA* had no effect on vegetative growth, conidiation, or appressorium formation. Deletion of *magC* reduced conidiation, but did not affect vegetative growth or appressorium formation. However, disruption of *magB* significantly reduced vegetative growth, conidiation, and appressorium formation. *magB*[−] transformants, unlike *magA*[−] and *magC*[−] transformants, exhibited a reduced ability to infect and colonize susceptible rice leaves. G protein α subunit genes are required for *M. grisea* mating. *magB*[−] transformants failed to form perithecia, whereas *magA*[−] and *magC*[−] transformants did not produce mature asci. These results suggest that G protein α subunit genes are involved in signal transduction pathways in *M. grisea* that control vegetative growth, conidiation, conidium attachment, appressorium formation, mating, and pathogenicity.

Additional keywords: cAMP, *Pyricularia oryzae*.

Magnaporthe grisea (T. T. Hebert) Yaegashi & Udagawa (anamorph *Pyricularia oryzae* Cavara) causes rice blast disease, which severely reduces rice (*Oryza sativa* L.) production and continuously threatens food supplies worldwide, especially in many Asian nations (Ou 1985; Teng 1994). Controlling this fungal disease is an ongoing priority of plant pathologists (Ou 1985; Dobinson and Hamer 1992; Ford et al. 1994; Teng 1994). Recently, considerable research has been focused on the molecular mechanisms of rice blast infection (Valent 1990; Bourett and Howard 1990; Lee and Dean 1993; Talbot et al. 1993; Mitchell and Dean 1995; Shi and Leung 1994; Xu and Hamer 1996). The infection process is com-

posed of several key steps: conidiation, conidia dispersal, conidium attachment to the host surface, appressorium formation, penetration, and invasive growth. The goal is to fully understand the mechanisms crucial for each key step and use this information to develop durable and environmentally sound new strategies to control this devastating disease.

Previous research has demonstrated that the formation of the appressorium, a specialized infection cell, produced by *M. grisea* can be induced by environmental signals, including both physical (Bourett and Howard 1990; Dean 1992; Lee and Dean 1994; Gilbert et al. 1996) and/or chemical (Gilbert et al. 1996) cues. cAMP and signal transduction pathways have been shown to be central to appressorium formation (Lee and Dean 1993; Mitchell and Dean 1995; Xu and Hamer 1996). Taken together, these findings suggest that *M. grisea* conidia and/or germ tubes possess receptor(s) that sense suitable host plants in order to initiate the infection process. Regulatory guanine nucleotide-binding proteins (G proteins) are likely candidates to play an important role in the signal transduction cascade.

G proteins are involved in the transduction of signals from activated cell-surface receptors to intracellular effectors, such as adenylate cyclase, phospholipases, kinases, and ion channels (Stryer and Bourne 1986; Gilman 1987; Simon et al. 1991). Many of these G protein-mediated signals are essential for proper regulation of cell function, division, and differentiation.

The structure and function of heterotrimeric G proteins has been extensively reviewed (Gilman 1987; Strathmann et al. 1989; Simon et al. 1991; Coleman and Sprang 1996). The G protein complex of α , β , and γ subunits is anchored to the cytosolic face of the plasma membrane through hydrophobic regions on the β and γ subunits. The interaction of a G protein with an activated agonist-receptor complex promotes the exchange of GTP for bound GDP on the α subunit. This exchange causes a conformational change by dissociation of the α and $\beta\gamma$ subunits that enables the α subunit (or in some instances the $\beta\gamma$ complex) to modulate the activity of the appropriate effector protein. Therefore, the α subunit is responsible for the specific interactions with both the receptor and effector molecules. The $\beta\gamma$ subunit complex is usually responsible for inactivation of the α subunit and is also necessary for interaction of the G protein with receptors (Gilman 1987; Strathmann et al. 1989). The lifetime of the activated α subunit is inherently set by its GTPase activity. The cycle is thus reset when GTP is hydrolyzed to GDP.

Corresponding author: Ralph A. Dean; E-mail: rdean@clemson.edu

Present address of Shaohua Liu: MBD/NIDDK, NIH, Bldg 10/Room 8C-101, 10 Center Dr., Bethesda, MD 20892-1752, U.S.A.

Nucleotide and/or amino acid sequence data are to be found at GenBank as accession numbers AF011340 (*magA*), AF011341 (*magB*), and AF011342 (*magC*).

Numerous genes encoding the G protein subunits have been cloned, especially for the α subunits. Sequence comparisons have identified three major groups of G α genes, termed GPA, G α -I, and G α -II. The GPA group is primarily composed of G α genes from fungi and plants. Within both G α -I and G α -II, four classes have been identified that have functional similarities: Gi, Gq, G12, and Gs (Wilkie and Yokoyama 1994). Certain G α s appear to be restricted to a particular cell type and are involved in a single function (Brown and Birnbaumer 1990). In contrast, other α subunits are ubiquitous and serve multiple functions (Lerea et al. 1988). Sequence alignment of α subunit genes from fungi, plants, and animals reveals several highly conserved amino acid motifs that are thought to contribute to GTP binding. Degenerate polymerase chain reaction (PCR) primers, which target these motifs, have been used to selectively amplify genes encoding G protein α subunits from many organisms (Strathmann et al. 1989; Yatani et al. 1988; Tolkacheva et al. 1994). Several G protein α subunit genes have been isolated from filamentous fungi such as *Emmericella nidulans* and *Neurospora crassa* as well as from plant pathogens such as *Cryphonectria parasitica* and *Ustilago maydis* (Choi et al. 1995; Regenfelder et al. 1997; Turner and Borkovich 1993; Yu et al. 1996). Growing evidence has revealed that G α proteins are involved in regulating both development and pathogenicity. Choi et al. 1995) and Chen et al. 1996) reported that hypovirulence of the chestnut blight fungus *Cryphonectria parasitica*, which results from harboring RNA viruses of the genus *Hypovirus*, is mimicked by suppressing the expression of or deleting the G protein α subunit

gene *CPG-1*. CPG-1 appeared to function as a negative modulator of adenylate cyclase. In *U. maydis*, disruption of the G protein α subunit gene *gpa3* resulted in strains that were unable to respond to pheromone and were thus mating deficient. Gpa3 is also required for pathogen development (Regenfelder et al. 1997).

Here, we report on the identification of three G protein α subunit genes from *M. grisea* and present the effects of targeted deletion or disruption of these genes on growth, development, and pathogenicity.

RESULTS

Identification of G protein α subunit genes in *Magnaporthe grisea*.

Putative G protein α subunit gene fragments were obtained by PCR amplification of *M. grisea* genomic DNA with degenerate oligonucleotide primers to conserved regions in mammalian G protein α subunit genes. One forward and two different reverse primers were used. The PCR products were cloned into pBluescript II KS⁺. The relationship among the $\approx$ 200 recombinant colonies was determined by Southern hybridization with both the PCR products and representative clones as probes. Three distinctive groups of clones were found. Representative clones were sequenced, and three G protein α subunits were identified: *pSL* 1-1, *pSL* 2-3, and *pSL* 8. All three were highly similar to the conserved portion of mammalian subunits, but were different from each other. Percent identities of amino acid sequence between *pSL* 1-1 and *pSL* 8, *pSL* 1-1 and *pSL* 2-3, and *pSL* 8 and *pSL* 2-3, were about 60% (40/67), 70% (47/67), and 67% (45/67), respectively.

Southern blot analysis of restriction enzyme-digested *M. grisea* genomic DNA confirmed that the cloned PCR DNA fragments were present in the *M. grisea* genome (Fig. 1). Clamped homogeneous electric field (CHEF) gel analysis revealed that the three G protein α subunit clones are located on three separate chromosomes (data not shown). All three genes are single copy in the *M. grisea* genome.

Nucleotide sequence and features of *magA*, *magB*, and *magC*.

Cosmid clones corresponding to the three G protein α subunit PCR fragments were identified by Southern hybridization analysis and *Hind*III fragments were subcloned into pBluescript II KS⁺ (*pB*11 1-1, *pE*12 2-3, and *pD*10 8). The nucleotide sequence and deduced amino acid sequence for three G protein α subunit genes named *magA*, *magB*, and *magC* are shown in Figure 2. Amino acid sequence comparison between *magA* and *magB*, *magA* and *magC*, and *magB* and *magC* revealed relatively low levels of similarity: 48, 40, and 48%, respectively. However, as described below, each deduced polypeptide is closely related to other G protein α subunit genes found in fungi.

M. grisea magA encodes a polypeptide of 356 amino acids, contains five introns, and has 91% amino acid identity to *CPG-2* of *Cryphonectria parasitica*; *magB* encodes a polypeptide of 353 amino acids, contains three introns, and has 99% identity to *CPG-1* of *C. parasitica*, 98% identity to *gna1* of *Neurospora crassa*, and 93% identity to *fadA* of *Emmericella nidulans*; *magC* encodes a polypeptide of 355 amino acids, contains four introns, and has 81% identity to *gna2* of *N. crassa*.

Fig. 1. Southern blots of *M. grisea* wild-type strain 70-15 genomic DNA, which had been digested with *Hind*III (H), *Eco*RI (E), or *Bam*HI (B) and probed with ³²P-labeled *pSL* 1-1 (*magA*), *pSL* 2-3 (*magB*), and *pSL* 8 (*magC*), respectively. Blots were hybridized at 55°C as described in Materials and Methods.

magA

ctgtactact taggtctagg gaccacagtag attgttttct tatcctcgaac acgagacccgc caaattttacg gtcgcactgc tegtttgtctc tetgtctcgt
 tgctgcactc tgcagagtgc actagattga cacagcaatt cccaccagcc acccgccccc ccaagggttt gtttccgacg gagcggtgact gattaatcag
 attgtctgctc caattctctg tgcaaacccg ccgtacatac acaaccgact ctgtcatcaa ccaatacgtc tttgtcttgg tgccgctcac taaccgcgct
 atacctccga gccatccgaat ttctggcccg ggacctatcc tctccactcc gaagacgtat ttggcgctctt gctcgtctgg cgaaggtcta cgaaggtcta
 cttgaccgcc acgcccctggc acagttggcg atcaagttgc tatcccgggc gaccctgtgc cgcccgctcta tcacaatcgc togcacgccc tttgttgacg
 agtctgtctg tcacaggcca gctacgtgct gtcttaacaa ttcttccacg acttggcgatc agccggcgagc cttccatctc gggtaatcga
 acgaacgacg aagcatatcc catccgctct cgacaacaaa gcatccgcaa cgggggactt cgtatctccta cgttctgcta gacagaaagg tttatccgag
 acgactcgtc gatggggtcc gattaggctg ggtgctgttg ccattgttaag ccgcccgggc aacgaccttt gccagcgagc ctcgagtcoc
 agggcccccag ccattaaggg cgcaacgaaa gaggccttga aacccaatac aattgggttc cctggccaac aggaaccattc ATGATGGGCG CTTCATGAG

1
 21 TGCGAGCTCG GAGGAGTCGG AACAGAAGAA ACGCAGTCAA AAGATTGATA AGGATCTCGA GGAGGATCTA AAGAGATTGA GGCGAGAATG CAAGATTCTA
 8 A S S E E S E Q K K R S Q K I D K D L E E D S K R L R R E C K I L
 121 CTGTTGGTA agtttacctc qctctgtctt gacgagaggt tggtaaacgac gacgggtcat ttctcttaac aaattcacta caagtttqaa acgacgaggg
 41 L L G INTRON I
 221 tcatctctct taacaattca ctacagGTTCT TGGCAGAGT GCGAAGTCCA CAATCGTCAA GCAGATGAAG ATTATACATC TGAAAGGCTA CTCGGATGAC
 43 S G E S G K S T I V K Q M K I I H L K G Y S D D
 321 GAACTTTTACA ACTACCGGCC AACGGTCTTC AAGAATTAA TTGAATGTGC GAAAGCCGTG ATCAGCGCCA TGAGGCAATT CGACATAGAA CCGCAGCTGG
 68 E L Y N Y R P T V F K N L I E C A K A V I S A M R Q F D I E P Q L E

421 AAGGAACAA AGAAGTTTGC GACTATCTGC TCGACTACAT GGTCTGAGTC GGACACAGG CAGAGATAGA CCCCAGGTC GGTCAAGCTG TGCAGTCAAT
 102 G N K E L C D Y L L D Y M V E S G P Q A E I D P K V G Q A V Q S I
 521 ATGGAATGAT CCTGCAAGGG AACAGTTGAT GGACAGGCAG ACCGAGTTTT ACCTGATGGA CTCGGCAGAA Tagtaagttt caggttqaa atcgttqaa
 135 W N D P A R E Q L M D R Q T E F Y L M D S A E Y INTRON II
 621 taccgccgga agtttgcga ttgtgtcga aagctaacat tggctgtctt gacccaacca gcttcttcca AGAGTTTCTG AGGATTGTCT CTCGCCACTA
 159 CCTGCCCAAT GAGATGGATG TTCTTCGCGC AAGAACAAAA ACTACAGGCA TATACGAGAC ACGCTTTCAG ATGGGCCAAC TGAGCATACA gtaagcaagc
 721 L P N E M D V L R A R T K T T G I Y E T R F Q M G Q L S I H
 172 aaagaatcct qtcggaatat qcacgtgata aatcagatta ctgacttqca ttgcccqcca caqCATGTTT GATGTAGGAG GCCAGGAGG TGAGCGTAAG
 821 INTRON III
 202 AAGTGGATCC ACTGCTTCGA GAACGTGACA TCCATCATCT TTTGCTTGC ACTGATGAA TACGATCAAG TTTTACTTGA AGAAAGCAGT CAAGTCACTT
 921 K W I H C F E N V T S I I F C V A L S E Y D Q V L L E E S S Q
 214 taqcaataag ccccccagcc aaaaaatgca qttactgctt tctgacagcc caatgcatte ctatttggcc qcatgctaac ctaatatcgg acatacagaa
 1021 INTRON IV
 245 CCGCATGATG GAAAGCCTTC TGCTCTTCGA TTCGGTGGTA AATTCGAGGT GGTTCATGAG GACCAAGTATC ATATTGTTC TCAACAAGGT GGATATTTTC
 1121 R M M R S L L L F D S V V N S R W F M R T S I I L F L N K V D I F
 246 AAGCAGAAC TTGGCCGATC GCCTCTCGGC AACTACTTCC CAGATTACTC TGGAGGCAAC GATGTTAACA AGGCACAA GTACTTCTG TGGCGTTTA
 1221 K Q K L G R S P L G N Y F P D Y S G N D V N K A D K Y L L N R F N
 279 ACCAAGTTAA CAGGGCTCAC CTAAACCTAT ATCTCTCagt agtccaccct agaccaccaat gctgctactt aataatgttg ctgactttta taatcaccca
 1321 Q V N R A H L N L Y P H INTRON V
 313 GTCTTACGA AGTACCGAC ACCTCGAACA TTGCGATTGT CTTTGGCGCA GTAAAAGAGA CTATACTCAA CAATGCCCTG AAGGACTCAG GTATTCTGTA
 1421 L T Q A T D T S N I R L V F A A V K E T I L N N A L K D S G I L *
 325 Atcccgacga attctgacct acatttttct tcttatctct tcttatctacg ccaacttatg tttttatttg ataccacag caacttcata
 1521 ctgaatctat cttgcccagc gacttatatt atactagcat cagctcggag ttatggtggg cggtctctotg ctgttatgac taaaaacaa tttccctcaa

magB

cgatttgatt ctgtgcgcgc gatattcttc tctccctcct aacagtcata atattgcgat aacacatttt aggtacgaat gtcaatcgag gttataagag
 ggaatcctac caaaataaaa ataaaaagac aaaaacatca gactgaatga ctgagagctg gccccccgca tcagcagctc ccaatcgaa taattgactt
 gttgacctgg ttgcagctgg gttgcaaaaga acaattgggt gtagagctgc aagcagcagg acgtcgtctg acatagcgca gattggccag
 acctgtggcc acgcccggcc cccctctcca aattacagtc ctgtgtcttc tccaccaatt ttccgagctc ccaatttcaac ttaattcttc
 cctattgatt cgtgtgcgca acacettatt cttccgaaga agcggctctt cagccatcgc cagccatcgc cttgcttacc atacctctc
 ttctctactt gttgtttgt aaacccgcta ccaagccgto accggtatagg ctctgtttgg aattcggggc acgtcgggta cttgtcaatt cgccttattg
 taggcctgcc caattcccaa ttccccccct tccggtcgca tgcacaaaac attcagctgc cgtccacctc cctgcgaccc tgtcactagg tgcagctttt
 gatcccgagc caaatcgaaa agcaaatcct ctgcgcacac tggacacgga caccgctga caccgctga cagccattg cgaacagtg aataactctg
 tctacctaac catcgcttgt attaattatc gttcgacaaa caatcgcgca caattgggtg CGGAATGAGC ACGGAGGAGA AGGAGGCGAA GGCCCGGAAC
 -52 T E E K E S G K M S G A R N
 1 GAGGAGATTG AGAACCAGCT CAAGAGGGAC AGGTTGCAAC AGCGCAATGA GATCAAGATG CTCTGCTCG GTACATTCGAG GTGAATCCGG AAAGTCCACC ATTCTCAAGC
 49 E E I L K R D R L Q R N E I K M L L L G
 17 qggccagcat qcccgagcat aaggtttttt ttttctttat gtcgctaaca tctcgggtct acagGAGCCG GTGAATCCGG AAAGTCCACC ATTCTCAAGC
 149 INTRON I
 41 AGATGAAGCT CATTACAGAA GGCGGCTACT CACGGGACGA GCGCGAATCT TTCAAGAA TCACTCTCAG CAACACTGTA CAGTCTATGC GTGTCTCTT
 249 M K L I H E G G Y S R D G R E S F K E I I F S N T V Q S M R V I L
 53 GGAGGCCATG GAGTCGCTCG AGCTGCCTCT GGAGGACCAA AGGATGGAGT ACCAGTACA GACCATCTTC ATGCAGCCTG CTCAGATCGA AGGCGACGTC
 349 E A M E S L E L P L E D Q R M E Y H V Q T I F M Q P A Q I E G D V
 86 TTGCCACAG AGTCCGCAA CGCCATCGAG GCCTGTGGA AGGATCTGGT TGTCCAGGAG TGCTTCAAGC GTTCAAGGA ATACCACTTG AACCACTCTG
 449 L P P E V G N A I E A L W K D R G V Q E C F K R S R E Y Q L N D S A

549 CCAGATGata ggtttttaccg ttggtctcgaq aaaaaccttat gctgcagcta cccctctgct tgaacgttga atgatgctaa cgcacaatga agCTACTTCG
 153 R Y INTRON II
 649 ACAATATTGC TCGATTGCT GCACCAAGCT ACATGCCTAA CGACCAAGAC GTGCTTCGAT CTCGCGTCAA GACAACCGGT ATCACTGAGA CGACTTTCAT
 158 N I A R I A A P D Y M P N D Q D V L R S R V K T T G I T E T T F I
 749 CATTGGCCAC CTCACCTACC GAATGTTCTGA TGTGTTGGC CAACGTTCTG AGCGAAAGAA GTGAGTTCAC TGCTTCGAGA ACGTTACAC GATTCTCTTC
 191 I G D L T Y R M F D V G G Q R S E R K K W I H C F E N V T T I L F
 849 CTCGTCGCCA TTTCCGAGTA CGATCAGCTA TTGTTCCGAG ACAGAGCGGT CAACCGATG CAGGAGGCTC TCACACTATT CGACTCCATC TGCAACTCGA
 224 L V A I S E Y D Q L L F E D E T V N R M Q E A L T L F T L F C N S R
 949 GGTGTTCAT CAAACCTCC ATCACTCTCT TCTTCAACAA GATCGATCGC TTAAAGGAGA AGTGCCCAT CAGCCCGATG AAGACTACT TCCCGACTA
 258 W F I K T S I T L F L N K I D R F K E K L P I S P M K N Y F P D Y
 1049 CGAGGGTGGC GAGCACTAG CTGAGCCTG CGACTACATC CTCACCGAT CTGATCGCT TTTAAGGAGA AGTGCCCAT CAGCCCGATG AAGACTACT TCCCGACTA
 291 E G G D Y A A A C D Y I L N R F V S L N Q H E T K Q I Y T H F T
 1149 TGCGCCACAG ACACCAACA GATCCGCTTC GTCATGGCG CTGTTAATG taagaaatca tgcgtccctg gattggaatt actgatcgc caqacattct
 324 C A T D T T Q I R F V M A A V N D
 1249 aactgttaca aatagACATC ATCATAAG AGAACCTCCG TCTCTGTTGT CTGATCTAAa atcagtgccc INTRON III
 341 ctcagctttt cctcgggata I I Q E N L R L C G L I *
 tctgtactc cccccaaag atcaataatc gacgagcctt ccaaacgttc cctcctcttg tgcacctctc tgccaacgga gtcacaggcg
 tgggggtttt tgtgggtgtt ttttttactt caaaactctg ttggtatttt ccaagcctt ccaactcttg aaactcttg gtcacaggcg cctgtttttg

Fig. 2. Nucleic acid and deduced amino acid (italic letters) sequences for *magA*, *magB*, and *magC*. Introns are in underlined lowercase letters. The GenBank accession numbers are AF011340, AF011341, and AF011342, respectively.

Figure 2 continued on next page.

Like CPG-1 of *C. parasitica*, *gna1* of *N. crassa*, and *fdaA* of *E. nidulans*, *magB* contains a consensus myristylation site (MGXXXS) (Buss et al. 1987) and pertussis toxin-labeling site (CXXX) (West et al. 1985) at its N and C termini, respectively: features characteristic of members of the inhibitory G (Gi) protein α subunit family of mammals (West et al. 1985; Simon et al. 1991; Spiegel et al. 1991). *magA* contains a potential consensus myristylation site (MMGXXXS) at its N terminus but does not have a pertussis toxin-labeling site at its C terminus. *magC* does not contain either site.

Gene deletion of *magA*, *magB*, and *magC*.

To determine the functions of *magA* and *magC*, homologous recombination was employed to replace the target gene with a selectable marker. For *magB*, a fragment internal to the target gene was used for disruption (Timberlake and Marshall 1989). *magA* was replaced by plasmid pB11pan, which contains the hygromycin phosphor transferase (*hph*) gene cassette flanked by 5' and 3' sequences from the *magA* gene (Fig. 3A). Southern analysis confirmed that six out of 65 hygromycin-resistant transformants had undergone the double crossover event to delete the *magA* gene (Fig. 4A). *magC* was replaced in a similar manner by plasmid p8-span, which contains the *hph* cassette flanked by 5' and 3' sequences from the *magC* gene (Fig. 3C). Six out of 54 hygromycin-resistant transformants were confirmed by Southern analysis to be deleted for *magC* (Fig. 4C). *magB* was disrupted by plasmid pAN2-3-6, which contains an internal ≈ 0.7 -kbp fragment (amino acids 15 to 196) of *magB* and the *hph* (Fig. 3B). Four out of 92 hygro-

mycin-resistant transformants were shown to be disrupted for *magB* (Figure 4B). Based on additional genomic Southern analysis, two of these transformants (2-9 and 2-11) had undergone a single plasmid DNA insertion event, whereas the other two (2-2 and 2-12) contained multiple insertions.

Effects of disruption of *magA*, *magB*, and *magC* on fungal growth, conidiation, and appressorium formation.

The effects of gene disruption on fungal growth, conidiation, and appressorium formation are shown in Table 1 and Figure 5, unless otherwise indicated. Targeted deletion of *magA* did not affect vegetative growth on different media (V8 or CM; data not shown), conidiation, or appressorium formation. Deletion of *magC* reduced conidia formation by greater than twofold, but did not affect vegetative growth on different media (CM data not shown) or appressorium formation. However, disruption of *magB* significantly reduced vegetative growth on different growth media, conidiation, and appressorium formation. Following 72 h of incubation, appressorium formation by *magB*⁻ transformants was still less than 5%.

Targeted deletion/disruption of G protein α subunit genes (*magA*, *magB*, or *magC*) did not affect conidium germination on the hydrophobic surface of GelBond (data not shown). The morphology of germ tubes of *magA*⁻ and *magC*⁻ transformants appeared similar to that of wild type (Fig. 5A, C, and D), i.e., the germ tubes were short and curvy; however, the germ tubes of *magB*⁻ transformants were long and straight (Fig. 5B).

Conidium attachment of *magB*⁻ transformants was reduced on the hydrophobic surface of GelBond, compared with that

Figure 2 continued from previous page.

	magC										
	tggaggctgg	acagataatg	tcagcgcaca	tacataat	tactgtcctt	tactttgggt	acacaacacc	ggaagagatg	tccgttgtat	caagccgaag	
	caaaagccaat	ctccocagcg	cgcgcacac	taccgcctt	ggccctccca	gacccttggc	atttaaccca	gcctaaaaatt	ctggtgtgat	ctagtctgggt	
	tccaggtaacc	cacctaggtta	cctatgtatc	tctcgtaggc	tgcaaggctt	ggtaggttaa	cttgcaacacc	accacgggtgc	ttgtccttat	ccaccttgtc	
	acacctgttcc	acctatccac	ccgctccgta	ccttgccgtg	ctttacatcca	ctccccaaca	aaccgcgtacg	gttagcttcc	ttgtgttatt	cgcttcccg	
	gcctgtgatat	cgctgcacat	cttcacgcac	gtgcattgga	gatccctcat	ctctttatcc	cgcccccgcg	cgccccaaga	caagcaaaa	aataataata	
	atagggggag	ccgcttctgt	caagtcacgc	gattgcgact	aacgcctatg	cgaccctccg	cgctcaattt	gtttcattcc	atcaccattt	ggttctcctc	
	ccatagcagg	caagacggac	ccggacctgt	cagaagacgt	gcgacacaca	caaattctga	ccaattatta	cctaggttagc	cacaaagcga	caaaacagtc	
	gcaccgcaga	aatcgacctt	gaataggtct	ttttactcgt	tagtttaatac	aagcaactcag	ttgcttcgac	ctcatcgac	cacttgtctg	gagcaatccg	
	ccaattgccc	tctggcaaga	GTGCTTCGG	CAGCCGCAAC	GTCCACGATG	AGGCGGGGGC	TGCCCGCAGC	AGGGAGCTCG	ACAAGGCCATC	CGCGCCGAC	
-19			M	C F G	S R N	V H D E	A G A	A R S	R E L D	K A I R	A R A D
1	GAGAAGCGCT	TGCAAAAGGA	GGTCAAAGT	CTGCTGCTGG	gtatgtcgc	gagcaatgtt	tacattcgaq	qccactcgtt	atctggtcca	aatcgaagac	
81	E K R L	Q K E	V K L	L L L G	INTRON I						
28	actctgctaa	ccaagtgtgt	tttttcgata	ccatagcgcg	CGGTGAATCG	GGAAAATCGA	CCGTCTCTCA	GCAATGAAG	CTGATATATG	CTCAAGGCTT	
181				A	G E S	G K S T	V L K	Q M K	L I Y A	Q G F	
42	CAGCAAGAAC	GAAAGATCG	AATGGAAGCC	TGTCGTGTTT	CAACACATTT	TGCAATCTTT	CAGGCTTATC	CATGATGCCA	TGCAAGAGTT	GGACATCGAA	
281	S K N	E K I E	W K P	V V F	Q H I V	Q S F	R L I	H D A M	Q E L D	I E	
63	TTTCGAGAAC	AGGAGAACGA	Ggtatgttaa	aactctgcac	tttctattt	catgacaaat	aggtctcgcg	cactcgattc	cctaccggac	tcgccccat	
381	F E N K	E N E	INTRON II								
96	catcctgacg	tccatccttc	cgcccccacg	aatgcactac	aaagcaacaa	aaagcctgac	gggttttqaa	catataacta	acatccattt	ttggtctacag	
481	AGGAACATAA	CCCTGATCAT	GGAGGAGCAG	GAATCGGGC	CGCATGACGC	CTTACCAATC	GAGTTTCTAG	ACCCAATCAA	AGCACTATGG	GTGGATGCTG	
581	R N I T	L I M	E E Q	E L G A	H D A	L P I	E F L D	P I K A	A L W V	D A G	
103	CGGTGAAGAA	GGCGATTGCC	AAGGGCAACG	AGTTTGCCTG	ACATGACAA	CTGGACTAGt	gagtgcat	tccccaaact	tcggctaatc	atgcctttat	
681	V K K A	I A I	K G N E	F A L	H D N	L D Y	INTRON III				
137	tttgtttctt	tacgaagccc	ctaatacaca	qTTTCTGTGA	CGACCTCGAC	CGTATTGTTG	ATAAGAGCTA	CGTTCGACA	GACCAAGATT	TGCTCCGCTC	
781				F C D	D L D	R I W D	K S Y	V P T	D Q D L	L R S	
156	GAGACTCAGA	ACGACCGGCA	TTACCGAGAC	CGTGTGTTGAC	TTGGGCCAGC	TCACATACCG	AATGTTTCAT	GTGGTGGCC	AAAGTTCGGA	AAGGAAGAAG	
881	R L R T	T T G I	T E T	V F D	L G Q L	T Y R	M F D	V G G Q	R S E R	R K K	
179	TGGATACATT	GTTTCGAGAA	CGTCAACTGT	CTTCTTTTCC	TAGTCGCAAT	TTCCGGCTAC	GATCAGTGT	TAGTTGAGGA	TAAGGACGGA	gtaagttctca	
981	W I H C	F E N	V N C	L L F L	V A I	S G Y	D Q C L	V E D	K D G		
212	acaccgcaag	cacacattta	taaaacagca	caqcttggca	acagtttttg	tcagccctag	gaqctcatat	ctaaccataa	cccqctqga	caqAACCAAA	
1081	INTRON IV										
242	TGAACGAAGC	CTTGATGCTG	TGGGAGTCGA	TCGCCAASCT	GCATTGGTTC	ACAAAATCGG	CGCTGATCTT	GTTCCTCAAC	AAGATGGACC	TGTTCAAGGA	N Q M
1181	N E A L	M L M	W E S I	A A N S	H W F	T K S A	L I L	F L N	K M D L	F K E	
245	GAAAATCACA	AAGAGTCCCA	TTACCGAATA	TGGGTTCACC	GACTATACG	GACCATCAGA	CGAGCTGGAAG	CAGACTAGCA	AGTACTTTAT	GGACAAGTTC	
1281	K I T	K S P I	T E Y	G F T	D Y H C	P S D	D W K	Q T S K	Y F M D	D K F	
278	CGAGCACTTA	ACCGAAACCC	AGAGAAAGAG	ATATACGGCC	ACTTCACAAA	CGTACCGGAC	ACGAACCTAC	TCAAGATCAC	GATGACGTCT	GTGCAGGACA	
1381											
311	R A L N	R N P	E K E	I Y G H	F T N	A T D	T N L L	K I T	M T S	V Q D M	
1481	TGATTATACA	GCGCAACCTT	AAGCAACTCA	TCITGTGAGt	gctactccgg	cggtcccggt	taataaccta	actggcggtc	gaatcttcac	tatcaccagc	
345	I I Q	R N L	K Q L I	L							
	acactcacac	ctctggacac	cattctagcg	tttgattaat	cctagacggg	gcaaaattctg	ccacatcgcc	gggtcttgta	cttcgacaaa	ccaactgcac	
	catctaccgc	gctgtccgct	accgctttct	tgaataacct	cagataaccga	aggttcttat	ctgtgttttt	atttcgtttt	tttcattttt	tttcattttt	
	tttctggctt	tggtcttatt	ttgtccatta	tcacattgga	cttttcgctg	aagctacctg	gctgcattgg	cttggttaaa	ttgtatgat	tggtgtattc	
	ctctttcagg	gtgtacatac	atggccgctgc	accagcatct	tggtctt						

of controls (Table 2). Fewer than 20% conidia of *magB*⁻ transformants remained attached to GelBond whereas about 50% of conidia from controls remained attached when rinsed with water after a 2-h incubation.

Appressorium formation by *magB*⁻ transformants could be restored by adding cAMP (Table 3; Fig. 6A) or 1,16-hexadecanediol (Table 3; Fig. 6B). Chemically induced appressoria formed on the tips of long, straight germ tubes.

Effect of deletion of *magB* on the ability to infect and colonize susceptible rice.

Deletion of *magB* reduced *M. grisea* virulence, as shown in Figure 7. Disease symptoms (necrotic lesions) on rice leaves were greatly reduced when inoculated with 10⁴ conidia either by injection (Fig. 7B,C) or by spray, compared with the wild-type (Fig. 7E) and an ectopic integration control (Fig. 7D). A few spreading lesions were observed when leaf sheaths were injected with *magB*⁻ transformants, although wound sites on the emerging leaves were clearly visible.

Direct observation of inoculated rice leaves under the microscope revealed that appressorium formation of *magB*⁻ transformants was reduced. When young rice seedlings (2 weeks old) were spray inoculated, germinated conidia of *magB*⁻ transformants formed very few appressoria (<3%) on leaves (Fig. 8B), compared with wild-type and ectopic integration controls (>95; Fig. 8A). However, when older rice plants (4 to 5 weeks old) were spray inoculated with *magB*⁻ transformants, appressorium formation increased with the age of the leaves; only 20 to 30% of germinated conidia produced appressoria on the young leaves (Fig. 8D), compared with >50% on the oldest leaf (Fig. 8C). For wild-type and ectopic integration transformant controls, no difference in appressorium formation was observed on either young or old leaves; appressorium formation always exceeded 95%. Appressoria produced by *magB*⁻ transformants on rice leaves always formed on the tips of long, straight germ tubes and were very similar to those induced by chemicals on the hydrophobic surface of GelBond (Fig. 6).

The deletion of *magA* or *magC* did not appear to affect the ability to infect and spread with the host tissues (data not shown), both gene deletion transformants caused typical disease symptoms indistinguishable from wild-type and ectopic integration controls.

Recovery of the wild-type phenotype by restoring *magB* to a gene-disrupted transformant.

magB⁻ transformant 2-12 was transformed with a plasmid (pE12Bar) containing the *magB* gene with the *bar* gene as a selective marker. Southern blot analysis confirmed the introduction of the plasmid (Fig. 9, lanes 9 to 11). Resulting transformants exhibited typical wild-type characteristics, including normal vegetative growth on different media, normal conidiation, normal appressorium formation on the hydrophobic surface of GelBond, and pathogenicity indistinguishable from that of the wild type (Table 4; Fig. 7F,G). *magB*-restored transformants produced appressoria on short, curved germ tubes.

G protein α subunits are required for *M. grisea* mating.

In repeated three-way crossing experiments with 70-6, 70-15, and *magB*⁻ transformants, perithecia failed to form be-

tween 70-6 and *magB*⁻ strains, whereas 70-6 and 70-15 formed perithecia and visual asci that germinated normally. Similar crosses involving *magA*⁻ and *magC*⁻ transformants formed perithecia; asci were visible but ascospores were indistinct and failed to germinate.

DISCUSSION

Our results indicate that *M. grisea* has multiple G protein α subunit genes. Each gene appears to be involved in different signal transduction pathways, as deletion of each gene affected different aspects of growth, development, and pathogenicity.

magB was disrupted with a DNA fragment encoding an internal part of the protein (amino acids 16 to 196). Transformants that had undergone a single homologous recombination event thus contained 2 truncated copies of the gene, one lacking the promoter and the first 192 amino acids, and the second lacking amino acids 193 to 353. It is highly unlikely that disruption strains would retain functional *magB* activity because

Fig. 3. Strategy for disruption of G protein α subunit genes in *Magnaporthe grisea*. A, Deletion of *magA*; B, Disruption of *magB*; C, Deletion of *magC*. B = BamHI, E = EcoRI, H = HindIII, S = SacI, Sp = SphI, St = StuI, Xb = XbaI, X = XhoI.

essential features including GTP-binding domains were lacking in both truncated copies. Indeed, *magB* apparently controls multiple functions, including vegetative growth, conidium formation, mating, and appressorium formation. Reduction of appressorium formation in *magB*⁻ transformants clearly indicates *magB* contributes to reduced pathogenicity, but based on the sheath injection experiments, *magB* also contributes to invasive growth.

Upon amino acid sequence inspection, *magB* contains certain features common to Gi protein α subunits. In mammalian systems, release of Gi protein α subunit lowers intracellular cAMP levels by inhibition of adenylate cyclase activity (Childers and Deadwyler 1996; Chen et al. 1996). Lee and Dean 1993, 1994) found that cAMP induces appressorium formation in *M. grisea*. Therefore, we may have expected that the deletion of *magB* might stimulate appressorium formation by removing an inhibitor of adenylate cyclase, and consequently increasing the intracellular cAMP level. However, deletion of *magB* in *M. grisea* blocked appressorium formation. Furthermore, adding external cAMP (or IBMX) induced appressorium formation of *magB*⁻ transformants. An adenylate cyclase gene of *M. grisea* has been cloned in our laboratory (W. Choi and R. A. Dean, *unpublished data*). The relationship between *magB* and adenylate cyclase in *M. grisea* is under investigation. It should be noted that, based on sequence alignments, the G protein α subunits of simple eukaryotes form a unique cluster (GPA) distinct from any mammalian G α class genes (Wilkie and Yokoyama 1994). Therefore, the function of G α subunit genes in simple eukaryotes may differ from that in their mammalian counterparts.

The attachment of conidia of *magB*⁻ to hydrophobic surface was substantially reduced; whether this is related to the failure to form appressoria is unknown. Surface proteins with possible adhesive properties such as mucilage, hydrophobin, or other secreted proteins have been postulated to be involved in recognizing environmental signals (Hamer et al. 1988; Beckerman and Ebbole 1996). These signals may be transduced into cells through *magB*, which then activates the differentiation pathway resulting in appressorium formation. How deletion of *magB* could affect surface attachment remains to be determined.

We checked whether *magB*⁻ transformants have an easily wettable phenotype, as described by Talbot et al. 1993) for strains lacking the hydrophobin MPG1. All *magB*⁻ mutant colonies on CM agar showed no wettability; drops (200 μ l) of H₂O remained in place on colonies up to 7 days without any visible change. Beckerman and Ebbole 1996) reported that, when mixed with wild-type conidia, strains deleted for *MPG1* could produce appressoria by utilizing the hydrophobin from wild-type conidia. When mixed with wild-type (70-15) conidia, *magB*⁻ transformants were clearly identifiable and continued to form distinctive straight, long germ tubes without appressoria. Apparently, deletion of *magB* did not affect hydrophobin production, and hydrophobin did not compensate the loss of *magB*. Thus, *magB*⁻ transformants may be defective in attachment independent of hydrophobins.

The difference between young and old rice leaves for inducing appressorium formation of *magB*⁻ transformants suggests that there is a difference in surface cues (either chemical or physical) presented between young and older leaves. Young

Fig. 4. Southern blots of G protein α subunit disruptions. All genomic DNA was digested with *Hind*III, and probed with a *Hind*III fragment of *magA* (4.8 kbp) for blot **A**, *magB* (4.3 kbp) for blot **B**, or *magC* (6.8 kbp) for blot **C**, respectively. **A**, Lane 1 is wild-type strain DNA (70-15); lane 2, ectopic integration strain (B1); lanes 3–8, *magA*⁻ transformants (B3, B6, B8, B14, B17, and B18). As expected, the 4.8-kbp band from *magA*⁻ transformants is no longer present. **B**, Lane 1, wild-type (70-15); lanes 2–4, ectopic integrations (2-3, 2-5, and 2-10); lanes 5–8, *magB*⁻ transformants (2-9, 2-11, 2-2, and 2-12). As expected, the 4.3-kbp band disappeared in *magB*⁻ transformants. For 2-9 and 2-11, a single integration event had occurred, whereas for 2-2 and 2-12, multiple insertion events occurred, as evidenced by an additional 7.5-kbp band (the size of the disruption plasmid). **C**, Lane 1, wild-type; lanes 4 and 8, ectopic integrations (8-8 and 8-18); lanes 2, 3, 5–7, and 9, *magC*⁻ transformants (8-2, 8-7, 8-10, 8-12, 8-17, and 8-19). As expected, the 6.8-kbp band is missing in *magC*⁻ transformants.

leaves may contain some chemical inhibitor(s), which can prevent appressorium formation of *magB*⁻ transformants. More likely, young leaves lack certain chemical stimulatory compounds. As leaves mature, the appropriate chemicals may be produced and deposited on to the surface. Potential candidates may be chemically similar to 1,16-hexadecanediol. This molecule, a minor cutin component, induced appressorium formation in *magB*⁻ transformants with a morphology similar to that observed on mature rice leaves; appressoria developed at the tips of long, straight germ tubes. Gilbert et al. 1996) reported that 1,16-hexadecanediol induced appressorium formation of *M. grisea* at nanomolar levels. This molecule and related chemicals commonly exist on the surface of plants, including on mature rice leaves. Further investigation is required to examine the presence of stimulating chemicals from young and mature leaves before firm conclusions can be drawn.

The mechanism of action of 1,16-hexadecanediol is unknown. Regardless, it appears to be independent of *magB*. *magB*, on the other hand, appears to be linked to the reception and processing of physical signals, such as hydrophobicity. These data reinforce previous evidence that there are at least dual pathways regulating the induction of appressoria (Gilbert et al. 1996).

Table 1. Vegetative growth, conidiation, and appressorium formation assay for G protein α subunit gene transformants^a

Strain	Colony diameter (cm)			Conidia ($\times 10^4$)	Appress. (%)
	5-day-old	7-day-old	10-day-old		
Transformants lacking <i>magA</i> ^V					
Ck ^w 70-15	2.2 $\pm$ 0.3	3.4 $\pm$ 0.3	5.0 $\pm$ 0.3	65.0 $\pm$ 12.0	>95
E ^x B1	2.2 $\pm$ 0.3	3.3 $\pm$ 0.3	4.7 $\pm$ 0.4	82.2 $\pm$ 5.8	>95
E B2	2.2 $\pm$ 0.3	3.3 $\pm$ 0.3	4.6 $\pm$ 0.3	55.8 $\pm$ 6.5	>95
M ^y B3	2.5 $\pm$ 0.3	3.1 $\pm$ 0.3	4.6 $\pm$ 0.3	67.6 $\pm$ 5.5	>95
M B6	2.3 $\pm$ 0.3	3.5 $\pm$ 0.3	4.9 $\pm$ 0.3	72.8 $\pm$ 2.3	>95
M B8	2.3 $\pm$ 0.3	3.4 $\pm$ 0.3	4.9 $\pm$ 0.3	53.4 $\pm$ 1.0	>95
M B14	2.3 $\pm$ 0.2	3.4 $\pm$ 0.2	5.0 $\pm$ 0.3	70.6 $\pm$ 11.2	>95
M B17	2.1 $\pm$ 0.3	3.3 $\pm$ 0.3	4.7 $\pm$ 0.3	70.2 $\pm$ 10.6	>95
M B18	2.3 $\pm$ 0.3	3.4 $\pm$ 0.3	4.8 $\pm$ 0.3	70.2 $\pm$ 9.9	>95
Transformants lacking <i>magB</i>					
Ck 70-15	2.7 $\pm$ 0.3 a ^z	4.2 $\pm$ 0.3 a	6.0 $\pm$ 0.3 a	61.3 $\pm$ 4.6 a	>95 a
E 2-3	2.6 $\pm$ 0.3 a	4.0 $\pm$ 0.3 a	5.9 $\pm$ 0.3 a	68.1 $\pm$ 8.0 a	>95 a
E 2-10	2.7 $\pm$ 0.3 a	4.0 $\pm$ 0.3 a	5.9 $\pm$ 0.3 a	75.5 $\pm$ 6.9 a	>95 a
M 2-2	2.3 $\pm$ 0.3 b	3.4 $\pm$ 0.3 b	4.8 $\pm$ 0.3 b	0.3 $\pm$ 0.2 c	2.5 $\pm$ 0.6 b
M 2-9	2.3 $\pm$ 0.4 b	3.5 $\pm$ 0.4 b	5.0 $\pm$ 0.3 b	21.7 $\pm$ 4.0 b	1.8 $\pm$ 0.5 b
M 2-11	2.4 $\pm$ 0.3 b	3.6 $\pm$ 0.2 b	5.1 $\pm$ 0.3 b	1.2 $\pm$ 0.6 c	3.7 $\pm$ 0.9 b
M 2-12	2.3 $\pm$ 0.3 b	3.4 $\pm$ 0.2 b	5.0 $\pm$ 0.3 b	1.9 $\pm$ 0.6 c	1.2 $\pm$ 0.9 b
Transformants lacking <i>magC</i>					
Ck 70-15	2.3 $\pm$ 0.3	3.4 $\pm$ 0.3	5.1 $\pm$ 0.3	76.8 $\pm$ 7.0 a	>95
E 8-8	2.2 $\pm$ 0.3	3.1 $\pm$ 0.4	4.7 $\pm$ 0.3	71.5 $\pm$ 5.0 a	>95
E 8-18	2.5 $\pm$ 0.3	3.5 $\pm$ 0.3	5.2 $\pm$ 0.3	76.3 $\pm$ 13.1 a	>95
M 8-7	2.2 $\pm$ 0.3	3.1 $\pm$ 0.3	4.7 $\pm$ 0.5	31.4 $\pm$ 8.6 b	>95
M 8-17	2.1 $\pm$ 0.4	3.0 $\pm$ 0.4	4.5 $\pm$ 0.4	21.9 $\pm$ 7.9 b	>95
M 8-19	2.4 $\pm$ 0.3	3.5 $\pm$ 0.3	5.1 $\pm$ 0.3	10.2 $\pm$ 5.2 b	>95

^a Colonies were grown on V8 media under fluorescent light. For appressorium assays, conidia were collected from 7-day-old cultures and appressorium formation was recorded after 20 h on the hydrophobic surface of GelBond. The data from three independent experiments, each with two or three replicates; mean $\pm$ standard deviation.

^b No statistically significant differences were observed between strains as determined by Student's *t* test.

^w Ck is the wild-type control strain 70-15.

^x E designates ectopic integration transformants.

^y M designates gene disrupted mutants.

^z Different letters in each data column indicate significant differences at *P* = 0.05.

In a recent report, Beckerman et al. 1997) discovered that the sex pheromone, α -mating factor from *Saccharomyces cerevisiae*, and a cell-free extract from *M. grisea* inhibit appressorium formation in a mating type-specific manner. The evidence reveals a possible relationship between the signal pathways for mating and appressorium formation.

Deletion of *magB* blocked sexual development as well as inhibiting appressorium formation. All *magB*⁻ transformants tested failed to form perithecia when crossed with the appropriate mating-type strain. This may suggest that *magB* interacts with sex pheromone receptor(s). Thus, removing *magB* might break the pathway between the receptor and effector(s), and prevent the initiation of the mating process. Whether the putative sex pheromone receptor(s) in *M. grisea* have dual functions will require further investigation.

It is also noteworthy that deletion of *magB* homologs in other filamentous fungi has similar phenotypic consequences. Deletion of *CPG1* of *C. parasitica* resulted in a marked reduction in growth and a loss of virulence, asexual reproduction, female fertility, and laccase activity (Gao and Nuss 1996); deletion of a G protein α subunit gene with high homology to *magB* in the corn pathogen *Cochliobolus heterotrophus* resulted in a loss of the ability to form appressoria and the production of infertile pseudothecia (Horowitz 1997). In *N. crassa*, deletion of *gna-1* caused abnormal vegetative growth and aerial hyphae as well as female sterility (Ivey et al. 1996). Disruption of *fadA* in *A. nidulans* also reduced vegetative growth, but did not affect conidiation or sexual competency (Yu et al. 1996).

Deletion of *magA* or *magC* on the other hand had minimal effect, although loss of these genes clearly affected a later stage of sexual development: perithecia formed but ascospores failed to germinate. Therefore these genes appear to function in ascospore maturation and/or germination.

Overall, our results indicate that different G protein α subunits control unique signal transduction pathways in *M. grisea*, involving vegetative growth, conidiation, appressorium formation, mating, and pathogenicity. It is possible the deletion of one gene may be compensated for by another, i.e., functional redundancy or pathway cross-talk. Research is presently addressing this possibility. Future insight into the role of G

Fig. 5. Appressorium formation (at 24 h) on the hydrophobic surface of GelBond. **A**, Wild-type strain (70-15); **B**, *magB*⁻ transformant; **C**, *magA*⁻ transformant; and **D**, *magC*⁻ transformant. Magnification = $\times 200$.

proteins in mediating morphogenesis will be provided by specific mutagenesis of the α subunits affecting their activity.

MATERIALS AND METHODS

Strains, growth, and maintenance of *Magnaporthe grisea*.

Strains of *M. grisea*, 70-15 (MAT1-1) and 70-6 (MAT1-2), used in this study were originally obtained from A. Ellingboe (University of Wisconsin-Madison) and maintained on complete medium (CM) with 1.5% agar (Lau et al. 1993; Mitchell and Dean 1995). All transformant isolates were maintained on CM agar with 200 μ g of hygromycin B per ml or 500 μ g of basta per ml.

M. grisea conidia were produced on V8 agar medium (4% vol/vol V8 juice, 15 g of agar per liter, pH 7.0, with or without 200 μ g of hygromycin B per ml), and grown under fluorescent light to induce conidiation.

Mycelia were produced by transferring conidia of *M. grisea* isolates into 20 ml of liquid CM in petri dishes (1.0×10^6 conidia per petri dish), and grown at 23°C in the dark for 4 to 5 days.

PCR cloning of G protein α subunit genes.

The degenerate oligonucleotide primers oMP 19, oMP 20, and T α 29 [oMP 19: CGGATCCAA(AG)TGGAT(AC)CA

(CT)TG(CT)TT; oMP 20: GGAATTC (AG)TC(CT)TT(CT)TT(AC)TT(ACGT)AG(AG)AA; T α 29: GGGATCC(ACT)GT(AG)TC(CT)GT(CT)GC(AG)CA(CT)GT], complementary to a conserved portion of mammalian G protein α subunits, were provided by T. M. Wilkie (University of Texas Southwestern Medical Center, Dallas). The underlined regions identify added restriction sites. PCR was performed on *M. grisea* genomic DNA with primer pairs oMP 19 $\times$ oMP 20 or oMP 19 $\times$ T α 29.

Fifty-microliter PCR reactions contained 5 μ l of 10 $\times$ PCR buffer (Perkin-Elmer Applied Biosystems Division, Foster City, CA), 8 μ l of 25 mM MgCl₂, 1 μ l each of dATP, dCTP, dGTP, and dTTP (10 mM stock), 1 μ l of *M. grisea* genomic DNA (5 ng of stock per μ l), 1 μ l each of forward or reverse primer (50 ng of stock per μ l), 30 μ l of distilled H₂O, and 0.2 μ l of AmpliTaq Polymerase (Perkin-Elmer Applied Biosystems Division). PCR conditions were as follows: 3 min at 94°C, followed by 35 cycles (1 min at 94°C; 1.5 min at 45°C; 2 min at 72°C), with a final extension at 72°C for 10 min.

PCR products were digested with *Bam*HI and *Eco*RI and cloned into *pBluescript* II KS⁺, and then transformed into *Escherichia coli* strain DH 5 α . The relationship among all white colonies was determined by Southern hybridization. Nucleotide sequence of representative clones was determined

Table 2. Attachment of conidia of *magB*[−] transformants to the hydrophobic surface of GelBond^x

Strain	Attachment (%)
Ck 70-15 ^y	38.8 $\pm$ 17.4 a ^z
E 2-3	55.8 $\pm$ 13.0 a
E 2-10	58.0 $\pm$ 7.8 a
M 2-2	17.7 $\pm$ 4.4 ab
M 2-9	9.7 $\pm$ 4.7 b
M 2-11	16.9 $\pm$ 5.4 b
M 2-12	10.8 $\pm$ 6.6 b

^x Conidial suspension was placed on GelBond. After incubation in a moisture chamber at room temperature for 2 h, all conidia in a marked area were counted with a light compound microscope. The GelBond was rinsed three times with a gentle stream of double-distilled H₂O from a squeeze bottle and conidia in the marked areas were recounted. Results are the combined data from two experiments, each with three replicates; mean $\pm$ standard deviation.

^y Ck = wild-type control strain. E = ectopic integration transformant. M = gene disrupted mutant.

^z The same letter indicates no significant effect of treatment at *P* = 0.05 as determined by Student's *t* test.

Table 3. Induction of appressorium formation (%) by *magB*[−] transformants by cAMP, or 1,16-hexadecanediol

Strain	GelBond ^y		Glass slide ^z	
	H ₂ O	10 mM cAMP	Ethanol	1 μ M Diol
Ck 70-15	12.1 $\pm$ 7.8	>95	0	>95
E 2-5	19.4 $\pm$ 3.5	93.4 $\pm$ 1.7	0	>95
E 2-10	...	...	0	>95
M 2-2	1.1 $\pm$ 0.7	62.1 $\pm$ 18.7	0	90.0 $\pm$ 5.0
M 2-9	2.1 $\pm$ 1.0	83.0 $\pm$ 10.7	0	75.0 $\pm$ 25.0
M 2-11	1.0 $\pm$ 0.6	76.0 $\pm$ 3.4	0	75.0 $\pm$ 25.0
M 2-12	0.8 $\pm$ 0.5	70.7 $\pm$ 3.7	0	75.0 $\pm$ 17.5

^y cAMP assay was conducted on GelBond; hydrophobic surface for gene disrupted mutants (M), and hydrophilic for controls: Ck = wild type; E = ectopic transformants (Lee and Dean 1993).

^z 1,16-hexadecanediol (Diol) assay was conducted on glass slides (Gilbert et al. 1996). Ethanol (5%) was used as the control for 1,16-hexadecanediol treatment. Results are the combined data from two experiments, each with three replicates; mean $\pm$ standard deviation. Over 100 conidia were counted for each replicate.

Fig. 6. cAMP (A) and 1,16-hexadecanediol (B) restore the ability to form appressorium (at 24 h) by *magB*[−] transformant (2-12) on the hydrophobic surface of GelBond. Magnification = $\times$ 200.

by the Taq DyeDeoxy terminator method (Amersham LifeScience, Arlington Heights, IL) and products separated on an ABI373 sequencer (Perkin-Elmer Applied Biosystems Division). Genomic clones of putative G α subunit genes were identified in a *pUI-NM554* cosmid library of *M. grisea* strain 70-15 (Lee and Dean 1993).

Nucleotide sequencing and alignments of *magA*, *magB*, and *magC* genes.

Cosmid clones corresponding to the three PCR clones were digested with *HindIII* and the appropriate fragments subcloned into *pBluescript II KS⁺*. A restriction map of each subclone was also constructed to localize the position of the putative

Fig. 7. Disease symptoms of *Magnaporthe grisea* on rice leaves after 7 days. Rice stem sheath was injected with 100 μ l of 1×10^5 conidia/ml or water, with a hypodermic needle and syringe. Inoculated plants were checked daily for symptom development; symptoms typically began to appear by 4 days. **A**, H₂O control; **B**, *magB⁻* transformant (2-12); **C**, *magB⁻* transformant (2-2); **D**, ectopic integration strain (2-10); **E**, Wild-type (70-15); **F**, *magB* rescued transformant (E2); **G**, *magB* rescued transformant (E4).

Fig. 8. Appressorium formation (at 24 h) on rice leaves. Ectopic integration strain (2-10) (**A**) or *magB⁻* transformant (2-12) (**B**) inoculated on the leaves from 2-week-old rice seedlings; *magB⁻* transformant (2-12) inoculated on an old leaf (**C**) or a young leaf (**D**) from a 4-week-old rice plant. Magnification = $\times 300$.

gene. The nucleotide sequence was determined and genes were named *magA*, *magB*, and *magC*, respectively. The data have been deposited at GenBank, with the following accession numbers: *magA*, AF011340; *magB*, AF011341; and *magC*, AF011342. On-line data base comparisons were performed with the BLAST algorithm (Altschul et al. 1990). Additional nucleotide and amino acid sequence comparisons and alignments were made with GeneJockey (Biosoft, Cambridge, UK) and GeneWorks (v2.3) (IntelliGenetics, Mountain View, CA) software packages.

Fig. 9. Southern blots of *magB* rescue. All genomic DNA was digested with *Hind*III and probed with a *Hind*III fragment of *magB* (4.3 kbp). Lane 1, wild-type (70-15); lane 2, ectopic integration strain (2-10); lane 3, *magB*⁻ transformant (2-12); lanes 5–8, transformants from 2-12 containing the *bar* gene only (P1, P2, P3, and P7); lanes 9–11, transformants from 2-12 containing *magB* and *bar* genes (E2, E4, and E6). As expected, the 4.3-kbp band reappeared in *magB* rescued transformants (E2, E4, and E6), and is lacking in *magB*⁻ transformant (2-12) and *magB*-transformants containing only the *bar* gene (P1, P2, P3, and P7).

Table 4. Vegetative growth, conidiation, and appressorium formation assay for strains restored for *magB*^x

Strain	Colony diameter (cm)		Conidia ($\times 10^4$)	Appress. (%)
	CM	V8		
Ck 70-15 ^y	3.6 $\pm$ 0.1 a ^z	4.9 $\pm$ 0.1 a	68.3 $\pm$ 14.1 a	>95 a
E 2-10	3.7 $\pm$ 0.1 a	4.9 $\pm$ 0.1 a	70.0 $\pm$ 10.5 a	>95 a
E 2-3	3.6 $\pm$ 0.1 a	4.9 $\pm$ 0.1 a	71.3 $\pm$ 7.8 a	>95 a
M 2-2	2.9 $\pm$ 0.1 b	4.1 $\pm$ 0.1 b	0.4 $\pm$ 0.2 b	2.2 $\pm$ 0.6 b
M 2-12	2.8 $\pm$ 0.1 b	4.0 $\pm$ 0.1 b	0.9 $\pm$ 0.4 b	2.5 $\pm$ 1.2 b
ME P1	2.8 $\pm$ 0.1 b	3.9 $\pm$ 0.1 b	0.6 $\pm$ 0.4 b	1.6 $\pm$ 0.5 b
ME P3	2.7 $\pm$ 0.1 b	3.9 $\pm$ 0.1 b	0.8 $\pm$ 0.4 b	0.4 $\pm$ 0.2 b
MR E2	3.6 $\pm$ 0.1 a	5.0 $\pm$ 0.1 a	64.7 $\pm$ 4.8 a	>95 a
MR E4	3.5 $\pm$ 0.1 a	4.9 $\pm$ 0.1 a	71.6 $\pm$ 10.4 a	>95 a

^x Colonies were grown on complete medium (CM) or V8 medium under fluorescent light for 7 days before colony diameters (cm) were measured. Conidia were collected from V8 plates and counted. Appressorium formation on the hydrophobic surface of GelBond was measured at 20 h. Data are from two independent experiments, each with two or three replicates; mean $\pm$ standard deviation.

^y Ck = wild-type control strain. E = ectopic integration transformant. M = gene disrupted mutant. ME = transformants of the *magB*⁻ strain 2-12 containing the *bar* gene alone. MR = transformants of the *magB*⁻ mutant 2-12 containing both *magB* and *bar* genes.

^z Different letters in each data column indicate significant differences at $P = 0.05$ as determined by Student's *t* test.

DNA and RNA extraction and manipulation.

Plasmid DNA was isolated by the standard alkaline lysis method (Sambrook et al. 1989). Restriction site mapping and subcloning were performed according to standard methods (Sambrook et al. 1989). For nucleotide sequencing, plasmid DNA was isolated with a Wizard DNA miniprep kit (Promega, Madison, WI). Fungal DNA was extracted from 4- to 5-day-old mycelia following the protocol of Mitchell and Dean 1995). RNA was extracted from mycelia or spores with the Qiagen RNeasy Plant mini kit (Qiagen, Chatsworth, CA). DNA digestion, agarose gel fractionation, RNA formaldehyde denatured gel electrophoresis, radiolabeling, and hybridization were according to the manufacturers' instructions and standard methods (Sambrook et al. 1989). Hybridization was carried out in 6 $\times$ SSC (1 $\times$ SSC is 0.15 M NaCl, 0.015 M sodium citrate), 5 $\times$ Denhardt's solution (1 $\times$ Denhardt's solution is 0.002% Ficoll 400, 0.02% polyvinylpyrrolidone [PVP], 0.02% bovine serum albumin [BSA]), 0.1% sodium dodecyl sulfate (SDS), and 50 mM phosphate buffer, pH 6.6, at 65 $^{\circ}$ C. Low-stringency hybridization was carried out at 55 $^{\circ}$ C. Membranes were washed down to between 2 $\times$ and 0.1 $\times$ SSC, 0.1 SDS, 55 to 65 $^{\circ}$ C, before being exposed to autoradiographic film.

Construction of gene replacement vectors *pB11pan* and *p8-8pan* and gene disruption vector *pAN2-3-6*.

Plasmid *pB11pan* was constructed by ligating a 3.6-kb *Stu*I-*Sph*I fragment from a partial digest of *pAN* 7-1 containing the complete hygromycin phosphor transferase (*hph*) gene (Punt et al. 1987) into *pB11-12*, digested with *Stu*I-*Sph*I. This resulted in the replacement of a 1.2-kb *Stu*I-*Sph*I fragment that contained the majority of the *magA* gene with the *hph* gene cassette.

Plasmid *p8-8pan* was constructed by first inserting a 2.3-kb *Xba*I fragment from the 5' end of *magC* into the *Xba*I site of *pAN* 7-1 to produce plasmid *p8pan*. After the correct orientation of the *Xba*I fragment in *p8pan* was confirmed, a 3.4-kb *Xho*I-*Bam*HI fragment from the 3' end of *magC* was ligated into *Xho*I- and *Bgl*II-digested *p8pan* to produce *p8-8pan*.

Plasmid *pAN2-3-6* (7.5 kbp) was constructed by inserting a 0.7-kb *Hind*III-*Xba*I PCR-amplified fragment of *magB* containing the amino half of the G α subunit. The PCR product was obtained with primers 2-3PF and 2-3PR (2-3PF GGGAAAGCTT⁴⁴GGAACGAGGAGATTGAGGAGAACC⁶⁴, 2-3PR 5' GGATCTAGA⁷⁶⁰AGG'TCGCCAATGATGAAAGT CG⁴³⁹) with the conditions described above, except the annealing temperature was raised to 55 $^{\circ}$ C.

Construction of plasmid for restoring *magB*.

Plasmid *pE12Bar* was constructed by inserting a 1.9-kb *Xba*I fragment containing the complete phosphinothricin acetyltransferase (*bar*) gene (Avalos et al. 1989) from *pBarKS1* (Pall and Brunelli 1993) into a unique *pE12* 2-3 *Xba*I that contained the complete *magB* gene.

Transformation of *M. grisea*.

Transformation protocols were adapted from established procedures (Leung et al. 1990; Sweigard et al. 1992). Conidia (5.0×10^6) were germinated in liquid CM overnight (18 h) at room temperature with agitation. Mycelium was harvested

with Miracloth (Calbiochem, San Diego, CA), rinsed, and wrung dry. The pellet was suspended in 30 ml of 1.0 M sorbitol to which 1 ml of Novozyme 234 (20 mg/ml; InterSpec Products, Foster City, CA) was added. After 90 min of agitation at room temperature, protoplasts were collected by filtering through cheesecloth and pelleted by centrifugation at $4,800 \times g$ for 10 min. Protoplasts were resuspended in STC (1.0 M sorbitol, 50 mM Tris-HCl, pH 8.0, and 50 mM CaCl_2) and were counted and adjusted to 5×10^7 cells/ml. Five micrograms of plasmid DNA (linearized DNA for gene replacement, or nonlinearized DNA for gene disruption or restoration) was mixed with 1.0×10^7 protoplasts in 200 μl of STC for 10 min before 1 ml of PTC (40% polyethylene glycol 4000 in STC) was added. After 20 min, 3 ml of TB3 (3 g of yeast extract, 3 g of casein acid hydrolysate, 10 g of glucose, and 200 g of sucrose per liter, containing 200 μg of hygromycin B or 500 μg of basta per ml) was added and incubated for 6 h at room temperature. Twenty milliliters of molten TB3 agar with hygromycin B (200 $\mu\text{g}/\text{ml}$) or basta (500 $\mu\text{g}/\text{ml}$) was mixed with 0.5 ml of treated protoplast mixture, and plated onto a petri dish. After culturing at 23°C in the dark for 5 to 7 days, hygromycin B or basta resistant clones were transferred to CM and V8 plates containing hygromycin B or basta. In CM plates containing basta, casamino acids were replaced with 0.5% L-proline. Conidia were streaked for single colonies.

Sexual crosses.

Crosses were performed by established procedures (Lau et al. 1993). Plugs of 0.5 cm^2 of 70-6 (MAT1-2), 70-15 (MAT1-1), and a strain of interest were placed 5 cm from each other on an oatmeal agar plate and incubated at $23 \pm 1^\circ\text{C}$ under fluorescent light. Between 14 and 28 days, asci were released from crushed perithecia on 2% distilled H_2O agar. Individual germinated asci were transferred to V8 agar plates and allowed to conidiate. Conidia were streaked for single colonies.

Vegetative growth and conidiation assays.

A 0.5- cm^2 agar block of *M. grisea* strains, grown on CM agar, was transferred onto CM or V8 agar, and grown under fluorescent light. Colony diameters were measured in centimeters at 5, 7, and 10 days. Conidia formed on V8 agar were harvested and counted with a hemacytometer. Each treatment contained 2 to 3 replicates, and the assays were conducted at least three times.

Appressorium formation assays.

Appressorium development was assayed by placing conidia on GelBond (Lee and Dean 1993; Mitchell and Dean 1995), or on rice leaf surfaces. Three 50- μl droplets of a conidial suspensions (1×10^5 conidia/ml) from 7- to 10-day-old cultures were placed on each piece of prewashed GelBond. After incubation in a moisture chamber at room temperature for a period of time, the germination and appressorium formation rates were scored with a light compound microscope (Mitchell and Dean 1995).

For observation of appressorium formation on the surface of rice leaves, freshly cut rice leaves (8 cm long) from 2- to 5-week-old rice plants were taped at their tips to glass slides and placed face up on a layer of moistened paper towel in a clear plastic box. Multiple 10- μl droplets of conidia (2×10^5 conidia/ml with 500 ppm Tween 20) were placed on the surface.

Conidia, germ tubes, and appressoria on the rice leaves were stained with cotton blue in lactophenol solution (5 ml of lactose, 5 ml of phenol, 15 ml of H_2O , and 10 mg of cotton blue) for 2 min, and counted with a light compound microscope. Germination and appressorium formation rates were checked periodically up to 72 h.

cAMP and 1,16-hexadecanediol treatments.

Fifty microliters of conidia suspension containing 10 mM cAMP or 1 μM 1,16-hexadecanediol was placed on the hydrophobic and hydrophilic surfaces of GelBond (controls only had H_2O or 5% ethanol) and allowed to germinate and form appressoria in a moisture chamber at room temperature (Lee and Dean 1993). Percent appressoria was recorded after 20 h of incubation.

Pathogenicity assays.

Conidia suspensions were prepared from 7-day-old V8 plates and adjusted to 1.0×10^5 conidia/ml in a 500-ppm Tween 20 solution. Approximately 8 ml was sprayed evenly onto 3-week-old rice plants (susceptible rice variety S-201), with a fine mist aspirator. The inoculated rice plants were placed in an unlit dew chamber at $25 \pm 2^\circ\text{C}$. After 24 h, a photoperiod of 15 h with fluorescent and incandescent lighting was resumed. Plants were rated for disease at 5 to 7 days after inoculation.

Conidia suspensions (100 μl) were also injected into the leaf with a 26 gauge, 1.27-cm-long needle and 3-ml syringe (Xu and Hamer 1996). After injection, the rice plants were covered with a plastic bag. After 24 h, the bag was removed and the plants were incubated for 5 to 7 days before being examined for disease.

ACKNOWLEDGMENTS

We are grateful to T. M. Wilkie and M. I. Simon for generously providing the degenerate oligonucleotide primers. We thank Robert D. Gilbert for conducting all diol induction assays, and Woo-Bong Choi for preparing the rice seedlings. This is technical contribution no. 4143 of the South Carolina Agricultural Experiment Station. This research was supported in part by National Science Foundation Grant nos. STI-9108772 and IBN-9219561 and U.S. Department of Agriculture National Research Initiative Grant no. 91-37303-6653 to R.A.D.

LITERATURE CITED

- Altschul, S. F., Gish, W., Willer, W., Myers, E. W., and Lipman, D. J. 1990. Blast local alignment search tool. *J. Mol. Biol.* 215:403-410.
- Avalos, J., Geever, R. F., and Case, M. E. 1989. Bialaphos resistance as a dominant selectable marker in *Neurospora crassa*. *Curr. Genet.* 16: 369-372.
- Beckerman, J. L., and Ebbole, D. J. 1996. *MPG1*, a gene encoding a fungal hydrophobin of *Magnaporthe grisea*, is involved in surface recognition. *Mol. Plant-Microbe Interact.* 9:450-456.
- Beckerman, J. L., Naider, F., and Ebbole, D. J. 1997. Inhibition of pathogenicity of the rice blast fungus by *Saccharomyces cerevisiae* α -factor. *Science* 276:1116-1119.
- Bourett, T. M., and Howard, R. J. 1990. *In vitro* development of penetration structures in the rice blast fungus *Magnaporthe grisea*. *Can. J. Bot.* 68:329-342.
- Brown, A., and Birnbaumer, L. 1990. Ionic channels and their regulation by G protein subunits. *Annu. Rev. Physiol.* 52:197-213.
- Buss, J. E., Mumby, S. M., Casey, P. J., Gilman, A. G., and Sefton, B. M. 1987. Myristoylated α subunits of guanine nucleotide-binding regulatory proteins. *Proc. Natl. Acad. Sci. USA* 84:7493-7497.
- Chen, B., Gao, S., Choi, G. H., and Nuss, D. L. 1996. Extensive altera-

- tion of fungal gene transcript accumulation and elevation of G-protein-regulated cAMP levels by a virulence-attenuation hypovirus. *Proc. Natl. Acad. Sci. USA* 93:7996-8000.
- Childers, S. R., and Deadwyler, S. A. 1996. Role of cyclic AMP in the actions of cannabinoid receptors. *Biochem. Pharmacol.* 52:819-827.
- Choi, G. H., Chen, B., and Nuss, D. L. 1995. Virus-mediated or transgenic suppression of a G-protein alpha subunit and attenuation of fungal virulence. *Proc. Natl. Acad. Sci. USA* 92:305-309.
- Coleman, D. E., and Sprang, S. R. 1996. How G proteins work: A continuing story. *TIBS* 21:41-44.
- Dean, R. A. 1992. Regulation of gene expression and signals in fungal development. Pages 3-26 in: *Molecular Signals in Plant-Microbe Communications*. D. P. S. Verma, ed. CRC Press, Boca Raton, FL.
- Dobinson, K. F., and Hamer, J. E. 1992. *Magnaporthe grisea*. Pages 67-86 in: *Molecular Biology of Filamentous Fungi: Proceedings of The EMBO-Workshop*, Berlin. U. Stahl and P. Tudzynski, eds. VCH, Weinheim, Germany.
- Ford, T. L., Cooley, J. T., and Christou, P. 1994. Current status for gene transfer into rice utilizing variety-independent delivery systems. Pages 195-208 in: *Rice Blast Disease*. R. S. Zeigler, S. A. Leong, and P. S. Teng, eds. CAB Int., Wallingford, UK.
- Gao, S., and Nuss, D. 1996. Distinct role for 2 G alpha subunits in fungal virulence, morphology, and reproduction revealed by targeted gene disruption. *Proc. Natl. Acad. Sci. USA* 93:14122-14127.
- Gilbert, R., Johnson, A., and Dean, R. A. 1996. Chemical signals responsible for appressorium formation in the rice blast fungus *Magnaporthe grisea*. *Physiol. Mol. Plant Pathol.* 48:335-346.
- Gilman, A. G. 1987. G proteins: Transducers of receptor-generated signals. *Annu. Rev. Biochem.* 56:615.
- Hamer, J., Howard, R. H., Chumley, F. G., and Valent, B. 1988. A mechanism for surface attachment in spores of a plant pathogenic fungus. *Science* 239:288-290.
- Horowitz, B. A. 1997. G protein α subunit genes of *Trichoderma larzianum* and *Cochliobolus heterostrophus*. (Abstr.) Page 87 in: *Fungal Genet. Conf.*, 19th.
- Ivey, F. D., Hodge, P. N., Turner, G. E., and Borkovich, K. A. 1996. The G alpha I homologue gna-1 controls multiple differentiation pathways in *Neurospora crassa*. *Mol. Biol. Cell.* 7:1283-1297.
- Lau, G. W., Chao, C. T., and Ellingboe, A. H. 1993. Interaction of genes controlling avirulence/virulence of *Magnaporthe grisea* on rice cultivar Katy. *Phytopathology* 83:375-382.
- Lee, Y.-H., and Dean, R. A. 1993. cAMP regulates infection structure formation in the plant pathogenic fungus *Magnaporthe grisea*. *Plant Cell* 5:693-700.
- Lee, Y.-H., and Dean, R. A. 1994. Hydrophobicity of contact surface induces appressorium formation of *Magnaporthe grisea*. *FEMS Microbiol. Lett.* 115:71-76.
- Lerea, C. L., Somers, D. E., Hurly, J. B., Klock, I. B., and Bunt-Mliam, A. H. 1988. Identification of specific transducin alpha subunits in retinal rod and cone photoreceptors. *Science* 234:77-80.
- Leung, H., Lehen, U., Karjalainen, R., Skinner, D., Tooley, P., Leong, S., and Ellingboe, A. 1990. Transformation of the rice blast fungus *Magnaporthe grisea* to hygromycin B resistance. *Curr. Genet.* 17:409-411.
- Mitchell, T. K., and Dean, R. A. 1995. The cAMP-dependent protein kinases catalytic subunit is required for appressorium formation and pathogenesis by the rice blast pathogen *Magnaporthe grisea*. *Plant Cell* 7:1869-1878.
- Ou, S. H. 1985. *Rice Disease*. 2nd ed. Commonw. Mycol. Inst., Surrey, UK.
- Pall, M. L., and Brunelli, J. P. 1993. A series of six compact fungal transformation vectors containing polylinkers with multiple unique restriction sites. *Fungal Genet. Newsl.* 40:59-62.
- Punt, P. J., Oliver, R. P., Dingemanse, M. A., Pouwels, P. H., and van den Hondel, C. A. M. J. J. 1987. Transformation of *Aspergillus* based on the hygromycin B resistance marker from *Escherichia coli*. *Gene* 56:117-124.
- Regenfelder, E., Spellig, T., Hartmann, A., Lauenstein, S., Bolker, M., and Kahmann, R. 1997. G proteins in *Ustilago maydis*: Transmission of multiple signals? *EMBO J.* 16:1934-1942.
- Sambrook, J., Fritsch, E. F., and Maniatis, T. A. 1989. *Molecular Cloning: A Laboratory Manual*. 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Shi, Z., and Leung, H. 1994. Genetic analysis and rapid mapping of a sporulation mutation in *Magnaporthe grisea*. *Mol. Plant-Microbe Interact.* 7:113-120.
- Simon, M. I., Strathmann, M. P., and Gautam, N. 1991. Diversity of G proteins in signal transduction. *Science* 252:802-808.
- Spiegel, A. M., Backlund, P. S., Butrynski, J. E., Jones, T. L. Z., and Simonds, W. F. 1991. The G protein connection: Molecular basis of membrane association. *Trends Biochem. Sci.* 16:338-341.
- Strathmann, M., Wilkie, T. M., and Simon, M. I. 1989. Diversity of the G-protein family: Sequences from five additional alpha subunits in the mouse. *Proc. Natl. Acad. Sci. USA* 86:7407-7409.
- Stryer, L., and Bourne, H. R. 1986. G protein: A family of signal transducers. *Annu. Rev. Cell Biol.* 2, 391-419.
- Sweigard, J. A., Chumley, F. G., and Valent, B. 1992. Disruption of a *Magnaporthe grisea* cutinase gene. *Mol. Gen. Genet.* 232:183-190.
- Talbot, N. J., Ebbole, D. J., and Hamer, J. E. 1993. Identification and characterization of MPG1, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *Plant Cell* 5:1575-1590.
- Teng, P. S. 1994. The epidemiological basis for blast management. Pages 409-434 in: *Rice Blast Disease*. R. S. Zeigler, S. A. Leong, and P. S. Teng, eds. CAB Int., Wallingford, UK.
- Timberlake, W. E., and Marshall, M. A. 1989. Genetic engineering of filamentous fungi. *Science* 244:1313-1317.
- Tolkacheva, T., McNamara, P., Piekarz, E., and Courchesne, W. 1994. Cloning of a *Cryptococcus neoformans* gene, GPA1, encoding a G-protein alpha-subunit homolog. *Infect. Immun.* 62:2849-2856.
- Turner, G. E., and Borkovich, K. A. 1993. Identification of a G protein alpha subunit from *Neurospora crassa* that is a member of the Gi family. *J. Biol. Chem.* 268:14805-14811.
- Valent, B. 1990. Rice blast as a model system for plant pathology. *Phytopathology* 80:33-36.
- West, R. E., Jr., Moss, J., Vaughan, M., Liu, T., and Liu, T.-Y. 1985. Pertussis toxin-catalyzed ADP-ribosylation of transducin. Cysteine 347 is the ADP-ribose acceptor site. *J. Biol. Chem.* 260:14428-14430.
- Wilkie, T. M., and Yokoyama, S. 1994. Evolution of the G protein alpha subunit multigene family. Pages 249-270 in: *Molecular Evolution of Physiological Processes*. Rockefeller University Press, New York.
- Xu, J.-R., and Hamer, J. E. 1996. MAP kinase and cAMP signaling regulate infection structure formation and pathogenic growth in rice blast fungus *Magnaporthe grisea*. *Gene Dev.* 10:2696-2706.
- Yatani, A., Imoto, Y., Codina, J., Hamilton, S. L., Brown, A. M., and Birnbaumer, L. 1988. The stimulatory G protein of adenylyl cyclase, Gs, also stimulates dihydropyridine-sensitive Ca²⁺ channels. Evidence for direct regulation independent of phosphorylation by cAMP-dependent protein kinase or stimulation by a dihydropyridine agonist. *J. Biol. Chem.* 263:9887-9895.
- Yu, J.-H., Wieser, J., and Adams, T. H. 1996. The *Aspergillus* FlbA RGA domain protein antagonizes G protein signaling to block proliferation and allow development. *EMBO J.* 15:5184-5190.