Population Genetic Analysis of *Puccinia striiformis* f. sp. *tritici* Suggests Two Distinct Populations in Tibet and the Other Regions of China

Xiaoping Hu and **Lijie Ma**, State Key Laboratory of Crop Stress Biology for Arid Areas, College of Plant Protection, Northwest A&F University, Yangling, Shaanxi 712100, China; **Taiguo Liu**, State Key Laboratory for the Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Science, Beijing 100193, China; **Conghao Wang**, College of Plant Protection, Northwest A&F University, Yangling, Shaanxi 712100, China; **Yuelin Peng**, Tibet Agricultural and Animal Husbandry College, Linzhi 860000, China; **Qiong Pu**, Tibet Autonomous Region Agricultural Technology Extension and Service Center, Lhasa 850000, Tibet; and **Xiangming Xu**, NIAB East Malling Research, New Road, East Malling, ME19 6BJ, Kent, U.K.

Abstract

Wheat stripe rust, caused by *Puccinia striiformis* f. sp. *tritici* (*Pst*), is an important disease on wheat, seriously threatening wheat production worldwide. China is one of the largest stripe rust epidemic regions in the world. The pathogen sexual reproduction and migration routes between Tibet and the other regions in China are still unknown. In this study, we obtained 961 *Pst* isolates from 1,391 wheat leaf samples from Gansu (277), Shaanxi (253), Sichuan (172), and Tibet (259), comprising 13 natural populations, and genotyped them with simple sequence repeat (SSR) markers. The isolates can be divided into two distinct clusters based on DAPC and STRUCTURE analyses. The genetic diversity of Longnan

(in Gansu) and Yibin (in Sichuan) populations was the highest and lowest among the 13 populations, respectively. The hypothesis of multilocus linkage disequilibrium was rejected for the populations from Linzhi in the Himalayan, Longnan, Hanzhong, Guangyuan, Mianyang, Liangshan, and Chendu in the south Qinling Mountains at the level of $P = 0.01$, which indicated significant linkage among markers in these populations. Populations in the other regions had extensive gene exchange ($N_m > 4$); little gene exchange was found between Tibet and the other regions ($N_m < 1$). The results suggest that the Tibet epidemic region of *Pst* is highly differentiated from the other epidemic regions in China.

Wheat stripe rust, caused by the basidiomycete fungus *Puccinia striiformis* Westend. f. sp. *tritici* Eriks. (*Pst*), is one of the most damaging diseases on wheat (*Triticum aestivum*) worldwide (Chen 2005; Hovmøller et al. 2010, 2011). It is a globally distributed and wind-dispersed pandemic disease. Urediniospores of *Pst* can be dispersed by atmospheric circulation up to 1,000 km, leading to rust spreading across large areas within a single growing season (Zeng and Luo 2006). *Pst* is a plant pathogen that evolves rapidly and can cause serious yield losses. In China, wheat stripe rust caused yield losses of 6.00, 3.20, 2.65, and 1.40 million metric tons when severe epidemics occurred in 1950, 1964, 1990, and 2002, respectively (Li and Zeng 2002; Wan et al. 2004).

The epidemics of wheat stripe rust in China can be subdivided into three epidemiological regions based on geography, climatic conditions, agronomy, *Pst* race composition, and epidemic characteristics (Hu et al. 2012; Li and Zeng 2002; Zeng and Luo 2006): the main epidemiological region (geographic ranges of 87 to 135°E and 22 to 53°N, covering Gansu, Shaanxi, Sichuan, Hubei, Henan, Yunnan, and Guizhou provinces), the Xinjiang epidemiological region, and the Tibet epidemiological region (Fig. S1). Dispersal of *Pst* inoculum within the main epidemiological region has been studied based on climatic conditions (Wang et al. 2010a) and fungal population structure at the molecular level (Chen 2008; Lu et al. 2011; Shan et al. 1998). However, fungal population genetic structure in the Tibet epidemiological region and its genetic relationship with the other epidemic regions are still unknown.

Several studies revealed an overall clonal population structure of *Pst* with low genetic diversity worldwide (Chen et al. 1993; Cheng

and Chen 2014; Enjalbert et al. 2005; Hovmøller et al. 2002; Wellings 2007), except in China, Nepal, and Pakistan (Ali et al. 2014a,b; Duan et al. 2010; Mboup et al. 2009; Zhao et al. 2013). However, Duan et al. (2010) demonstrated that Chinese *Pst* isolates from Gansu Province exhibited high genetic diversity and evidence of recombination. A very recent worldwide population genetic structure analysis of *Pst* also indicated a high regional heterogeneity in levels of recombination, with clear signatures of recombination in the Himalayan and near-Himalayan regions (Ali et al. 2014b). However, the sexual reproduction of *Pst* in China remains unknown although Zhao et al. (2013) demonstrated that *Pst* can infect some *Berberis* spp. through greenhouse work, which may lead to sexual recombination on *Berberis* spp.

The information on factors affecting fungal population genetic structure is critical if we are to develop disease management strategies based on the knowledge of the patterns of pathogen migration among regions. This paper reports a study comparing *Pst* populations from Tibet and other regions (excluding the Xinjiang region) using SSR markers to infer *Pst* population biology, migration routes, and potential sexual reproduction.

Materials and Methods

Sampling *Pst*-infected leaves and multiplying urediniospores.

A total of 1,391 *Pst* samples were randomly collected from adult plants of 96 fields in Gansu (347), Sichuan (349), Shaanxi (200), and Tibet (495) provinces in 2009 and 2010. These sampling areas were classified into 13 geographical regions (Table 1). Leaves with fresh uredinia pustules but without signs of other pathogens were sampled. Leaf surfaces were gently dried with a piece of absorbent paper, enclosed in an envelope, and kept inside a desiccator at room temperature for not more than 7 days before inoculum multiplication. To produce fresh urediniospores on those sampled leaves, the leaves were washed with water, placed on water-soaked tissue paper in petri dishes, and then incubated in the dark for 12 h at 10 to 13°C. Urediniospores were then collected to inoculate seedlings of cv. Mingxian 169 (a cultivar highly susceptible to all known races of *Pst* identified in China) for multiplication.

Ten seeds were sowed in a 10 cm diameter pot, containing cow dung compost and soil (1:2, v/v). When the first leaf of seedlings had fully expanded (approx. 8 to 10 days after sowing), a single urediniospore from a freshly sporulating lesion on the sampled leaf was

Corresponding author: X. Hu; E-mail: xphu@nwsuaf.edu.cn

X. P. Hu, L. J. Ma, and T. G. Liu contributed equally to this study.

*The e-Xtra logo stands for “electronic extra” and indicates that three supplementary figures are available online.

Accepted for publication 12 October 2016.

picked up with a sterile insect needle under a microscope (Olympus BX51, Japan) at 200× and transferred onto a leaf. A single seedling received a single urediniospore. Immediately after inoculation, seedlings were kept in a dew chamber at 10°C for 24 h and then placed on benches in a glasshouse compartment with the temperature set to 16:14°C (day/night) and a photoperiod of 10:14 h (day/night). When a lesion became visible on each inoculated leaf, seedlings inoculated with spores from the same lesion on each sampled leaf were separated from other seedlings with a plastic cylinder to prevent cross contamination. Fresh urediniospores produced on these inoculated seedlings were collected separately, vacuum dried, and then stored at –80°C for future use.

After collecting and multiplying urediniospores, we successfully recovered 961 samples from the 1,391 samples, including 702 samples from Gansu, Shaanxi, and Sichuan provinces collected in June 2009, and 259 samples from Tibet collected in June 2010 (Table 1).

DNA extraction. DNA was extracted directly from urediniospores of each isolate using the CTAB method (Wang et al. 2010b). A sample of 10 mg fresh urediniospores were placed into a 1.5-ml centrifuge tube and ground by electric drill on ice. Thirty microliters of lysis buffer containing 20% SDS, 20 µl of 5 mol/liter NaCl, and 30 µl of CTAB/NaCl were added into each tube. The tubes were then incubated in a 65°C water bath for 1 h, and inverted every 10 min. Next, 400 µl of phenol/chloroform/isoamyl alcohol (25:24:1, pH > 7.8) was added to each tube; the tubes were shaken gently and then centrifuged at 12,500 rpm for 10 min at 4°C. The aqueous phase was collected and DNA was precipitated with equal volume of isopropyl alcohol at –20°C for 1 to 2 h. The tubes were centrifuged at 12,500 rpm for 15 min at 4°C, the supernatant was discarded, and DNA pellet was washed twice with 70% ethanol, air-dried, and dissolved in 30 µl TE solution. DNA was purified by adding 4 µl RNase A (10 mg/ml) and

incubated for 1 h at 37°C. DNA concentration was then measured with a NanoDrop 2000 spectrophotometer (Thermal Co., U.S.A.), and DNA integrity was checked using agarose gel electrophoresis in a 1.0% agarose gel and visualized under AlphaImager System (CA, U.S.A.) after staining in an ethidium bromide solution (0.5 µg/ml) for 15 min. DNA was diluted to 50 ng/µl and stored at –80°C for future use.

PCR amplification and electrophoresis of SSR markers. Ten pairs of published SSR primers were used to genotype each isolate (Table 2): RJ18, RJ20, RJ21, and RJ24 were used as described in Enjalbert et al. (2002); CPS08, CPS09, CPS10, CPS15, CPS34, and CPS36 were used according to Chen et al. (2009). All SSR primers were synthesized by Life Technologies co., Shanghai, China. The PCR reaction system consisted of 25 µl: 10× reaction buffer 2.5 µl, MgCl₂ (25 mM) 1.7 µl, dNTP (2.5 mM) 2.0 µl, forward primer (10 µM) 0.5 µl, reverse primer (10 µM) 0.5 µl, Taq polymerase (5 U/µl) 0.2 µl, DNA (50 ng/µl) 1.0 µl, ddH₂O 16.6 µl. The amplification conditions were as the following: a predenaturation step at 94°C for 4 min followed by 10 amplification cycles consisting of denaturation at 94°C for 45 s, annealing at 64°C for 45 s (the temperature decreased by 0.7°C for every cycle), and extension at 72°C for 45 s, with another 25 cycles of denaturation at 94°C for 45 s, annealing at 54°C for 45 s, extension 72°C for 45 s, a final extension step at 72°C for 10 min.

PCR products were separated on a 6% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) in a sequence electrophoresis tank (DYCZ-20C, Beijing, China). The SDS-PAGE with electrophoresis glass plate was soaked in a 10% acetic acid and gently shaken until color indicator disappeared (approx. 30 min), washed for 10 min in ddH₂O twice, transferred into silver-staining solution (1.7 g AgNO₃ and 1,700 ml ddH₂O) and gently shaken for 30 min, washed with

Table 1. Information of *Puccinia striiformis* f. sp. *tritici* samples collected from different regions in China

Province	Geographical region	County	No. of fields	No. of samples collected	No. of samples recovered	Altitude (m)
Gansu	Tianshui	Qinzhou	5	347	72	1,320–1,750
		Qinan	3		33	
		Maiji	3		36	
	Pingliang	Kongtong	4		49	1,100–1,600
	Longnan	Wenxian	4		56	
		Wudu	3		31	
Subtotal			22		277	
Sichuan	Aba	Jiuzhaigou	3	349	24	2,850–3,000
		Songpan	4		24	
		Heishui	3		23	
	Liangshan	Dechang	3		16	1,100–1,700
		Huili	2		42	
	Guangyuan	Yuanba	3		9	1,000–1,950
		Jiange	3		22	
		Zitong	3		18	
	Mianyang	Jintang	2		28	500–1,800
	Chengdu	Yibin	3		7	450–900
		Pingshan	3		30	
		Cuiping	2		10	
Subtotal			34		253	
Shaanxi	Baoji	Longxian	3	200	26	700–1,100
		Chencang	3		20	
		Qianyang	4		20	
		Fenxiang	2		28	
	Hanzhong	Ningqiang	3		26	300–550
		Mianxian	3		30	
	Ankang	Hanbin	2		22	800–1,900
Subtotal			20		172	
Tibet	Linzhi	Miling	7	495	94	2,450–3,760
		Linzhi	5		69	
		Bomi	5		91	
		Gongbujiangda	3		5	
Subtotal			20		259	
Total			96	1,391	961	

ddH₂O, soaked into a developer solution (40 g NaOH, 400 µl ammonium persulfate, 3 ml formaldehyde, and 2,000 ml ddH₂O) until bands of desired intensity appeared (approx. 10 min). Then, the gel was immersed in a 5% acetic acid solution for about 5 min. The SDS-PAGE was dried and scanned into computer using a scanner. The bands were scored using automatic molecular weight calculation based on standard molecular ladder under the AlphaView Software (Alpha Innotech, CA, U.S.A.). The experiment was repeated twice. Only stable bands were recorded. To verify the motifs, 50 SSR products (representing a range of sizes for each SSR marker) were randomly selected and sequenced with an ABI 373A automated DNA sequencer (Life Technologies co., Shanghai, China).

Amplicon size categories at each locus were considered unique alleles that were compiled across loci into multilocus genotypes (MLGs). Isolates with identical MLGs in the same population were considered clonets (Tibayrenc and Ayala 2002, 2012). Data from only one representative of each clonet was incorporated into clone-corrected datasets for analysis using GenoDive 2.0 beta version (Meirmans and Hedrick 2011) and Multilocus version 3.1 (Agapow and Burt 2001).

Analysis of population subdivision. We used multivariate and model-based Bayesian clustering approaches to infer population subdivision. Multivariate discriminant analyses of principal components (DAPC), implemented in the ADEGENET package in R environment (Jombart 2008), were performed to identify partitions within the dataset, regardless of the geographic origin of samples. This method makes no assumption regarding data structure or underlying population genetics model. The *k*-means were used to run sequentially with increasing values of *k* (number of clusters), and different clustering solutions were compared using Bayesian Information Criterion (BIC) to identify the optimal number of clusters (Jombart et al. 2010).

In addition to nonparametric analysis (i.e., DAPC), a model-based Bayesian method implemented in STRUCTURE 2.2 (Falush et al. 2003; Pritchard et al. 2000) was used to identify genetic clusters and to evaluate the extent of admixture among them. A model allowing admixture and independent allele frequencies among populations was used. For each simulated cluster *K* (*K* = 1–20), 50 runs were performed with 400,000 iterations in Monte Carlo Markov Chain replications and a burn-in period of 100,000. The outputs were processed with CLUMPP 1.1.2 (Jakobsson and Rosenberg 2007) and *G'*-statistic >80% (Nei 1973) was used as a criterion to assign runs into a common clustering pattern. STRUCTURE output was used as an input file to STRUCTURE HARVESTER (<http://taylor0.biology.ucla.edu/structureHarvester>) to estimate the optimal number of clusters (*K*). The ad hoc statistic ΔK , which is based on the rate of change in

the log probability, was calculated to identify the optimal number of clusters (*K*) (Evanno et al. 2005).

Analysis of genetic variation within and among clusters. Two classifications were used to group strains for within- and between-group analysis. The first one is the genetic clusters identified by DAPC and STRUCTURE; the second was the geographic location. Clonal fractions (CF) were calculated as:

$$CF = 1 - [(\text{number of different MLGs})/(\text{total number of isolates})]$$

Allele frequencies, the number of alleles per locus, and gene diversity (*H*) were calculated using POPGENE 1.32 (Yeh et al. 1999). Allelic richness (AR) and private allelic richness (PAR) were calculated using ADZE (Szpiech et al. 2008), which uses a rarefaction procedure to adjust variation in sample size across populations. AR is the average number of alleles per locus corrected for sample size (Dilmaghani et al. 2012). PAR is the number of unique alleles exclusive to single populations (Dilmaghani et al. 2012; Szpiech et al. 2008). Gene diversity was calculated for each group of isolates, which estimates the probability that two alleles chosen at random from a given population are different (Nei 1973, 1978). Pairwise measures of differentiation between clusters were estimated from a corrected standardized fixation index (G''_{ST}) using GENODIVE 2.0 beta version (Hedrick 2005; Meirmans and Hedrick 2011). Gene flow (N_m , number of migrants) between *Pst* populations was estimated as: $N_m = 0.25 \times (1 - G''_{ST})/G''_{ST}$. $N_m < 1$ and ≥ 1 imply little and considerable amount of gene flow between populations, respectively (Culley et al. 2002; Wright 1951).

Linkage disequilibrium analysis. The linkage disequilibrium (LD) tests were performed for each cluster (or subpopulation) separately. Only clusters and subpopulations with at least 10 clone corrected isolates were used for LD analysis. The standardized index of association (rBarD), an alternative measure of the index of association (I_A) adjusted for the number of loci, was used to assess the significance of LD using Multilocus version 3.1 (Agapow and Burt 2001). The rBarD values range from 0 to 1; an rBarD value of 0 corresponds to a random association of alleles at different loci (linkage equilibrium), whereas an rBarD value of 1 indicates a complete association of alleles at different loci (complete linkage disequilibrium) (Agapow and Burt 2001). The significance of rBarD was assessed with a randomization test. Under the null hypothesis of rBarD = 0, 1,000 sets of data were randomly drawn from the data, representing 1,000 realizations under the null hypothesis; rBarD was calculated for each realization. Statistical significance of the observed rBarD value was assessed by its comparison with the 1,000 rBarD values generated under the rBarD = 0 hypothesis.

Table 2. SSR primer pairs of *Puccinia striiformis* f. sp. *tritici* used in this study

Primer name	Primer sequence (5'–3')	Repeat motif	Annealing temperature (°C)	Expected range of fragment sizes (bp)	No. of alleles	Reference
CPS08F	GATAAGAAACAAGGACAGC	(CAG) ₁₄	55	221–227	3	Chen et al. 2009
CPS08R	CAGTGAACCAATTACTCAG					
CPS09F	CGGGAGAAGACCTGAGC	(GTT) ₉	58	379–407	9	Chen et al. 2009
CPS09R	AGAAAACGGAATGTAATGTG					
CPS10F	TCTACTGGGCAGACTGGTC	(TAG) ₈	56	335–341	4	Chen et al. 2009
CPS10R	CGGTTTGTTCGCTTC					
CPS15F	GATGGGGAAAAGTAAGAAGT	(TTC) ₄	58	236–245	3	Chen et al. 2009
CPS15R	GGTGGGGATGTAAGTATGTA					
CPS34F	GTTGGCTACGAGTGGTCATC	(TC) ₉	58	131–138	3	Chen et al. 2009
CPS34R	TAACACTACACAAAAGGGGTC					
CPS36F	TCCAGGCAGTAAATCAGACGC	(GCA) ₆	54	145–149	2	Chen et al. 2009
CPS36R	ATCAGCAGGTGTAGCCCCATC					
RJ18F	CTGCCCATGCTCTTCGTC	(TGT) ₅	56	220–376	4	Enjalbert et al. 2002
RJ18R	GATGAAGTGGGTGCTGCTG					
RJ20F	AGAAGATCGACGCACCCG	(CAG) ₄	56	300–309	4	Enjalbert et al. 2002
RJ20R	CCTCCGATTGGCTTAGGC					
RJ21F	TTCCTGGATTGAATTCGTCG	(GTT) ₆	58	200–226	2	Enjalbert et al. 2002
RJ21R	CAGTTCTCACTCGGACCCAG					
RJ24F	TTGCTGAGTAGTTTGGCGGTGAG	(GTT) ₅₊₉	54	287–315	9	Enjalbert et al. 2002
RJ24R	CTAAGCCCATCCTCCAACC					

cluster 1 contained isolates from Qinzhou, Qinan, Maiji, and Kongtong in Gansu Province; cluster 2 contained isolates from Wenxian and Wudu in Gansu Province; Heishui, Dechang, Huili, Yibin, Pingshan in Sichuan Province; Chengcang, Mianxian, and Hanbin in Shaanxi Province; cluster 3 contained all the other isolates.

Nonparametric DAPC also suggested two clusters (Fig. 2) though the support is not as strong as the STRUCTURE analysis (Fig. 1). Cluster 2 and 1 comprised mainly isolates from the Tibet region and the other regions, respectively, consistent with the clusters derived from the STRUCTURE analysis (Table 3). DAPC grouped four isolates from Tibet with isolates from the other regions, compared with such a classification for only one isolate by STRUCTURE analysis. The 700 isolates in cluster 1 comprised three subclusters (Fig. 3). Subcluster 1 consisted mainly of those isolates from Mianyang, Pingliang, Baoji, Ankang, and Hanzhong; subcluster 2 mainly of isolates from Longnan, Yibin, Liangshan, and Ankang; subcluster 3 mainly of isolates from Guangyuan, Chengdu, and Tianshui. Isolates from Tianshui, Pingliang, Longnan, Baoji, Hanzhong, Ankang, Guangyuan, Mianyang, Aba, Chengdu, Liangshan, and Yibin were distributed into three subclusters (Fig. 4).

Genetic diversity. Table 4 gives clonal fractions (CF), gene diversity (H), and allelic richness (AR) and private allelic richness (PAR) for the two DAPC clusters. The clonal fraction was 0.30 for cluster 1 ($N = 700$) and 0.26 for cluster 2 ($N = 261$). The CF of all *Pst* isolates was 0.29. The gene diversity was 0.307 for cluster 1 and 0.336 for cluster 2. The allelic richness was 1.749 for cluster 1 and 1.830 for cluster 2; the corresponding value of private allelic richness was 0.153 and 0.235, respectively.

Cluster 1 included 12 subpopulations (Tianshui, Pingliang, Longnan, Aba, Liangshan, Gyuangyuan, Mianyang, Chengdu, Yibin,

Baoji, Hanzhong, and Ankang). CF values of these subpopulations ranged from 0.11 for Longnan ($N = 58$) to 0.74 for Yibin ($N = 74$) (mean: 0.338). H values ranged from 0.179 for Liangshan ($N = 58$) to 0.307 for Tianshui ($N = 140$) (mean: 0.243) (Table 4). AR values ranged from 1.514 for Liangshan to 1.843 for Longnan (mean: 1.697); PAR values were close to 0 (Table 4) for all 12 subpopulations. Cluster 2 included Linzhi subpopulation only; CF, H , AR, and PAR values were 0.26, 0.336, 1.830, and 0.235, respectively.

The relationship between population differentiation (G''_{ST}) and geographical distance was approximately linear ($R^2 = 0.626$) (Fig. 5). The distance between the Linzhi subpopulation and each of Tianshui, Pingliang, Longnan, Aba, Liangshan, Gyuangyuan, Mianyang, Chengdu, Yibin, Baoji, Hanzhong, and Ankang subpopulation is >1,500 km; the G''_{ST} values were all >0.3. The distances among Tianshui, Pingliang, Longnan, Aba, Liangshan, Gyuangyuan, Mianyang, Chengdu, Yibin, Baoji, Hanzhong, and Ankang subpopulation are all <1,500 km; the G''_{ST} values were all <0.3 (Fig. 5).

Population differentiation. Corrected standardized population differentiation (G''_{ST}) values of two DAPC clusters indicated that the two clusters were highly differentiated, with G''_{ST} values of 0.323 ($P < 0.001$). At the subpopulation level, most G''_{ST} values between subpopulations from different clusters were greater than zero, namely P13 (Linzhi from cluster 2) with other 12 subpopulations from cluster 1 ($P < 0.001$) (Table 5). There was no significant difference ($P > 0.05$) between Pingliang subpopulation (P2) and each of (Mianyang) P7, (Chengdu) P8, (Yibin) P9, (Baoji) P10, and (Ankang) P12 subpopulation; Tianshui (P1) and Mianyang (P7); Longnan (P3) and Ankang (P12); Liangshan (P5) and Yibin (P9); Guangyuan (P6) and each of Mianyang (P7) and Chengdu (P8); Mianyang (P7) and each of Chengdu (P8), Yibin (P9), Baoji (P10), and

Table 3. Classification of 961 strains of *Puccinia striiformis* f. sp. *tritici* using STRUCTURE version 2.2.3 (Pritchard et al. 2000) and the discriminant analysis of principal components (DAPC) (Jombart et al. 2010)

Method	$K = 2$				$K = 3$					
	Cluster 1		Cluster 2		Cluster 1		Cluster 2		Cluster 3	
	Individual number from main epidemic region/% ^a	Individual number from Tibet epidemic region/% ^b	Individual number from main epidemic region/% ^a	Individual number from Tibet epidemic region/% ^b	Individual number from main epidemic region/% ^a	Individual number from Tibet epidemic region/% ^b	Individual number from main epidemic region/% ^a	Individual number from Tibet epidemic region/% ^b	Individual number from main epidemic region/% ^a	Individual number from Tibet epidemic region/% ^b
STRUCTURE	702 / 100.0	0 / 0.0	0 / 0.0	259 / 100.0	296 / 42.2	0 / 0.0	395 / 56.3	2 / 0.8	11 / 1.6	257 / 99.2
DAPC	696 / 99.1	4 / 1.5	6 / 0.9	255 / 98.5	288 / 41.0	0 / 0.0	410 / 58.4	4 / 1.5	4 / 0.6	255 / 98.5

^a The proportion was calculated using the individual number of the cluster and total individual number of the main epidemic region.

^b The proportion was calculated using the individual number of the cluster and total individual number of the Tibet epidemic region.

Fig. 3. Discriminant analysis of principal components (DAPC) (Jombart et al. 2010) on 702 *Puccinia striiformis* f. sp. *tritici* (*Pst*) genotypes from the main epidemic region. **A**, Value of Bayesian information criterion (BIC) versus number of clusters for 702 *Pst* genotypes from the main epidemic region. **B**, Scatter-plot of the DAPC on a set of 702 *Pst* genotypes from the main epidemic region. At the bottom right, the eigenvalues of the first two axes are represented. Genotypes are represented by dots and clusters as ellipses.

Ankang (P12); Chengdu (P8) and Yibin (P9); Baoji (P10) and Ankang (P12); and Hanzhong (P11) and Ankang (P12) (Table 5).

Linkage disequilibrium. The clusters determined by DAPC analysis were used for multilocus linkage disequilibrium (LD) analysis.

Fig. 4. Subclusters based on the discriminant analysis of principal components (DAPC) (Jombart et al. 2010) performed on 702 individuals of *Puccinia striiformis* f. sp. *tritici* genotypes from main epidemic region in China.

The LD value was significantly greater than 0 for both clusters ($P = 0.01$). However, subpopulations from Longnan, Liangshan, Mianyang, Chengdu, and Hanzhong in cluster 1 had LD values that were not statistically different from 0 ($P > 0.05$) (Table 4; Fig. S3).

Gene flow of *Pst* among populations. Populations in the other regions including Gansu, Shaanxi, and Sichuan had extensive gene exchanges as N_m were 6.17, 7.55, and 4.71 for Gansu/Sichuan, Gansu/Shaanxi, and Sichuan/Shaanxi, respectively; whereas low gene flows were detected between Tibet and other provinces (Gansu, Shaanxi, and Sichuan): N_m values were all close to 0.4.

Discussion

The *Pst* epidemiology in the epidemic regions except Tibet and Xinjiang in China had been well studied (Hu et al. 2000; Li and Liu 1957; Lu et al. 2011; Shen et al. 2008; Wang et al. 2010a,b; Xie et al. 1986, 1988; Zeng and Luo 2006, 2008). Zeng and Luo (2006) proposed 15 different epidemic subregions in the main epidemiological region in China based on geographic, climatic, cropping, and rust epidemic features. In the main *Pst* epidemiological region, the genetic diversity was relatively high and the majority of genetic variation occurred within populations. The Longnan population had the highest genetic diversity in the main epidemiological region in China (Chen 2008; Lu et al. 2011). In contrast, few reports were on the Xinjiang and Tibet epidemic regions. The *Pst* population in Xinjiang had a higher genetic diversity than populations in Qinghai, Gansu, Ningxia, and Tibet. Twenty-seven previously known and eight unknown races were detected in Xinjiang, higher than in other epidemiological regions (Zhan et al. 2016). The composition of

Table 4. Genetic diversity of *Puccinia striiformis* f. sp. *tritici* populations analyzed with 43 microsatellite loci

Population		<i>Berberis</i> sp. (Zhao et al. 2013; Wang et al. 2016)	Sample size	Clone corrected	Clonal fraction ^a	Genotypic richness	Gene diversity (H^b)	Allelic richness ^c	Private allelic richness ^c	$rBarD^d$
Cluster 1	Tianshui	<i>B. brachypoda</i>	140	119	0.15	0.85	0.307 (± 0.19)	1.773 (± 0.08)	0	0.1757**
		<i>B. aggregate</i>								
	Pingliang	NI ^e	49	39	0.21	0.79	0.254 (± 0.18)	1.735 (± 0.09)	0	0.054**
	Longnan	<i>B. aggregate</i>	86	77	0.11	0.89	0.301 (± 0.16)	1.843 (± 0.06)	0	0.005797
		<i>B. soulieana</i>								
		<i>B. stenostachya</i>								
	Aba	NI	70	50	0.29	0.71	0.257 (± 0.19)	1.718 (± 0.08)	0.001 $\pm$ 0.001	0.02643**
	Liangshan	NI	58	27	0.54	0.46	0.179 (± 0.21)	1.514 (± 0.10)	0	0.01584
	Guangyuan	<i>B. soulieana</i>	31	24	0.23	0.77	0.263 (± 0.19)	1.753 (± 0.09)	0	0.01763
	Mianyang	<i>B. soulieana</i>	18	10	0.47	0.53	0.197 (± 0.17)	1.700 (± 0.11)	0	0.02573
	Chengdu	<i>B. atrocarpa</i>	28	21	0.26	0.74	0.228 (± 0.17)	1.715 (± 0.10)	0	0.004653
	Yibin	NI	47	13	0.74	0.26	0.213 (± 0.22)	1.550 (± 0.11)	0	0.05625**
	Baoji	<i>B. shensiana</i>	93	54	0.42	0.58	0.247 (± 0.19)	1.711 (± 0.09)	0	0.01674**
		<i>B. circumserrata</i>								
		<i>B. potaninii</i>								
Cluster 2	Hanzhong	<i>B. dasystachya</i>	55	40	0.28	0.72	0.229 (± 0.17)	1.701 (± 0.09)	0	0.001043
	Ankang	<i>B. soulieana</i>	21	14	0.35	0.65	0.242 (± 0.21)	1.650 (± 0.11)	0	0.1202**
	Linzhi		4	4	0.00	1.00		NA ^f	NA	NA
	Cluster 1 total		700	492	0.30	0.70	0.307 (± 0.17)	1.749 (± 0.07)	0.153 (± 0.04)	0.01624
	Tianshui		1	1	NA	NA	NA	NA	NA	NA
	Longnan		1	1	NA	NA	NA	NA	NA	NA
	Aba		1	1	NA	NA	NA	NA	NA	NA
	Baoji		1	1	NA	NA	NA	NA	NA	NA
	Hanzhong		1	1	NA	NA	NA	NA	NA	NA
	Ankang		1	1	NA	NA	NA	NA	NA	NA
Cluster 2	Linzhi	<i>B. polyantha</i>	255	187	0.27	0.73	0.332 (± 0.14)	1.901 (± 0.03)	0.056 (± 0.04)	0.02083
	Cluster 2 total		261	193	0.26	0.74	0.336 (± 0.14)	1.830 (± 0.04)	0.235 (± 0.07)	0.02362

^a Clonal fraction was calculated as $1 - [(\text{number of MLGs}) / (\text{total number of isolates})]$ (Zhan et al. 2002).

^b Gene diversity within populations was calculated from clone-corrected data at 20 microsatellite loci (Yeh et al. 1999). Value inside the parenthesis is standard deviation.

^c Allelic richness and private allelic richness were calculated using ADZE that uses rarefaction. Value inside the parentheses is standard deviations.

^d Multilocus linkage disequilibrium (*LD*) was assessed using the standardized index of association (*rBarD*) and estimated for each population as described previously (Agapow and Burt 2001). The significance of *rBarD* was tested with 1,000 randomizations of the data by comparing the observed value to that expected under the null hypothesis of *rBarD* = 0. The null hypothesis of *LD* was rejected if $P < 0.01$. ** Significant at $P = 0.01$.

^e NI, not investigated.

^f NA, not analyzed.

Pst races in Tibet was different from that in Yunan, Qinghai, and Shaanxi (Hu et al. 2012).

Several studies have showed dispersal of *Pst* in different geographic regions or temporal dynamics using DNA markers. The clonal structure in the Northwest European *Pst* population was confirmed using AFLP (Hovmöller et al. 2002). Similar results were obtained in Australia based on RAPD and AFLP markers (Steele et al. 2001). In China, Gansu and Sichuan provinces have been considered as the major sources of inoculum and new virulence (Wu and Niu 2000) as *Pst* can over-summer in these regions (including Qinghai Province). The genetic diversity in the Longnan region was much higher than in the Linxia and Qinghai regions (Lu et al. 2011). *Pst* population in Longnan is genetically diverse and undergoes extensive genetic exchange within the region (Lu et al. 2009; Zheng et al. 2005), which is supported by the present results. *Pst* genetic diversity in Southeast and Southwest Sichuan was lower than in the Sichuan basin and Gansu Province, which could be related to local climate, environment, geography, and wheat varieties. Chen (2008) studied 20 natural populations of *Pst* in the main epidemiological region and found that the N_m values of Shaanxi and Gansu ranged from 1.1 to 9.0, and the highest N_m value (8.9692) was observed between Piangliang (Gansu) and Baoji (Shaanxi). In contrast, the N_m values between Gansu and Sichuan varied from 1.1 to 2.5 and N_m values between Shaanxi and Sichuan varied from 1.1 to 2.7. These results are supported by the present study although the N_m values in this study were higher than previous study (Chen 2008).

Fig. 5. Relationship of population differentiation (G''_{ST}) with geographical distance among *PST* subpopulations.

Genetic differentiation of *Pst* mainly occurred within populations, agreeing with previous studies (Chen 2008; Lu et al. 2009; Wang et al. 2010a,b). The general consensus is that mutation and somatic recombination could generate increased levels of *Pst* variability (Park and Wellings 2012). Recently the *Pst* sexual stage was observed on barberry (Jin et al. 2010) and a very low level of *Pst* sexual reproduction was reported to have occurred in Shaanxi, Gansu, and Tibet (Wang et al. 2016). Ali et al. (2014b) showed heterogeneity in levels of *Pst* recombination in the Himalayan and near-Himalayan regions; the 71 Chinese *Pst* isolates used in that study originated from Longnan in Gansu Province, China (Ali, personal communication). We also demonstrated heterogeneity in Longnan, in the Himalayan Mountains, and in the southern Qinling Mountains, which might be due to admixtures as alternative hosts of *Pst*: *Berberis* spp. may serve as alternate hosts for *Pst* sexual reproduction. However, barberry is believed not to play a role for stripe rust in the U.S. Pacific Northwest (Wang and Chen 2015).

Tibet is situated in the Qinghai-Tibet Plateau with average altitude exceeding 4,500 m, covering most of Tibet and Qinghai Province. It stretches approximately 1,000 km north to south and 2,500 km east to west and is surrounded by large mountain ranges. The plateau is bordered to the north by the Kunlun Mountains with average altitude >5,500 m, and to the east by the Hengduan Mountains with average altitude >4,100 m. These mountains form a natural barrier in the southwest direction between Tibet and the main *Pst* epidemic regions in China, preventing/minimizing *Pst* gene flows, as indicated by the low N_m estimates in the present study. *Pst* race composition and frequency in Tibet also suggested Tibet as another center of inoculum source and virulence (Hu et al. 2012), agreeing with results based on AFLP markers (Kuang 2010). The Tibet *Pst* population could have been experiencing considerable gene flow with *Pst* populations in Nepal and India via Yarlung Zangbo Canyon. Further studies are necessary to assess this possible dispersal route.

In recent years, the “metapopulation” theory advocated by Burdon and Thrall (2009) has been a predominant theory in population genetics. This theory views the “host-pathogen” system as one entire population in which host and pathogen are the two subgroups. Host varieties exert a strong directional selection pressure on rust pathogens. This may explain why the *Pst* population from Tibet is more or less independent of the other populations because wheat cultivars used in Tibet are different (more correctly different resistance genes) from those used in the other regions. Fewer resistance genes are present in those wheat cultivars grown in Tibet region than in the other regions (Cao et al. 2011; Han et al. 2012; Li et al. 2008, 2015; Peng et al. 2015; Zhang et al. 2012).

In summary, the *Pst* epidemic in the Tibet region can be considered to be independent from the epidemics occurring in the other regions, with the exception of the Xinjiang region in China. In addition, there is extensive *Pst* gene flow within the main wheat production

Table 5. Estimates of population differentiation (G''_{ST}) (above diagonal) and its significance (i.e., P values: below diagonal), and geographical distance (i.e., value in kilometer in parenthesis) for all pairs of subpopulations within the two genetic clusters of *Puccinia striiformis* f. sp. *tritici* as determined from DAPC analysis (Jombart et al. 2010)

Subpop	P1 ^a	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	P13
P1	-	0.062 (233) ^f	0.143 (250)	0.156 (866)	0.146 (1,132)	0.087 (450)	0.087^b (571)	0.103 (683)	0.144 (934)	0.111 (176)	0.129 (406)	0.144 (555)	0.347 (2,365)
P2	0.001	-	0.073 (597)	0.182 (1,111)	0.142 (1,460)	0.084 (748)	0.034 (900)	0.076 (1,011)	0.07 (1,275)	0.02 (207)	0.116 (577)	0.093 (545)	0.339 (2,917)
P3	0.001	0.001	-	0.188 (824)	0.143 (885)	0.116 (202)	0.166 (324)	0.098 (435)	0.092 (401)	0.145 (364)	0.077 (572)	0.356 (2,117)	
P4	0.001	0.001	0.001	-	0.173 (611)	0.109 (678)	0.112 (510)	0.21 (403)	0.202 (664)	0.198 (1,042)	0.245 (839)	0.235 (1,047)	0.311 (1,605)
P5	0.001	0.001	0.001	0.001	-	0.163 (737)	0.205 (570)	0.182 (442)	0.102 (562)	0.178 (1,101)	0.282 (899)	0.269 (1,107)	0.421 (1,687)
P6	0.001	0.001	0.001	0.001	0.001	-	0.078 (175)	0.011 (286)	0.117 (531)	0.105 (387)	0.181 (184)	0.174 (392)	0.332 (1,968)
P7	0.008	0.097	0.001	0.001	0.001	0.015	-	0.129 (120)	0.082 (381)	0.06 (539)	0.155 (337)	0.156 (545)	0.314 (1,802)
P8	0.001	0.002	0.001	0.001	0.001	0.271	0.002	-	0.096 (269)	0.08 (651)	0.189 (448)	0.173 (656)	0.387 (1,675)
P9	0.001	0.014	0.001	0.001	0.004	0.001	0.036	0.01	-	0.07 (900)	0.178 (698)	0.114 (763)	0.373 (1,796)
P10	0.001	0.036	0.001	0.001	0.001	0.001	0.021	0.001	0.011	-	0.105 (213)	0.049 (389)	0.352 (2,333)
P11	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	-	0.014 (223)	0.363 (2,130)
P12	0.001	0.006	0.002	0.001	0.001	0.001	0.008	0.001	0.01	0.02	0.23	-	0.369 (2,341)
P13	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	-

^a P1 (Tianshui), P2 (Pingliang), P3 (Longnan), P4 (Aba), P5 (Liangshan), P6 (Guangyuan), P7 (Mianyang), P8 (Chengdu), P9 (Yibin), P10 (Baoji), P11 (Hanzhong), P12 (Ankang) subpopulations from Cluster 1; P13 (Linzi) subpopulation from Cluster 2.

^b Values in bold indicate no significant difference at $P = 0.001$.

^c Geographical distance in kilometer in the parentheses was estimated using Google Earth.

area in China. Further research is necessary to study the distribution of *Pst* alternative host *Berberis* spp. and their roles in *Pst* epidemiology and genetic diversity.

Acknowledgments

This work was supported by the National Key Basic Research Program of China (2013CB127704), the National Natural Science Foundation of China (31471731 and 31371884), the Program for Innovative Research in Northwest A&F University (X2015021), and partially supported by the 111 Project from Education Ministry of China (B07049). We pay a special thanks to Dr. Pierre Gladieux (Ecologie Systematique Evolution, Univ. Paris-Sud., France) and Dr. Sajid Ali (Institute of Biotechnology and Genetic Engineering, The University of Agriculture, Pakistan) for their valuable suggestions.

Literature Cited

- Agapow, P. M., and Burt, A. 2001. Indices of multilocus linkage disequilibrium. *Mol. Ecol. Notes* 1:101-102.
- Ali, S., Gladieux, P., Leconte, M., Gautier, A., Justesen, A. F., Hovmöller, M. S., Enjalbert, J., and de Vallavieille-Pope, C. 2014a. Origin, migration routes and worldwide population genetic structure of the wheat yellow rust pathogen *Puccinia striiformis* f. sp. *tritici*. *PLoS Pathog* 10:e1003903.
- Ali, S., Gladieux, P., Rahman, H., Saqib, M. S., Fiaz, M., Ahmad, H., Leconte, M., Gautier, A., Justesen, A. F., Hovmöller, M. S., Enjalbert, J., and de Vallavieille-Pope, C. 2014b. Inferring the contribution of sexual reproduction, migration and off-season survival to the temporal maintenance of microbial populations: a case study on the wheat fungal pathogen. *Mol. Ecol.* 23:603-617.
- Burdon, J. J., and Thrall, P. H. 2009. Co-evolution of plants and their pathogens in natural habitats. *Science* 324:755-756.
- Cao, S. Q., Zhang, B., Li, M. J., Xu, S. C., Luo, H. S., Jin, S. L., Jia, Q. Z., Huang, J., and Shang, X. W. 2011. Postulation of stripe rust resistance genes and analysis of adult resistance in 50 wheat varieties (lines) in Gansu Province. *Acta Agron. Sin.* 37:1360-1371.
- Chen, C. Q. 2008. Molecular population genetic structure of *Puccinia striiformis* f. sp. *tritici* in China. Ph.D. Dissertation, Northwest A&F University, Yangling, China (in Chinese with English abstract).
- Chen, C. Q., Zheng, W. M., Buchenauer, H., Huang, L. L., Lu, N. H., and Kang, Z. S. 2009. Isolation of microsatellite loci from expressed sequence tag library of *Puccinia striiformis* f. sp. *tritici*. *Mol. Ecol. Notes* 9:236-238.
- Chen, X. M. 2005. Epidemiology and control of stripe rust [*Puccinia striiformis* f. sp. *tritici*] on wheat. *Can. J. Plant Pathol.* 27:314-337.
- Chen, X. M., Line, R. F., and Leung, H. 1993. Relationship between virulence variation and DNA polymorphism in *Puccinia striiformis*. *Phytopathology* 83:1489-1497.
- Cheng, P., and Chen, X. M. 2014. Virulence and molecular analyses support asexual reproduction of *Puccinia striiformis* f. sp. *tritici* in the U.S. Pacific Northwest. *Phytopathology* 104:1208-1220.
- Culley, T. M., Wallace, L. E., Genger-Nowak, K. M., and Crawford, D. J. 2002. Comparison of two methods of calculating GST, a genetic measure of population differentiation. *Am. J. Bot.* 89:460-465.
- Dilmaghani, A., Gladieux, P., Gout, L., Giraud, T., Brunner, P. C., Stachowiak, A., Balesdent, M. H., and Rouxel, T. 2012. Migration patterns and changes in population biology associated with the worldwide spread of the oilseed rape pathogen *Leptosphaeria maculans*. *Mol. Ecol.* 21:2519-2533.
- Duan, X., Tellier, A., Wan, A., Leconte, M., de Vallavieille-Pope, C., and Enjalbert, J. 2010. *Puccinia striiformis* f. sp. *tritici* presents high diversity and recombination in the over-summering zone of Gansu, China. *Mycologia* 102:44-53.
- Enjalbert, J., Duan, X., Giraud, T., Vautrin, D., de Vallavieille-Pope, C., and Solignac, M. 2002. Isolation of twelve microsatellite loci, using an enrichment protocol in the phytopathogenic fungus *P. striiformis* f. sp. *tritici*. *Mol. Ecol. Notes* 2:563-565.
- Enjalbert, J., Duan, X., Leconte, M., Hovmöller, M. S., and de Vallavieille-Pope, C. 2005. Genetic evidence of local adaptation of wheat yellow rust (*Puccinia striiformis* f. sp. *tritici*) within France. *Mol. Ecol.* 14:2065-2073.
- Evanno, G., Regnaut, S., and Goudet, J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol. Ecol.* 14:2611-2620.
- Falush, D., Stephens, M., and Pritchard, J. K. 2003. Inference of population structure using multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics* 164:1567-1587.
- Han, D. J., Zhang, P. Y., Wang, Q. L., Zeng, Q. D., Wu, J. H., Zhou, X. L., Wang, X. J., Huang, L. L., and Kang, Z. S. 2012. Identification and evaluation of resistance to stripe rust in 1980 wheat landraces and abroad germplasm. *Sci. Agric. Sin.* 45:5013-5023.
- Hedrick, P. W. 2005. A standardized genetic differentiation measure. *Evolution* 59:1633-1638.
- Hovmöller, M. S., Justesen, A. F., and Brown, J. K. M. 2002. Clonality and long-distance migration of *Puccinia striiformis* f. sp. *tritici* in north-west Europe. *Eur. J. Plant Pathol.* 51:24-32.
- Hovmöller, M. S., Sørensen, C. K., Walter, S., and Justesen, A. F. 2011. Diversity of *Puccinia striiformis* f. sp. *tritici* on cereals and grasses. *Annu. Rev. Phytopathol.* 49:197-217.
- Hovmöller, M. S., Walter, S., and Justesen, A. F. 2010. Escalating threat of wheat rusts. *Science* 329:369.
- Hu, X. P., Li, J. J., Wang, Y., Wang, B. T., Li, Q., Kang, Z. S., Yang, M. N., Peng, Y. L., Liu, T. G., Chen, W. Q., and Xu, X. M. 2012. Race composition of *Puccinia striiformis* f. sp. *tritici* in Tibet, China. *Plant Dis.* 96:1615-1620.
- Hu, X. P., Yang, Z. W., Li, Z. Q., Deng, Z. Y., Chen, H. W., and Ke, C. H. 2000. Factors analysis affected the epidemics of wheat stripe rust in Hanzhong district Shaanxi province. *Acta Agr. Boreali-occidentalis Sin.* 9:36-38 (in Chinese with English abstract).
- Jakobsson, M., and Rosenberg, N. A. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* 23:1801-1806.
- Jin, Y., Szabo, L. J., and Carson, M. 2010. Century-old mystery of *Puccinia striiformis* life history solved with the identification of *Berberis* as an alternate host. *Phytopathology* 100:432-435.
- Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* 24:1403-1405.
- Jombart, T., Devillard, S., and Balloux, F. 2010. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genet.* 11:94.
- Kuang, W. J. 2010. Population diversity of *Puccinia striiformis* in Tibet. Master Dissertation, Sichuan Agricultural University, Yaan, China (in Chinese with English abstract).
- Li, F. Q., Han, D. J., Wie, G. R., Zeng, Q. D., Huang, L. L., and Kang, Z. S. 2008. Molecular detection of stripe rust resistant genes in 126 winter wheat varieties from the Huanghuai wheat region. *Sci. Agric. Sin.* 41:3060-3069.
- Li, M. Z., Li, Q., Cao, K. X., Shen, X. X., Fan, Y., Wang, Y., and Wang, B. T. 2015. Molecular detection of stripe rust resistance genes in 115 wheat varieties (lines) from Shaanxi Province. *Acta Phytopathol. Sin.* 45:632-640.
- Li, Z. Q., and Liu, H. W. 1957. Preliminary studies on the trend of occurrence and development of the stripe rust of wheat (*Puccinia glumarum* (Sch.) Erikss. and Henn.) in provinces Shensu, Kansu, and Chinghai. *J. Northwest Agr. Coll.* 1:34-46 (in Chinese with English abstract).
- Li, Z. Q., and Zeng, S. M. 2002. Pages 1-11 in: Wheat rust in China. China Agricultural Press, Beijing (in Chinese).
- Lu, N. H., Wang, J. F., Chen, X. M., Zhan, G. M., Chen, C. Q., Huang, L. L., and Kang, Z. S. 2011. Spatial genetic diversity and interregional spread of *P. striiformis* f. sp. *tritici* in Northwest China. *Eur. J. Plant Pathol.* 131:685-693.
- Lu, N. H., Wang, J. F., Zhan, G. M., Huang, L. L., and Kang, Z. S. 2009. SSR analysis of population genetic diversity of *Puccinia striiformis* f. sp. *tritici* in different altitude regions of Longnan, Gansu. *Mycosystema* 28:496-503.
- Mboup, M., Leconte, M., Gautier, A., Wan, A. M., Chen, W., de Vallavieille-Pope, C., and Enjalbert, J. 2009. Evidence of genetic recombination in wheat yellow rust populations of a Chinese overwintering area. *Fungal Genet. Biol.* 46:299-307.
- Meirmans, P. G., and Hedrick, P. W. 2011. Assessing population structure: F_{ST} and related measures. *Mol. Ecol. Resour.* 11:5-18.
- Nei, M. 1973. Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. USA* 70:3321-3323.
- Nei, M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89:583-590.
- Park, R. F., and Wellings, C. R. 2012. Somatic hybridization in the uredinales. *Annu. Rev. Phytopathol.* 50:219-239.
- Peng, Y. L., Yang, M. N., Dan, B., and Ci, Z. G. 2015. Molecular detection of stripe rust resistance genes in 41 wheat varieties from Tibet. *Acta Phytopathol. Sin.* 45:211-215.
- Pritchard, J. K., Stephens, M., and Donnelly, P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945-959.
- Shan, W. X., Chen, S. Y., Kang, Z. S., Wu, L. R., and Li, Z. Q. 1998. Genetic diversity in *Puccinia striiformis* Westend. f. sp. *tritici* revealed by pathogen genome-specific repetitive sequence. *Can. J. Bot.* 76:587-595.
- Shen, L., Luo, L. M., Chen, W. Q., Zhao, Z. M., and Wang, J. J. 2008. Epidemic zones and dispersal routes of wheat stripe rust caused by *Puccinia striiformis* West. in Sichuan. *Acta Phytophyllac. Sin.* 35:220-226 (in Chinese with English abstract).
- Steele, K. A., Humphreys, E., Wellings, C. R., and Dickinson, M. J. 2001. Support for a stepwise mutation model for pathogen evolution in Australasian *Puccinia striiformis* f. sp. *tritici* by use of molecular markers. *Plant Pathol.* 50:174-180.
- Szpiech, Z. A., Jakobsson, M., and Rosenberg, N. A. 2008. ADZE: a rarefaction approach for counting alleles private to combinations of populations. *Bioinformatics* 24:2498-2504.
- Tibayrenc, M., and Ayala, F. J. 2002. The clonal theory of parasitic protozoa: 12 years on. *Trends Parasitol.* 18:405-410.
- Tibayrenc, M., and Ayala, F. J. 2012. Reproductive clonality of pathogens: A perspective on pathogenic viruses, bacteria, fungi, and parasitic protozoa. *Proc. Natl. Acad. Sci. USA* 109:E3305-E3313.
- Wan, A. M., Zhao, Z. H., Chen, X. M., He, Z. H., Jin, S. L., Jia, Q. Z., Yao, G., Yang, J. X., Wang, B. T., Li, G. B., Bi, Y. Q., and Yuan, S. Y. 2004. Wheat stripe rust epidemic and virulence of *Puccinia striiformis* f. sp. *tritici* in China in 2002. *Plant Dis.* 88:896-904.
- Wang, H. G., Yang, X. B., and Ma, Z. H. 2010a. Long-distance spore transport of wheat stripe rust pathogen from Sichuan, Yunnan, and Guizhou in Southwestern China. *Plant Dis.* 94:873-880.

- Wang, J. F., Cheng, C. Q., Lu, N. H., Peng, Y. L., Zhan, G. M., Huang, L. L., and Kang, Z. S. 2010b. SSR analysis of population genetic diversity of *Puccinia striiformis* f. sp. *tritici* in Sichuan Province, China. *Mycosystema* 29:206-213 (In Chinese with English abstract).
- Wang, M. N., and Chen, X. M. 2015. Barberry does not function as an alternate host for *Puccinia striiformis* f. sp. *tritici* in the US Pacific Northwest due to teliospore degradation and barberry phenology. *Plant Dis.* 99:1500-1506.
- Wang, Z. Y., Zhao, J., Chen, X. M., Peng, Y. L., Ji, J. J., Zhao, S. L., Lv, Y. J., Huang, L. L., and Kang, Z. S. 2016. Virulence variations of *Puccinia striiformis* f. sp. *tritici* isolates collected from *Berberis* spp. in China. *Plant Dis.* 100:131-138.
- Wellings, C. R. 2007. *Puccinia striiformis* in Australian: A review of the incursion, evolution and adaptation of stripe rust in the period 1979-2006. *Aust. J. Agric. Res.* 58:567-575.
- Wright, S. 1951. The genetical structure of population. *Ann. Eugen.* 15:323-354.
- Wu, L. R., and Niu, Y. C. 2000. Strategies of sustainable control of wheat stripe rust in China. *Sci. Agric. Sin.* 33:1-7 (in Chinese with English abstract).
- Xie, S. X., Chen, Y. L., Chen, W. Q., Wang, K. N., Ding, X. P., He, X. L., Pu, J. C., Qi, S. W., Ou, Y., and Du, D. M. 1988. Studies on the epidemiology of wheat stripe rust (*Puccinia striiformis* West.) in Abazhou, Sichuan province. *Acta Phytophylac. Sin.* 15:85-91 (in Chinese with English abstract).
- Xie, S. X., Chen, Y. L., Li, H. Y., Lin, X. L., Sun, Y. H., Wang, D. S., and Zeng, D. X. 1986. Studies on the epidemiology of stripe rust of wheat in Mian-Yang area, Sichuan province. *Acta Phytophylac. Sin.* 13:31-38 (in Chinese with English abstract).
- Yeh, F. C., Yang, R., and Boyle, T. 1999. POPGENE version 1.32: Microsoft windows-based freeware for population genetic analysis, a quick user guide. University of Alberta, Center for International Forestry Research, Alberta, Canada.
- Zeng, S. M., and Luo, Y. 2006. Long-distance spread and interregional epidemics of wheat stripe rust in China. *Plant Dis.* 90:980-988.
- Zeng, S. M., and Luo, Y. 2008. Systems analysis of wheat stripe rust epidemics in China. *Eur. J. Plant Pathol.* 121:425-438.
- Zhan, G., Wang, F., Wan, C., Han, Q., Huang, L., and Kang, Z. 2016. Virulence and molecular diversity of the *Puccinia striiformis* f. sp. *tritici* population in Xinjiang in relation to other regions of western China. *Plant Dis.* 100:99-107.
- Zhan, J., Kema, G. H. J., Waalwijk, C., and McDonald, B. A. 2002. Distribution of mating type alleles in the wheat pathogen *Mycosphaerella graminicola* over spatial scales from lesions to continent. *Fung. Genet. Biol.* 36:128-136.
- Zhang, Z. Y., Ji, H. L., Shen, L., Xu, S. C., Ni, J. Y., and Peng, Y. L. 2012. Postulation of resistance genes and evaluation of adult resistance to stripe rust in 58 cultivars from Sichuan. *Acta Phytophylac. Sin.* 39:13-23.
- Zhao, J., Wang, L., Wang, Z. Y., Chen, X. M., Zhang, H. C., Yao, J. N., Zhan, G. M., Chen, W., Huang, L. L., and Kang, Z. S. 2013. Identification of eighteen *Berberis* species as alternate hosts of *Puccinia striiformis* f. sp. *tritici* and virulence variation in the pathogen isolates from natural infection of barberry plants in China. *Phytopathology* 103:927-934.
- Zheng, W. M., Chen, S. Y., Kang, Z. S., Wang, Y., Wu, L. R., and Li, Z. Q. 2005. DNA fingerprinting of natural population of *Puccinia striiformis* f. sp. *tritici* in Tianshui area. *Mycosystema* 24:199-206 (in Chinese with English abstract).