Distribution and Characterization of *Monilinia* spp. Causing Apple Fruit Decay in Serbia

Miljan Vasić and Ivana Vico, University of Belgrade, Faculty of Agriculture, Belgrade 11080, Serbia; Wayne M. Jurick II, USDA-ARS, Food Quality Laboratory, Beltsville, MD; and Nataša Duduk,[†] University of Belgrade, Faculty of Agriculture, Belgrade 11080, Serbia

Abstract

Brown rot, caused by *Monilinia* spp., is an economically important pre- and postharvest disease of pome and stone fruits worldwide. In Serbia, apple is the most widely grown pome fruit, and the distribution of economically important *Monilinia* spp. responsible for apple brown rot is unknown. Hence, we conducted a three year survey, from 2010 to 2012, where 349 isolates were obtained from six orchards and four storage facilities from five different apple cultivars with brown rot symptoms. Morphological characterization of the isolates, multiplex PCR, and phylogenetic

analysis revealed four species: *M. fructigena*, *M. laxa*, *M. fructicola*, and *Monilia polystroma*. All species were found in the orchard and in storage, with *M. fructigena* predominating, followed by *M. polystroma*. Representative isolates were analyzed in vitro and in vivo where differences in growth rate, sporulation, and virulence on apple fruit were observed. Findings from this investigation demonstrate diversity in the species responsible for pre- and postharvest apple brown rot, which has significant implications for pathogen detection and for developing disease-specific management strategies.

Brown rot is caused by *Monilinia* spp. and is one of the most destructive pre- and postharvest diseases of pome and stone fruits (Batra 1991; Byrde and Willetts 1977; Holb and Scherm 2007; van Leeuwen et al. 2000). Economic losses ranging from 7 to 25% and from 0.6 to 8% were found in orchards and storage facilities in Europe, respectively (Berrie and Holb 2014). Species distribution and prevalence differ depending on the geographic region, origin, and host. For instance, *M. fructigena* (Aderhold and Ruhland) is the most important causal agent of brown rot of pome fruits in Europe, while it is a quarantine pathogen for North America and Australia (Batra 1991; Byrde and Willetts 1977). *M. fructicola* (G. Winter) Honey and *M. laxa* (Aderhold and Ruhland) are prevalent species on stone fruits occasionally affecting pome fruits (Batra 1991; Byrde and Willetts 1977; Peter et al. 2015). Until 2001, *M. fructicola* was endemic in North and South America and Oceania (Batra 1991; Byrde and Willetts 1977; OEPP/EPPO 2002), when it was introduced in France. In the last decade, this pathogen was detected in most European countries including Serbia (EFSA Panel 2011; Hrustić et al. 2013; Vasić et al. 2012) and China (Zhu et al. 2011, 2016). *M. laxa* is globally distributed (Batra 1991; Byrde and Willetts 1977). A fourth species, *Monilia polystroma* van Leeuwen is newly described, originating from Japan, and is closely related to *M. fructigena* (van Leeuwen et al. 2002). *M. polystroma* was detected on different pome and stone fruits in Hungary (Petróczy and Palkovics 2009), Czech Republic (OEPP/EPPO 2011), Poland (Poniatowska et al. 2016), Serbia (Vasić et al. 2013), Italy (Martini et al. 2015), Croatia (Di Francesco et al. 2015), and Slovenia (Munda 2015). In Asia, two additional species of *Monilinia* are distributed on pome and stone fruits: *M. yunnanensis* (M. J. Hu and C. X. Luo. sp) and *M. mumecola* (Y. Sasaki and T. Sano) (Harada et al. 2004; Hu et al. 2011; Zhu et al. 2016). A recent study by Zhu et al. (2016) reported a high frequency (20%) of *M. polystroma* isolates during a survey for *Monilinia* spp. in pome fruit producing regions in China.

In Serbia, apple is one of the most widely grown fruit crops. In 2014, harvested area was 23,737 ha with production about 336,313 tons (<http://www.fao.org/faostat/en/#data>). Yield is threatened by numerous pathogens including *Monilinia* spp. Ivanović and Ivanović (2001) reported losses up to 80% in Serbian organic apple orchards caused by *M. fructigena*. Distribution of newly introduced pathogens could

change species composition and may increase pre- and postharvest fruit decay. In Serbia, four species are known as causal agents of apple brown rot: *M. fructigena*, *M. laxa*, *M. polystroma*, and *M. fructicola* (Vasić et al. 2012, 2013, 2016), but their frequency and geographic distribution is currently unknown. In another study in Serbia, Hrustić et al. (2015) observed an increased number of *M. fructicola* isolates causing brown rot of stone fruits in 2012 and 2013 following its initial detection in 2011. Similarly, the composition of *Monilinia* spp. in apple fruit production may also differ. In complex pathosystems, variability in biological features of closely related species could help determine the pathogen's ability to adapt and thrive (Byrde and Willetts 1977). For example, *M. polystroma* is capable of forming thicker, more compact and complex stromata on apple fruit than *M. fructigena*. This factor alone may enhance the survival of the fungus and increase the amount of primary inoculum available for the next season (Vasić et al. 2016).

Monilinia spp. also have different lifestyles as *M. fructicola* and *M. laxa* cause both blossom and twig blight and pre- and postharvest fruit rot, while *M. fructigena* and *M. polystroma* are primarily fruit pathogens (Byrde and Willetts 1977; Holb and Scherm 2007; van Leeuwen et al. 2000; Vasić et al. 2016). Fungal growth, as influenced by different ecological factors and available nutrients, contribute to the basic knowledge of fungal pathogens that might provide key information for developing disease-specific management strategies. Sugar content and pH value of host tissue influences the pathogen's ability to colonize its host, while temperature plays an important role in disease development and epidemiology. Ecological and cultural characteristics concerning newly described and emerging *Monilinia* spp. and its comparison with other well characterized species has not been conducted in detail. Holb (2004) showed that *M. fructigena* mycelial growth was faster under lower pH compared with *M. polystroma*, at pH values between 2.5 and 6.5. Holb and Chauhan (2004) observed that the optimal carbohydrate sources for mycelial growth rate of *M. polystroma* and *M. fructigena* were glucose, fructose, and saccharose, while the best nitrogen source was peptone. However, the influence of different temperatures, varying pH range, and commercially used media were not determined for *M. polystroma*. The main objectives for the current study were to: 1) identify *Monilinia* spp. diversity, both pre- and postharvest, from apple fruit of different production areas in Serbia; 2) compare cultural and morphological factors influencing *Monilinia* spp. growth and reproduction; and 3) assess pathogenic potential and virulence of *Monilinia* spp. on apple fruit.

Material and Methods

Sample collection, fungal isolation, and culture storage. In a 3-year survey from 2010 to 2012, apple fruit 'Idared,' 'Golden Delicious,' 'Granny Smith,' 'Elstar,' and 'Jonagold' with brown rot

[†]Corresponding author: Nataša Duduk; E-mail: natasadukic@yahoo.com

symptoms were collected from 15 apple orchards located in Svilajnac, Topola, Bela Crkva, Ruma, Grocka, and Smederevo and six storage facilities located in Svilajnac, Bela Crkva, Grocka and Smederevo (Fig. 1). Isolation was done from the lesion margin after surface sanitization using 70% ethanol and aseptic removal of the skin. Fragments of tissue were placed onto potato dextrose agar (PDA, EMD Chemicals Inc, Gibbstown, NJ, U.S.A.) amended with 2.5 ml of 25% lactic acid per liter of medium and incubated at 22°C in the dark for 4 days in an incubator. Subsequently, mycelial fragments taken from the growing colony margin were transferred to fresh PDA. Single spore cultures were obtained from conidia produced on V8 medium (200 ml V8 juice and 20 g agar per liter). One isolate, due to the inability to sporulate on either media or inoculated fruit, was propagated by hyphal tip transfer. The isolates were maintained in 15% glycerol at -80°C as mycelial plugs taken from 7-day-old cultures grown on PDA. Isolates of *M. fructigena* and *M. polystroma* (Vasić et al. 2016) were also included in this study for comparison (Table 1).

Pathogenicity test and reisolation of *Monilinia* spp. from apple fruit. Pathogenicity of the isolates was tested on mature Idared fruit. Apples were surface-sanitized by immersing in 1% solution of sodium hypochlorite (NaOCl) for 2 min and rinsed twice in sterile distilled water before inoculation. A 5-mm-diameter mycelial plug was taken

from the margin of 5-day-old colonies grown on PDA and inserted into a hole (6 mm diameter and 5 mm depth) in the cortex of each fruit using a cork borer. Apple fruit inoculated with uncolonized PDA plugs were used as controls. Inoculated and control fruit were placed in plastic containers at 22 ± 1°C under high humidity (RH 95 to 100%) and alternating 14 h light/10 h dark cycles and incubated for 5 days. Three replicates per isolate were used. Reisolation was performed as described above (see sample collection and fungal isolation).

Morphological identification of *Monilinia* spp. The morphology of each isolate was determined on PDA. Morphological characterization was performed according to van Leeuwen et al. (2002) with slight modifications. For morphological characterization, 5-mm-diameter mycelial plugs were taken from the edge of the 5-day-old colonies and inoculated in the center of a PDA plate. Colony characteristics (color, aerial mycelium, margin shape, presence of reproductive structures, and formation of stroma) of *Monilinia* spp. isolates were determined on PDA after 12 days of incubation in 12 h light/12 h dark cycle at 22 ± 1°C using a fluorescent lamp approximately 15 cm above the Petri dishes as described by Byrde and Willets (1977), Batra (1991), and OEPP/EPPO (2009).

DNA extraction and multiplex PCR. Fungal mycelium was obtained from isolates grown on PDA for 7 days. DNA extraction

Fig. 1. *Monilinia* spp. isolates obtained from apple fruit in Serbia.

was performed following the CTAB protocol of Day and Shattock (1997). Total DNA was precipitated with isopropanol, resuspended in 30 µl of TE buffer (10 mM Tris pH 8 and 1 mM EDTA), and stored at -20°C. DNA was amplified via multiplex PCR with primers based on a genomic region of *Monilinia* spp. with unknown function, forward MO368-5 and three reverse Laxa-R2, MO368-8R, and MO368-10R, which yield products of 402 bp for *M. fructigena*,

425 bp for *M. polystroma*, 535 bp for *M. fructicola* and 351 bp for *M. laxa* (Côté et al. 2004). PCR reaction mix (25 µl) contained 1 µl of template DNA, 1× PCR MasterMix (Fermentas, Vilnius, Lithuania), and 0.2 µM of each primer. PCR cycling parameters were as follows: initial denaturation at 95°C for 2 min, followed by 35 cycles of 95°C for 15 s, 60°C for 15 s, and 72°C for 1 min, with final extension at 72°C for 3 min. The PCR product (5 µl) was separated in

Table 1. *Monilinia* spp. isolates and rDNA-ITS region sequences used in this study

Isolates	Species	Geographic origin	Host	GenBank accession number
Isolates from Serbia				
MPSRB-2G ^{u,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	KP279655
MPSRB-20 ^{u,v,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	KP279656
MPSRB-13 ^{u,v,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	KP279657
MPSRB-13A ^{u,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	JX315717
MPSRB-3 ^{u,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	KP279658
MPSRB-2 ^{u,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	KP279659
MPSRB-36 ^{u,v,w,x}	<i>M. polystroma</i>	Bela Crkva	Apple	KP279660
MPSRB-6 ^w	<i>M. polystroma</i>	Bela Crkva	Apple	nd ^z
MPSRB-21 ^w	<i>M. polystroma</i>	Bela Crkva	Apple	nd
MFSRB-1 ^{u,w,x}	<i>M. fructigena</i>	Svilajnac	Apple	KP279661
MFSRB-10 ^{u,w,x}	<i>M. fructigena</i>	Bela Crkva	Apple	KP279662
MFSRB-17 ^{u,v,w,x}	<i>M. fructigena</i>	Grocka	Apple	KP279663
MFSRB-2 ^{u,w,x}	<i>M. fructigena</i>	Grocka	Apple	KP279664
MFSRB-32 ^{u,w,x}	<i>M. fructigena</i>	Svilajnac	Apple	KP279665
MFSRB-6 ^{u,v,w,x}	<i>M. fructigena</i>	Svilajnac	Apple	KP279666
MFSRB-3 ^{u,w,x}	<i>M. fructigena</i>	Topola	Apple	KP279667
MFSRB-22 ^{u,v,w,x}	<i>M. fructigena</i>	Bela Crkva	Apple	KP279668
MFSRB-6SD ^w	<i>M. fructigena</i>	Smederevo	Apple	nd
MFSRB-8SD ^w	<i>M. fructigena</i>	Smederevo	Apple	nd
MFC-20 ^{u,v,w}	<i>M. fructicola</i>	Grocka	Apple	JN176564
MFC-28 ^{u,w}	<i>M. fructicola</i>	Ruma	Apple	KN695550
MFC-1 ^{v,w}	<i>M. fructicola</i>	Smederevo	Apple	nd
MLX-SV1 ^{u,w}	<i>M. laxa</i>	Svilajnac	Apple	KP279669
MLX-7 ^{u,w}	<i>M. laxa</i>	Svilajnac	Apple	KN695549
MLX-13 ^{v,w}	<i>M. laxa</i>	Smederevo	Apple	nd
MLX-24 ^{v,w}	<i>M. laxa</i>	Ruma	Apple	nd
Sequences from NCBI ^y				
16FG	<i>M. fructigena</i>	Hungary	Apple	AM937110
SPBA	<i>M. fructigena</i>	Serbia	Plum	KC544805
LAI1B	<i>M. fructigena</i>	France	Apple	AF150677
759.K	<i>M. fructigena</i>	Unknown	Apple	Z73779
COY2M	<i>M. fructigena</i>	France	Plum	AF150680
770.S	<i>M. fructigena</i>	Unknown	Apple	Z73780
ES41	<i>M. fructigena</i>	Spain	Unknown	EF207429
W17	<i>M. fructigena</i>	Spain	Unknown	EF207424
UFT	<i>M. polystroma</i>	Hungary	Apple	AM937114
AP1	<i>M. polystroma</i>	Poland	Apple	JF820317
09-G4	<i>M. polystroma</i>	Switzerland	Apricot	JN128835
2319	<i>M. polystroma</i>	Japan	Unknown	HQ856916
HML-3	<i>M. polystroma</i>	China	Plum	GU067539
P169	<i>M. fructicola</i>	Italy	Nectarine	FJ411109
LMK12	<i>M. fructicola</i>	Unknown	Peach	Z73777
M 10020029	<i>M. fructicola</i>	Slovenia	Peach	GU967379
SPT16	<i>M. fructicola</i>	China	Peach	HQ846920
NPUD1	<i>M. fructicola</i>	Serbia	Nectarine	KC544808
M1PL	<i>M. fructicola</i>	Poland	Unknown	JX312665
SFT1	<i>M. fructicola</i>	China	Unknown	HQ846921
1067. K	<i>M. laxa</i>	Unknown	Apple	Z73784
BPZK	<i>M. laxa</i>	Serbia	Peach	KC544793
L2	<i>M. laxa</i>	Spain	Unknown	EF153013
T3	<i>M. laxa</i>	Spain	Unknown	EF153014
LUC1A	<i>M. laxa</i>	France	Plum	AF150673
BF-SZ-1	<i>M. laxa</i>	China	Peach	HQ908790

^u Isolates used in phylogenetic analysis.

^v Isolates used for cultural and ecological tests.

^w Isolates used in virulence and sporulation tests.

^x Identified isolates used in virulence test (Vasić et al. 2016).

^y National Center for Biotechnology Information.

^z nd: Sequence was not determined.

a 1.5% agarose gel, stained with ethidium bromide, and visualized with a UV transilluminator.

PCR-RFLP analysis. PCR-RFLP was conducted according to the method described by Vasić et al. (2016). The ITS1-5.8S-ITS2 region was amplified with the universal primer pair ITS1/ITS4 (White et al. 1990). PCR reaction mix (25 µl) contained 1 µl of template DNA, 1× PCR MasterMix (Fermentas, Vilnius, Lithuania), and 0.4 µM of each primer. Cycling parameters for the ITS region consisted of: initial denaturation at 94°C for 90 s, followed by 30 cycles of 94°C for 30 s, 55°C for 30 s, and 72°C for 30 s, and final extension at 72°C for 10 min. RFLP analysis of ITS1/ITS4-amplified PCR products (~550 bp) of *Monilinia* spp. isolates (Fig. 1) was performed using *HhaI* restriction endonuclease (Fermentas, Vilnius, Lithuania), according to the manufacturer's instructions. RFLP products were separated on a 2% agarose gel, stained using ethidium bromide, and visualized with a UV transilluminator.

Sequence analysis of rDNA-ITS region and phylogeny. Amplified ITS1/ITS4 products (~550 bp) were purified using mi-PCR Purification Kit (Metabion International AG, Germany) and sequenced using ITS1 and ITS4 primers. Sequences were assembled using Pregap4 from the Staden program package (Staden et al. 2000), aligned using Clustal X (Thompson et al. 1997) under MEGA version 5 (Tamura et al. 2011), and 2× consensus was deposited in GenBank. Sequences were compared with others in GenBank using the MegaBlast algorithm. Phylogenetic analysis of 19 sequences from Serbian apple fruit and 24 reference sequences were retrieved from GenBank from published sources (Table 1) was performed for the noncoding region (ITS1-5.8S-ITS2). For constructing the ITS phylogenetic tree, *Botryotinia fuckeliana* sequence was designated as an outgroup. Maximum parsimony (MP) was performed to carry out phylogenetic construction using MEGA version 6.0 (Tamura et al. 2013). Bootstrap values were estimated by 1000 replicates to test clade stability.

Effect of different media on mycelial growth and sporulation of *Monilinia* spp. Mycelial growth of selected isolates was evaluated on five different media. PDA (EMB Chemicals Inc., Gibbstown, NJ, U.S.A.) and Czapeck dox agar (CYA, Honeywell Fluka, Seelze, Germany) were prepared according to the manufacturer's instructions. Malt extract agar (MEA) was prepared from 20 g malt extract (Sigma-Aldrich) and 15 g agar by adding 1 liter of distilled water according to procedure described by Kim et al. (2005). Apple juice agar (AJA) was prepared by adding 200 ml of apple juice in 800 ml of distilled water, and adding 3 g CaCO₃ and 15 g agar according to procedure described by Kim et al. (2005). V8 agar (V8) was prepared by adding 200 ml of V8 juice in 800 ml of distilled water, and adding 3 g CaCO₃ and 15 g agar according to procedure described by Gams et al. (1987). The media were sterilized for 20 min at 121°C using an autoclave. Isolates were maintained on PDA for 5 days. Mycelial plugs, 5 mm in diameter, were removed from the growing edge of the colony and inoculated in the center of a Petri plate containing PDA medium. Subsequently, plates were wrapped with Parafilm and placed in incubators at 25°C. Mean colony diameter, excluding the plug, was measured daily and growth rate was expressed as mm of growth per day. Sporulation intensity was assessed according to method described by Hu et al. (2011) with slight modifications. After 7 days of incubation, plates were flooded with 5 ml of sterile distilled water and the colonies were softly scraped with a glass rod. Conidial suspensions were filtered through a double layer of cheesecloth and conidial concentration was determined using a hemacytometer and compound light microscope. The number of conidia produced per cm² for each colony was calculated as a quotient of the number of conidia produced per colony and colony area. Five replicates per isolate were used and the entire assay was done twice.

Effect of temperature and pH on mycelial growth of *Monilinia* spp. The effect of various incubation temperatures ranging from 5 to 35°C on mycelial growth of select representative isolates was evaluated on PDA. Inoculum and inoculation of different media were done as described above (see effect of different media on mycelial growth and sporulation). Mean colony growth per day was calculated as described above (see effect of different media

on mycelial growth). Five replicates per isolate were used and the experiment was done twice. The effect of different culture pH on mycelial growth of representative isolates was evaluated on PDA. Different pH values ranging from 3 to 9 were adjusted using a pH-meter (InoLab pH 220, Xylem Analytics, Germany) by adding 0.1 N HCL or 0.1 N NaOH into the medium. After autoclaving and cooling, pH was verified with pH test strips (universal indicator, pH 0-14, Merck, Germany). Inoculum and inoculation of different media were done as described above (see effect of different media on mycelial growth). Inoculated plates were incubated at 25°C. Mean colony growth per day was calculated as described above (see effect of different media on mycelial growth). Five replicates per isolate were used and the experiment was done twice.

Virulence test and sporulation intensity on apple fruit. Analysis of pathogen virulence was conducted using a representative isolate of each of the four *Monilinia* spp. Mature apple fruit Golden Delicious, Idared, and Granny Smith were wound-inoculated as described in the pathogenicity test. Lesion size was measured at equatorial and longitudinal axes 5 days postinoculation. Sporulation intensity was assessed according to the method described by van Leeuwen et al. (2002) with slight modifications. Six days after incubation, conidia were collected by shaking fruit in 30 ml of sterile distilled water at 250 rpm for 30 s in an orbital shaker. Conidial suspensions were filtered through a double layer of cheesecloth and conidial concentration was determined using a hemacytometer and compound light microscope. The total number of conidia produced per fruit was calculated by multiplying the number of conidia per ml with the collected volume in ml. The number of conidia produced per cm² of lesion was calculated as a quotient of the number of conidia produced per fruit and lesion area. Sporulation data were log-transformed for statistical analysis. Six replicates per isolate were tested and the experiment was conducted twice.

Statistical analysis. All experiments were conducted at least twice. To determine whether the data from two experiments could be combined, the analysis of variance (ANOVA) was used to test homogeneity of variance. If data from both runs of each experiment were not significantly different, then they were combined. Conidial production data were log transformed before analysis so not to violate the ANOVA assumption. All analyses were performed with STATGRAPHICS (Centurion StatPoint 2005). Treatment means were separated using Tukey post hoc test. Virulence was analyzed with two-factor ANOVA testing the influence of species and cultivars on lesion size. Statistical significance for all tests was defined as $P < 0.05$.

Results

***Monilinia* spp. isolate origin and pathogenicity on apple.** During a three-year study from 2010 to 2012, 291 *Monilinia* spp. isolates were obtained from symptomatic apple fruit collected in six orchards and 58 from four storage facilities, totaling 349 isolates. All inoculated apples showed symptoms of brown rot 5 days postinoculation, while control fruit remained symptomless, indicating their pathogenic capability. Lesion areas were brown with entire to slightly undulate margins. On decayed areas, numerous yellow sporodochia or white mycelia with sparse sporulation were observed. On some fruit, around inoculation sites, grayish conidial masses were evident. In total, 349 reisolates of *Monilinia* spp. were obtained from symptomatic apples, both pre- and postharvest (Fig. 1).

Cultural morphology. Colony morphology of isolates on PDA indicated the presence of four distinct phenotypes (Fig. 2). It was observed that 236 isolates formed whitish colonies with entire to slightly undulate margins, with sparse aerial mycelia, did not sporulate on PDA medium, and lacked stromatization in the colony reverse. These isolates were later confirmed to be *M. fructigena*. Thirty-three isolates formed grayish colonies with entire to slightly undulate margins, while aerial mycelia were sparse, and sporulation was not observed. The reverse exhibited mycelial pigmentation along with minor stromatization and were later considered *M. fructigena*/ *M. polystroma*. Forty-eight isolates formed grayish colonies with entire to slightly undulate margins, while aerial mycelia were sparse and

sporulation was not evident. The colony reverse developed black stromatal plates in concentric rings around inoculation points and were later considered to be *M. polystroma*. Nineteen isolates were whitish to gray with lobed margins and were devoid of sporulation and stromatization. These isolates were later identified as *M. laxa*. Thirteen isolates formed grayish colonies, developing spore masses in concentric rings, colony margins were even, and colony reverse without stromatization and were labeled *M. fructicola*.

Molecular identification and phylogeny. Molecular identification using multiplex PCR confirmed the morphological identification

of four *Monilinia* spp. Two hundred sixty-nine isolates (first and second morphological group) were identified as *M. fructigena* (402-bp amplicon), 48 as *M. polystroma* (421-bp amplicon), 19 as *M. laxa* (325-bp amplicon), and 13 as *M. fructicola* (535-bp amplicon). RFLP analysis using ITS1/ITS4 amplicon digested with *Hha*I restriction enzyme yielded RFLP profiles of *M. polystroma*, which were distinct from other *Monilinia* spp. Phylogenetic analysis based on the ITS sequences revealed divergent clusters for each species, while *M. fructigena* and *M. polystroma* were more closely related (Fig. 3). A phylogenetic tree based on ITS sequences also

Fig. 2. Colony characteristics of *Monilinia* spp. grown on PDA in 12 h light/12 h dark regime after 12 days of incubation. **A, B, C,** *M. fructigena* (MFSRB-1, MFSRB-22, and MFSRB-32); **D,** *M. polystroma* (MPSRB-20); **E,** *M. laxa* (MLX-13); and **F,** *M. fructicola* (MFC-1).

revealed high bootstrap values within each clade, and provided an indication of strong phylogenetic relationships between *Monilinia* spp. causing pre- and postharvest apple decay.

Diversity of *Monilinia* spp. causing apple brown rot in Serbia. In Serbian apple orchards the dominant causal agent of brown rot was *M. fructigena* (76.98%), followed by *M. polystroma* (14.78%), *M. laxa* (5.15%), and *M. fructicola* (3.09%) (Table 2). In results from the 2011 survey for *Monilinia* spp. in Bela Crkva region, *M. polystroma* was detected in more samples (62.07%) than *M. fructigena* (33.93%), while similar trends were observed in 2012 as 69.45% were *M. polystroma* compared with *M. fructigena* at 30.55%. Under storage conditions, *M. fructigena* was the dominant species (77.58%), followed by *M. polystroma* (8.62%), *M. fructicola* (6.9%), and *M. laxa* (6.9%) (Table 2).

Characterization of growth rate on different media, temperatures, and pH levels in vitro. Culture media affected the mycelial growth of *Monilinia* spp. in vitro ($P < 0.05$). *M. fructigena*, *M. polystroma*, *M. laxa*, and *M. fructicola* exhibited the fastest growth on V8 and PDA, but grew slower on AJA, MEA, and CYA (Table 3). The slowest isolate MFSRB-6 (*M. fructigena*) had a similar growth rate on all media. Isolates of *M. fructigena*, *M. polystroma*, and *M. laxa* sporulated only on V8 media (Table 4), whereas isolates of *M. fructicola* sporulated on all media examined. Optimal media for *M. fructicola* sporulation were PDA and V8, followed by MEA, while CYA and AJA were all less favorable. Temperature also affected mycelial growth of *Monilinia* spp. in vitro ($P < 0.05$). All isolates grew at temperatures from 5 to 30°C, while no growth was observed at 35°C (Fig. 4). Slowest growth was observed at 5°C, and optimal temperature

Fig. 3. Phylogenetic tree of rDNA region ITS1-5.8S-ITS2 for 19 *Monilinia* spp. isolates originating from apple fruit and 24 reference isolates from GenBank. *Botryotinia fuckeliana* was used as an outgroup. The tree was constructed using the maximum parsimony method and bootstrap values more than 60 are shown above or below branches. The scale represents number of changes.

occurred at 25°C for all isolates. *M. polystroma* grew the fastest between 15 and 25°C, while isolates of *M. fructigena*, *M. laxa*, and *M. fructicola* grew optimally between 20 and 30°C. Culture pH significantly impacted fungal growth in vitro ($P < 0.05$) (Fig. 5). Optimal pH for *M. fructigena* occurred between 3 and 4, *M. polystroma* 3 to 6, *M. fructicola* between 6 and 8, and *M. laxa* between 5 and 8.

Analysis of virulence and sporulation in different commercial apple cultivars. A significant influence of species and apple cultivar on lesion size ($P < 0.05$) was detected via two-factor ANOVA. Interaction between isolates of four species and cultivars was also significantly different ($P = 0.004$). Four species produced larger lesions on Golden Delicious than on Idared and Granny Smith (Fig. 6A). On Golden Delicious, *M. fructigena* and *M. polystroma* were the most virulent and did not differ significantly ($P = 0.406$). *M. laxa* had lower virulence, while *M. fructicola* was the least virulent. On Idared, *M. fructigena* was the most virulent, *M. fructicola* the least, while *M. polystroma* exhibited differences in virulence where lesion size was not significantly different from *M. fructigena* ($P = 0.219$) and *M. laxa* ($P = 0.378$). On Granny Smith, *M. fructigena* was most virulent, followed by *M. polystroma*, as *M. fructicola* was least virulent. *M. laxa* exhibited variability in lesion size and was not significantly different from *M. polystroma* ($P = 0.938$) and *M. fructicola* ($P = 0.256$).

Table 2. Geographic distribution of *Monilinia* spp. isolates from apple fruit in Serbia

Species	Location	Number of isolates	
		Orchard	Storage
<i>M. fructigena</i>	Svilajnac	69	15
	Smederevo	49	17
	Grocka	20	9
	Bela Crkva	22	4
	Ruma	42	— ^z
	Topola	22	—
	Total (%)	224 (76.98)	45 (77.58)
<i>M. laxa</i>	Svilajnac	2	0
	Smederevo	5	4
	Grocka	4	0
	Ruma	4	—
	Total (%)	15 (5.15)	4 (6.9)
<i>M. fructicola</i>	Smederevo	3	0
	Grocka	3	4
	Ruma	3	—
	Total (%)	9 (3.09)	4 (6.9)
<i>M. polystroma</i>	Bela Crkva	43	5
	Total (%)	43 (14.78)	5 (8.62)

^z '—' indicates that diseased apple fruit were not collected and isolates were not obtained.

A two-factor ANOVA revealed a statistically significant effect of species and apple cultivar on sporulation intensity ($P < 0.05$). Interaction between isolates of four species and cultivars was significantly different ($P = 0.001$). Four species had higher sporulation on Idared and Golden Delicious compared with Granny Smith (Fig. 6B). On Idared, *M. fructigena* and *M. fructicola* sporulated abundantly and did not differ statistically ($P = 0.995$), followed by *M. polystroma*, which sporulated at a lower amount. On Golden Delicious, all species sporulated similarly as on Idared, whereas *M. fructicola* and *M. fructigena* had the highest sporulation intensity ($P = 0.357$), followed by *M. polystroma*, which had slightly lower sporulation. On Granny Smith, *M. fructicola* had the most abundant sporulation, followed by *M. fructigena* and *M. polystroma*, which did not differ significantly ($P = 0.528$). In comparison with other tested species, *M. laxa* had the lowest sporulation levels on three different apple cultivars.

Discussion

During a three-year sampling period from 2010 to 2012, *M. fructigena* was the dominant causal agent of apple brown rot both pre- and postharvest in Serbia, followed by *M. polystroma*, *M. laxa*, and *M. fructicola*. Distribution of *Monilinia* spp. in complex pathosystems, where more than one species is present, has been shown primarily in stone fruit-producing regions. Michailides et al. (1987) observed increased populations of *M. fructicola* compared with *M. laxa* in California, which mirrors what was found in European countries such as Spain, Greece, and Serbia (Hrustić et al. 2015; Papavasileiou et al. 2015; Villarino et al. 2013). However, in Poland, *M. fructigena* was most prevalent on apricot, peach, and plum, while the incidence of *M. laxa*, *M. fructicola*, and *M. polystroma* was low (Poniatowska et al. 2013). Distribution of *Monilinia* spp. in apple and pear fruit production has recently been described in China by Zhu et al. (2016). Interestingly, the brown rot species composition on pomes in China was different from Serbia, where the dominant species was *M. yunnanensis* (77%), followed by *M. polystroma* (20%) and *M. fructicola* (3%). *M. polystroma* was detected only in eastern China, while *M. fructigena* was undetected. In contrast, we have observed that *M. fructigena* is the main agent of apple brown rot in Serbia, which is in agreement with literature for other European countries (Holb and Scherm 2007; van Leeuwen et al. 2000).

M. polystroma is a newly threatening species for apple fruit production in Serbia, since more isolates of *M. polystroma* were detected 2012 compared with 2011 in Bela Crkva region. There is limited data about the potential spread of *M. polystroma*, which is very similar to *M. fructigena*, making it potentially dangerous since *M. fructigena* is a quarantine species for North America and Australia and also causes significant production losses in Europe. Research by Vasić et al. (2013) showed that *M. polystroma* occurred in apple orchards in Serbia, but in the current study, for the first time, we found this species in storage on apple fruit 4 months after harvest. *Monilinia* spp.

Table 3. Mycelial growth rate of *Monilinia* spp. isolates on five media at 25°C in the dark

Species	Isolate	Colony growth rate on culture media (mm/day $\pm$ SE) ^u				
		MEA ^v	AJA ^w	PDA ^x	CYA ^y	V8 ^z
<i>M. fructigena</i>	MFSRB-17	6.93 $\pm$ 0.43 a	6.83 $\pm$ 0.14 a	10.71 $\pm$ 0.25 bc	7.19 $\pm$ 0.37 a	11.92 $\pm$ 0.08 c
	MFSRB-22	4.07 $\pm$ 0.15 ab	3.56 $\pm$ 0.54 a	7.81 $\pm$ 0.92 bc	4.59 $\pm$ 0.42 ab	8.15 $\pm$ 0.26 c
	MFSRB-6	1.93 $\pm$ 0.4 6ab	0.96 $\pm$ 0.58 a	1.29 $\pm$ 0.10 ab	0.74 $\pm$ 0.36 a	1.33 $\pm$ 0.41 ab
<i>M. polystroma</i>	MPSRB-36	4.36 $\pm$ 0.89 ab	3.18 $\pm$ 0.27 a	6.12 $\pm$ 1.09 bc	3.99 $\pm$ 0.64 a	7.88 $\pm$ 0.60 c
	MPSRB-13	4.81 $\pm$ 0.10 ab	3.55 $\pm$ 0.14 a	7.81 $\pm$ 0.49 bc	3.4 $\pm$ 1.14 a	8.14 $\pm$ 0.58 bc
	MPSRB-20	5.02 $\pm$ 0.15 ab	4.71 $\pm$ 0.16 a	7.28 $\pm$ 0.31c	4.45 $\pm$ 1.59 a	7.68 $\pm$ 1.41 cd
<i>M. fructicola</i>	MFC-1	8.17 $\pm$ 0.19 b	5.11 $\pm$ 1.13 a	11.45 $\pm$ 0.06 cd	4.72 $\pm$ 0.70 a	13.67 $\pm$ 0.22 d
	MFC-20	8.12 $\pm$ 0.16 b	5.67 $\pm$ 0.8 a	10.02 $\pm$ 0.11 cd	4.49 $\pm$ 0.55 a	10.17 $\pm$ 0.63 d
<i>M. laxa</i>	MLX-13	5.05 $\pm$ 0.33 bc	3.98 $\pm$ 1.11 a	6.11 $\pm$ 0.13 cd	4.22 $\pm$ 0.90 ab	10.23 $\pm$ 1.1 e
	MLX-24	6.43 $\pm$ 0.07 b	4.38 $\pm$ 0.94 a	7.90 $\pm$ 0.10 bc	4.39 $\pm$ 0.52 a	9.21 $\pm$ 0.29 cd

^u Values within each row followed by the same letters are not significantly different based on Tukey's test at $P < 0.05$. SE: standard error of the mean.

^v Malt extract agar.

^w Apple juice agar.

^x Potato dextrose agar.

^y Czapeck Dox agar.

^z V8 juice agar.

could be devastating during the harvest period, but also in storage as well. *M. polystroma* could affect fruit in storage facilities and this finding may have important management implications as postharvest fungicides are not utilized in Serbia via EU regulations.

M. fructicola has been reported as less frequent apple brown rot fungi in Germany, Italy, the Czech Republic, Pennsylvania, and China (Grabke et al. 2011; Martini et al. 2015; OEPP/EPPO 2011; Peter et al. 2015; Zhu et al. 2016). In 2011, it was detected as postharvest apple

Table 4. Sporulation intensity of *Monilinia* spp. isolates on five media at 25°C in the dark

Species	Isolate	No. of conidia per cm ² of colony $\pm$ SE (log transformed) ^u				
		MEA ^v	AJA ^w	PDA ^x	CYA ^y	V8 ^z
<i>M. fructigena</i>	MFSRB-17	0	0	0	0	0
	MFSRB-22	0 a	0 a	0 a	0 a	4.37 $\pm$ 1.23 b
	MFSRB-6	0 a	0 a	0 a	0 a	4.23 $\pm$ 0.17 b
<i>M. polystroma</i>	MPSRB-36	0 a	0 a	0 a	0 a	4.17 $\pm$ 0.56 b
	MPSRB-13	0 a	0 a	0 a	0 a	4.51 $\pm$ 0.86 b
	MPSRB-20	0 a	0 a	0 a	0 a	4.56 $\pm$ 0.65 b
<i>M. fructicola</i>	MFC-1	4.49 $\pm$ 0.43 bc	3.17 $\pm$ 0.57 ab	4.8 $\pm$ 0.15 cd	3.12 $\pm$ 0.67 ab	4.87 $\pm$ 0.42 cd
	MFC-20	4.54 $\pm$ 0.53 bc	3.34 $\pm$ 0.82 ab	4.93 $\pm$ 0.75 cd	3.27 $\pm$ 0.54 ab	4.95 $\pm$ 0.52 cd
<i>M. laxa</i>	MLX-13	0 a	0 a	0 a	0 a	3.98 $\pm$ 0.41 b
	MLX-24	0 a	0 a	0 a	0 a	3.46 $\pm$ 0.24 b

^u Values within each row followed by the same letters are not significantly different based on Tukey's test at $P < 0.05$. SE: standard error of the mean.

^v Malt extract agar.

^w Apple juice agar.

^x Potato dextrose agar.

^y Czapeck Dox agar.

^z V8 juice agar.

Fig. 4. Effect of temperature on mycelial growth rate of representative *M. fructigena* isolates (MFSRB-17, MFSRB-22, MFSRB-6), *M. polystroma* (MPSRB-36, MPSRB-13, MPSRB-20), *M. fructicola* (MFC-1, MFC-20), and *M. laxa* (MLX-13, MLX-24) on potato dextrose agar in the dark. Vertical bars indicate standard error of mean (SE).

Fig. 5. Effect of pH on mycelial growth rate of *M. fructigena* isolates (MFSRB-17, MFSRB-22, MFSRB-6), *M. polystroma* (MPSRB-36, MPSRB-13, MPSRB-20), *M. fructicola* (MFC-1, MFC-20), and *M. laxa* (MLX-13, MLX-24) on potato dextrose agar at 25°C in the dark. Vertical bars indicate standard error of mean (SE).

pathogen in Serbia (Vasić et al. 2012). Since then, this study indicates the geographic spread of this species. *M. fructicola* is a pathogen capable of causing latent infections (Michailides et al. 2007), sporulates profusely (Luo et al. 2001), and can easily develop a perfect stage that plays an important role in epidemiology (Hong and Michailides 1998). Its adaptability to higher temperatures according to a study by Papavasileiou et al. (2015) can favor this pathogen. Since global temperatures are increasing, this pathogen could be more harmful for apple fruit production. Also, a recent study by Beckerman et al. (2016) reported severe crop losses over 50% on apple fruit caused by *M. fructicola* in Indiana; thus, this pathogen should be monitored in the future as well.

Monilinia spp. can be identified using traditional and molecular methods (Batra 1991; Byrde and Willetts 1977; Côté et al. 2004; OEPP/EPPO 2009). Morphological features are widely used as identification criteria, but they are time consuming and can sometimes be misleading, since variations within a species exist, leading to inaccurate identification. Cultural features of *M. polystroma*, *M. fructicola*, *M. laxa*, and one group of *M. fructigena* isolates were in accordance with other reports (Batra 1991; Byrde and Willetts 1977; OEPP/EPPO 2009; van Leeuwen et al. 2002). However, one group of *M. fructigena* isolates formed stomatal plates in culture, which is similar to *M. polystroma*. This observation is congruent with what Poniatowska et al. (2013) also noted that Polish isolates of *M. fructigena* formed stomatal plates in vitro. In light of these observations, this morphological feature may not be a reliable diagnostic character to distinguish solely between *M. fructigena* and *M. polystroma*.

In this study, we used both traditional and molecular methods for species identification, as molecular techniques are a reliable and

accurate tool to confirm species-level identification. Multiplex PCR developed by Côté et al. (2004) facilitated accurate identification and differentiation of *M. polystroma*, *M. fructigena*, *M. fructicola*, and *M. laxa*. Additionally, PCR-RFLP developed by Vasić et al. (2016) was able clearly to separate isolates of *M. polystroma* from other species as an additional layer of species-level confirmation. Phylogenetic analysis of the ITS rDNA sequence, routinely used for *Monilinia* spp. (Hrustić et al. 2015; Hu et al. 2011; Poniatowska et al. 2013; Zhu et al. 2016), confirmed the species-level identification and revealed that *M. fructigena* and *M. polystroma* are closely related.

Experiments examining fungal growth rate, sporulation, pH, and temperature in vitro for *M. fructigena*, *M. fructicola*, and *M. laxa* have been examined (Byrde and Willetts 1977). However, there is little knowledge about how these features for *M. polystroma* compare with other *Monilinia* spp. that cause apple fruit decay from Serbia. We showed that V8 and PDA were the most favorable media for the growth of all four *Monilinia* spp. Hu et al. (2011) also observed rapid growth of Chinese isolates of *M. fructicola*, *M. fructigena*, and *M. laxa* on V8 and PDA. Serbian isolates of *M. polystroma*, *M. fructigena*, and *M. laxa* only sporulated on V8 media, while Hu et al. (2011) did not observe *M. fructigena* and *M. laxa* sporulation on V8 media, indicating regional/geographic differences in these isolates. Optimal pH for *M. polystroma* mycelial growth was greater than for *M. fructigena*, as a pH between 7 and 9 were less favorable for proliferation. Similarly, Holb (2004) noticed that Japanese isolates of *M. polystroma*, originating from apple, grew fastest at pH 4.5. We observed that optimum temperature for mycelial growth of *M. polystroma* was 25°C, which is the first report for temperature

Fig. 6. Mean lesion diameter (A) and sporulation intensity (B) on inoculated apple fruit 'Idared,' 'Golden Delicious,' and 'Granny Smith' with representative isolates of *M. fructigena* (MF), *M. polystroma* (MP), *M. laxa* (MLX), and *M. fructicola* (MFC) after 5 and 6 days of incubation, respectively. Vertical bars represent standard error of mean (SE).

growth requirements of this species. The same temperature optimum was observed for mycelial growth of other tested species, which is in accordance with the literature for *M. fructigena*, *M. fructicola*, and *M. laxa* (Byrde and Willetts 1977).

Comparative analysis of *M. fructigena*, *M. polystroma*, *M. laxa*, and *M. fructicola* virulence and sporulation intensity in apple has not been conducted. In our previous investigation, *M. fructigena* was slightly more virulent than *M. polystroma* on Idared and Granny Smith cultivars, while there was no significant difference in aggressiveness on Golden Delicious (Vasić et al. 2016), which is also in agreement with the current study. *M. laxa* and *M. fructicola* were found to be less aggressive overall, which may implicate lower specialization toward apple fruit and may explain why they are more commonly found on stone fruit. Additionally, *M. fructigena*, *M. polystroma*, and *M. fructicola* were able to sporulate abundantly on all inoculated apple fruit, indicating their reproductive potential. Similarly, van Leeuwen et al. (2002) showed abundant sporulation of *M. fructigena* and *M. polystroma* on 'Conference' pear and 'Cox's Orange Pippin' apple, but quantification of *M. fructicola* sporulation on apple fruit was not conducted. Grabke et al. (2011) observed a zone of sporulation on diseased 'Jonagold' apples and artificially inoculated 'Gala' fruit. Presumably, *M. fructicola* can sporulate more abundantly and is more prevalent on stone fruit. However, based on our findings, *M. fructicola* has significant inoculum production potential and could compete with *M. fructigena* and *M. polystroma* in fungal-fungal interactions in the orchard.

On the basis of samples collected and isolates obtained during this investigation, we conclude that *M. fructigena* is the main causal agent of apple brown rot pre- and postharvest in Serbia. However, the species composition of brown rot fungi is in flux due to a variety of suspected, yet unknown factors (i.e., differences in host susceptibility, selection pressure exerted by different mode of action fungicides in the field, and varying environmental conditions). Thus, we hypothesize based on increased detection frequency and other competitive factors in the fungus (growth at different temperatures and pH) that *M. polystroma* has the potential to compete with *M. fructigena*. The presence of compact stromata on apple fruit coupled with intense melanized hyphal rinds (Vasić et al. 2016) implies that *M. polystroma* has the potential advantage to survive and produce inoculum compared with other *Monilinia* spp. Additionally, *M. polystroma* has a wider pH optima and higher temperature range than *M. fructigena*, which may contribute to increased adaptability on different hosts in diverse microenvironments. More information about the spread of newly introduced species, changes in pathogen populations, and epidemiology in complex pathosystems is needed to establish adequate disease management strategies.

Acknowledgments

This research was supported by project III 46008 financed by the Ministry of Education, Technological Development and Science, Republic of Serbia, and partially supported by USDA-ARS project no. 8042-42430-014-00D via National Program 303 Plant Diseases. Use of a company or product name by the U.S. Department of Agriculture does not imply approval or recommendation of the product to the exclusion of others that may also be suitable.

Literature Cited

Batra, L. R. 1991. World Species of *Monilinia* (Fungi). Their Ecology, Biosystematics and Control. J. Cramer, Berlin, Germany.

Beckerman, J., Albright, N., and Abbott, C. 2016. First report of brown rot (*Monilinia fructicola*) on apple (*Malus × domestica*). Plant Dis. 100:1949.

Berrie, A. M., and Holb, I. 2014. Brown rot diseases. Pages 43-45 in: Compendium of Apple and Pear Diseases and Pests, 2nd ed. T. B. Sutton, H. S. Aldwinckle, A. M. Angelo, and J. F. Walgenbach, eds. American Phytopathological Society, St. Paul, MN.

Byrde, R. J. W., and Willetts, H. J. 1977. The Brown Rot Fungi of Fruit. Their Biology and Control. Pergamon Press, Oxford.

Côté, M. J., Tardif, M. C., and Meldrum, A. J. 2004. Identification of *Monilinia fructigena*, *M. fructicola*, *M. laxa*, and *Monilinia polystroma* on inoculated and naturally infected fruit using multiplex PCR. Plant Dis. 88:1219-1225.

Day, J. P., and Shattock, R. C. 1997. Aggressiveness and other factors relating to displacement of populations of *Phytophthora infestans* in England and Wales. Eur. J. Plant Pathol. 103:379-391.

Di Francesco, A., Fruk, M., Martini, C., Jemrić, T., and Mari, M. 2015. First report of Asiatic brown rot (*Monilinia polystroma*) on apple in Croatia. Plant Dis. 99: 1181-1182.

EFSA Panel. 2011. Pest risk assessment of *Monilinia fructicola* for EU territory and identification and evaluation of risk management options. EFSA J. 9:2119.

Gams, W., van der Aa, H. A., van der Plaats-Niterink, A. J., Samson, R. A., and Stalpers, J. A. 1987. Page 136 in: CBS Course of Mycology. 3rd edition. Institute of Royal Netherlands Academy of Arts and Sciences, Centraalbureau voor Schimmelfcultures, Baarn, Netherlands.

Grabke, A., Hu, M. J., Luo, C. X., Bryson, P. K., and Schnabel, G. 2011. First report of brown rot of apple caused by *Monilinia fructicola* in Germany. Plant Dis. 95:772.

Harada, Y., Nakao, S., Sasaki, M., Sasaki, Y., Ichihashi, Y., and Sano, T. 2004. *Monilinia mumecola*, a new brown rot fungus on *Prunus mume* in Japan. J. Gen. Plant Pathol. 70:297-307.

Holb, I. J. 2004. Effect of acidity on growth rate and stroma formation of *Monilinia fructigena* and *M. polystroma* isolates. Int. J. Hort. Sci. 10:63-67.

Holb, I. J., and Chauhan, S. V. S. 2004. Effect of carbohydrate and nitrogen sources on the growth rates of *Monilinia fructigena* and *M. polystroma* isolates. J. Mycol. Plant Pathol. 35:128-131.

Holb, I. J., and Scherm, H. 2007. Temporal dynamics of brown rot in different apple management systems and importance of dropped fruit for disease development. Phytopathology 97:1104-1111.

Hong, C., and Michailides, M. J. 1998. Effect of temperature on the discharge and germination of ascospores by apothecia of *M. fructicola*. Plant Dis. 82:195-202.

Hrustić, J., Delibašić, G., Stanković, I., Grahovac, M., Krstić, B., Bulajić, A., and Tanović, B. 2015. *Monilinia* spp. causing brown rot of stone fruit in Serbia. Plant Dis. 99:709-717.

Hrustić, J., Tanović, B., Mihajlović, M., Delibašić, G., Stanković, I., Krstić, B., and Bulajić, A. 2013. First report of brown rot caused by *Monilinia fructicola* on nectarine in Serbia. Plant Dis. 97:147.

Hu, M. J., Cox, D. C., Schnabel, G., and Luo, C. X. 2011. *Monilinia* species causing brown rot of peach in China. PLoS One 6:e24990.

Ivanović, M., and Ivanović, D. 2001. Mikro i Pseudomikroze Biljaka. De-Em-Ve, Belgrade, Serbia.

Kim, Y. K., Xiao, C. L., and Rogers, J. D. 2005. Influence of culture media and environmental factors on mycelial growth and pycnidial production of *Sphaeropsis pyriputrescens*. Mycologia 97:25-32.

Luo, Y., Ma, Z., and Michailides, T. J. 2001. Analysis of factors affecting latent infection and sporulation of *Monilinia fructicola* on prune fruit. Plant Dis. 85:999-1003.

Martini, M., Di Francesco, A., Lantos, A., and Mari, M. 2015. First report of Asiatic brown rot (*Monilinia polystroma*) and brown rot (*Monilinia fructicola*) on pears in Italy. Plant Dis. 99:556.

Michailides, T. J., Luo, Y., Ma, Z., and Morgan, D. P. 2007. Brown rot of dried plum in California: new insight on an old disease. APSnet Feature.

Michailides, T. J., Ogawa, J. M., and Ogenorth, D. C. 1987. Shift of *Monilinia* spp. and distribution of isolates sensitive and resistant to benomyl in California prune and apricot orchards. Plant Dis. 71:893-896.

Munda, A. 2015. First report of brown rot on peach caused by *Monilinia polystroma* in Slovenia. Plant Dis. 99:1281.

OEPP/EPPO. 2002. First report of *Monilinia fructicola* in France. EPPO Reporting Service 2002/003. <http://archives.eppo.int/EPPOReporting/2002/Rse-0201.pdf>

OEPP/EPPO. 2009. *Monilinia fructicola*. Bull. OEPP/EPPO Bull. 39:337-343.

OEPP/EPPO. 2011. First reports of *Monilinia polystroma* in Hungary and the Czech Republic. EPPO Reporting Service. <https://gd.eppo.int/reporting/article-1703>

Papavasileiou, A., Testempasis, S., Michailides, T. J., and Karaoglanidis, G. S. 2015. Frequency of brown rot fungi on blossoms and fruit in stone fruit orchards in Greece. Plant Pathol. 64:416-424.

Peter, K., Gaskin, V., Lehman, B., and Jurick, W. 2015. First report of brown rot caused by *Monilinia fructicola* in Pennsylvania. Plant Dis. 99:1179.

Petróczy, M., and Palkovics, L. 2009. First report of *Monilinia polystroma* on apple in Hungary. Eur. J. Plant Pathol. 125:343-347.

Poniatowska, A., Michalecka, M., and Bielenin, A. 2013. Characteristic of *Monilinia* spp. fungi causing brown rot of pome and stone fruits in Poland. Eur. J. Plant Pathol. 135:855-865.

Poniatowska, A., Michalecka, M., and Puławska, J. 2016. Genetic diversity and pathogenicity of *Monilinia polystroma* - the new pathogen of cherries. Plant Pathol. 65:723-733.

Staden, R., Beal, K. F., and Bonfield, J. K. 2000. The Staden package, 1998. Methods Mol. Biol. 132:115-130.

Tamura, K., Stecher, G., Peterson, D., Filipowski, A., and Kumar, S. 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol. Biol. Evol. 30: 2725-2729.

Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., and Higgins, D. G. 1997. The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Res. 25: 4876-4882.

van Leeuwen, G. C. M., Baayen, B. P., Holb, I. J., and Jeger, M. J. 2002. Distinction of the Asiatic brown rot fungus *Monilinia polystroma* sp. nov. from *M. fructigena*. Mycol. Res. 106:444-451.

- van Leeuwen, G. C. M., Stein, A., Holb, I., and Jeger, M. J. 2000. Yield loss in apple caused by *Monilinia fructigena* (Aderh. & Ruhl.) Honey, and spatio-temporal dynamics of disease development. *Eur. J. Plant Pathol.* 106:519-528.
- Vasić, M., Duduk, N., and Ivanović, M. 2013. First report of brown rot caused by *Monilinia polystroma* on apple in Serbia. *Plant Dis.* 97:145.
- Vasić, M., Duduk, N., Ivanović, M. M., Obradović, A., and Ivanović, M. S. 2012. First report of brown rot caused by *Monilinia fruticola* on stored apple in Serbia. *Plant Dis.* 96:456.
- Vasić, M., Duduk, N., Vico, I., Rančić, D., Pajić, V., and Backhouse, D. 2016. Comparative study of *Monilinia fructigena* and *Monilinia polystroma* on morphological features, pathogenicity, RFLP analysis and histopathology. *Eur. J. Plant Pathol.* 144:15-30.
- Villarino, M., Egüen, B., Lamarca, N., Seggera, J., Usall, J., Melgarejo, P., and De Cal, A. 2013. Occurrence of *Monilinia laxa* and *M. fructigena* after introduction of *M. fruticola* in peach orchards in Spain. *Eur. J. Plant Pathol.* 137:835-845.
- White, T. J., Bruns, T., Lee, S., and Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Pages 315-322 in: *PCR Protocols: A Guide to Methods and Applications*. M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White, eds. Academic Press, San Diego, CA.
- Zhu, X. Q., Chen, X. Y., and Guo, L. Y. 2011. Population structure of brown rot fungi on stone fruits in China. *Plant Dis.* 95:1284-1291.
- Zhu, X. Q., Niu, C. W., Chen, X. Y., and Guo, L. Y. 2016. *Monilinia* species associated with brown rot of cultivated apple and pear fruit in China. *Plant Dis.* 100:2240-2250.