Bedding Plant Production and the Challenge of Fungal Diseases

Vladimiro Guarnaccia,^{1,2} Francesca Peduto Hand,³ Angelo Garibaldi,¹ and M. Lodovica Gullino^{1,†}

¹ Centre for Innovation in the Agro-Environmental Sector, AGROINNOVA, University of Torino, Largo Braccini 2, 10095 Grugliasco (TO), Italy

² Department of Agricultural, Forest and Food Sciences (DISAFA), University of Torino, Largo Braccini 2, 10095 Grugliasco (TO), Italy

³ Department of Plant Pathology, The Ohio State University, Columbus, OH 43220, U.S.A.

Keywords: foliar diseases, soilborne, soilless, soil suppressiveness, sustainable management

Bedding Plants, Production, and Market

Bedding plants are fast-growing annuals or herbaceous perennials (Fig. 1) used outdoors in flowerbeds or planters, in gardens, on balconies, or around buildings. The bedding plant industry is mainly local, with very limited export of the products but including massive distribution direct to consumers by large chain stores. The choice of plants often reflects trends and consumers' taste at the local level.

The total value of bedding and garden plants in the United States was \$2.16 billion in 2018, which was up 7% compared with 2015, and represented 47% of the total wholesale value of the floriculture crops sector (USDA 2019). In Europe, the production of bedding plants encompasses a very significant part of the ornamental plants sector that registered a wholesale value of €2.2 billion in 2019, with The Netherlands accounting for 30% of the total production, followed by Italy and Germany, each accounting for 13% (European Commission 2020).

Bedding plant cultivation in temperate climates such as those of both the United States and Europe starts under greenhouse conditions in late winter to early spring. Plants are grown in a variety of formats: in flats, pots, or hanging baskets. Sometimes different plant species are grown and sold as mixed items in the same container, for use as a hanging basket or patio container. A very wide variety of plant species is often grown within the same greenhouse range. Plants are sold in garden centers, big box stores, farm stands, or street markets and planted at the beginning of the growing season, as soon as the risk of frost has passed (Daughtrey and Buitenhuis 2020). Most genera and species are subject to a number of diseases, whose severity varies according to the species and cultivar as well as the characteristics of the production process of the industry. The production cycle is generally shorter in comparison with other floriculture crops, making disease management less complex, because plant

pathogens have less time to cause severe losses during the production time. The quality of the cultivation practices strongly influences the health status of the finished plant.

Recent Changes in this Production Sector

Genera and species grown. Many genera and species are grown as bedding plants, with a strong influence of local trends and continuous changes due to changes in customers' preferences. In addition to the most familiar and popular annuals, herbaceous perennial species are increasingly grown, leading to a very high total number of cultivated species. The species shown in Figure 1 are some of those commonly produced in the United States and Europe; nevertheless, different countries are specialized in production of certain species such as impatiens and begonia in the United States, bulb species (e.g., tulip, lily, hyacinth, gladiolus, narcissus, and so on) in The Netherlands, and cyclamen and Asteraceae (e.g., coneflower, chrysanthemum, and so on) in Northern Italy. New cultivars as well as new species are continuously introduced each year, making not only the market but also the phytosanitary situation quite variable and dynamic.

Production practices, including cultural methods, substrates, and inputs in relation to plant health. Because the industry is localized and offers an abundant variety of plant choices, it is not uncommon to find hundreds of species and cultivars originating from different specialist propagators grown in the same greenhouse. Propagules may be sourced from many parts of the world but then distributed locally after growth to transplant size. The practice of using seeds or unrooted cuttings from global sources strongly increases the risk of introduction of pathogens that may quickly spread from infected to healthy plants. Production practices, which are very fragmented and strongly related to the size of the facility, influence the health status of the final product.

Propagative material. Bedding plants can be started from seeds or cuttings and both of them can be a source of pathogens. In recent years, well-rooted, transplant-ready seedlings or cuttings in plug trays replaced the seeding step for growers of many crops such as begonia and dianthus. Global distribution of unrooted cuttings to rooting stations or direct to growers who finish the crops has sped up the process of introducing new lines. For example, in the United States, 10% of petunias are propagated from cuttings (Daughtrey and Buitenhuis 2020). Vegetative propagation material is shipped to growers directly as unrooted cuttings or first rooted in rooting stations and then sold to growers as rooted cuttings. Such early steps in plant production are very critical, and the health status of the final

[†]Corresponding author: M. L. Gullino; marialodovica.gullino@unito.it

Funding: This work was supported by the Italian Ministry of Education, Universities and Research (MIUR), Local research (ex 60%), and United States Department of Agriculture–National Institute of Food and Agriculture Hatch project number 1020446.

The author(s) declare no conflict of interest.

Accepted for publication 29 October 2020.

crop strongly depends upon the quality of the measures adopted at the propagation, rooting station, or plug production steps, before the young plant (as seedling or rooted cutting) reaches the finisher (Daughtrey and Buitenhuis 2020).

Cultivation systems and growing media. Soilless culture has been gaining popularity in the ornamental industry (Gullino and Garibaldi 2007) which, in most cases, allows starting a production cycle completely free of pathogens. Although such a growing system is ultimately intended to reduce production costs and maximize profits, precise environment and nutrition controls that push plants to new limits of growth and productivity can generate chronic stress conditions, which are difficult to measure, but are apparently conducive to diseases caused by pathogens such as *Penicillium* or *Pythium* spp. (Garibaldi and Gullino 2010).

A variety of growing media has been developed which provides growers with very broad options that can be tailored to the specific crop grown and the type of cultivation adopted. During the past two decades, much attention has been devoted to the selection of media that not only support good growth but are also suppressive of important pathogens (Pugliese et al. 2015).

Case studies

This section aims to review the most important groups of pathogens causing severe diseases on bedding plants, with special emphasis on those most recently reported (Fig. 2).

Fusarium wilts. The genus *Fusarium*, which was recently separated into several *Fusarium*-like genera (i.e., *Bisifusarium* [*Fusarium dimerum* species complex (SC)] or *Neocosmospora* [*F. solani* SC]), includes highly important plant pathogens affecting several hosts (Guarnaccia et al. 2018; Lombard et al. 2015; O'Donnell et al. 2008). *Fusarium* spp. are well-known causal agents of diseases of ornamental plants, including bedding plants and other herbaceous as well as woody ornamentals (Guarnaccia et al. 2019a; Gullino et al. 2015a), occurring at all plant production stages.

Fusarium wilts, caused by several formae speciales of *F. oxysporum* (Booth 1984) or novel species members of the *F. oxysporum*

SC recently described (Lombard et al. 2019), can generate economic losses in commercial greenhouses where a wide range of ornamental species are cultivated (Elmer 2008; Wang and Jeffers 2002).

Bedding ornamentals that are susceptible to *Fusarium* wilts include a number of species in the genera *Aster*, *Begonia*, *Dianthus*, *Cyclamen*, *Eustoma*, and *Gerbera* (Gullino et al. 2015a; Minuto et al. 2007; Sinclair and Lyon 2005). Woody ornamentals such as *Ailanthus*, *Albizia*, and *Bougainvillea* spp. can also be affected (Guarnaccia et al. 2019a; Sinclair and Lyon 2005).

F. oxysporum f. sp. *chrysanthemi* causes wilt of chrysanthemum (Armstrong et al. 1970; Jackson and McFadden 1961). Similarly, wilts caused by the same formae speciales were reported on three hosts in the Asteraceae family (Fig. 3): gerbera daisy (*Gerbera jamesonii*), Paris daisy, (*Argyranthemum frutescens*), and African daisy (*Osteospermum* sp.) (Garibaldi and Gullino 2012b; Garibaldi et al. 2004; Li et al. 2010; Minuto et al. 2007). A subsequent study revealed the presence of three physiological races in *F. oxysporum* f. sp. *chrysanthemi* causing different pathogenic reactions on various cultivars of the same host species (Troisi et al. 2013). Race 1 of this formae speciales has also been found associated with chrysanthemum in both the United States (Horst and Nelson 1997) and Italy (Garibaldi and Gullino 2012b). Moreover, *F. oxysporum* f. sp. *chrysanthemi*, has been reported as pathogenic on orange coneflower (*Rudbeckia fulgida*), which emphasizes its expanded host range (Table 1) (Garibaldi et al. 2017b; Matić et al. 2018). Gerbera and argyranthemum plants cultivated in Italy and Brazil have been found infected by *F. oxysporum* f. sp. *tracheiphilum* (Garibaldi and Gullino 2012b; Troisi et al. 2010b).

Fusarium wilt caused by *F. oxysporum* f. sp. *cyclaminis* commonly occurs on cyclamen (*Cyclamen* spp.), where it causes major losses. This has led several growers in the United States and Europe to abandon this crop (Fig. 4) (Elmer and Daughtrey 2012; Tompkins and Snyder 1972). No races have been identified. However, three clones have been identified based on vegetative compatibility groups and restriction fragment length polymorphism analysis (Woudt 1995).

The dianthus industry in Italy, France, and Spain suffered a strong reduction in terms of acreage cultivated due to *F. oxysporum* f. sp.

Fig. 1. Examples of the wide variety of species grown as bedding plants.

dianthi and *F. redolens* f. sp. *dianthi* (Garibaldi and Gullino 2012a). Moreover, *F. oxysporum* f. sp. *dianthi* also has been reported on plants cultivated in Colombia and the United States (Chase et al. 2018; Garibaldi et al. 2011b).

Lisianthus (*Eustoma grandiflorum*) is also affected by *F. oxysporum* f. sp. *eustomae* (Li et al. 2010) and the presence of three groups was demonstrated based on genetic diversity and vegetative compatibility (Bertoldo et al. 2015). *F. oxysporum* f. sp. *eustomae* is the typical example of a forma specialis restricted to only one host species, *lisianthus*. However, other formae speciales such as the abovementioned *F. oxysporum* f. sp. *chrysanthemi* have a broader host range (chrysanthemum and coneflower). Similarly, one host plant can be susceptible to multiple formae speciales, as occurs for *G. jamesonii*, which can show typical wilt symptoms caused by *F. oxysporum* f. sp. *chrysanthemi* and *F. oxysporum* f. sp. *tracheiphilum* (Garibaldi et al. 2004, 2008c; Minuto et al. 2007; Troisi et al. 2013). *F. oxysporum* causing wilting on *Lewisia cotyledon* and *F. oxysporum* f. sp. *ranunculi* infecting *Ranunculus* spp. have been reported in Italy (Garibaldi and Gullino 1985).

Oomycetes. Several species of oomycetes in the genera *Phytophthora* and *Pythium* are well known causes of root and foliar diseases on a wide range of ornamental plants (Aiello et al. 2018; Erwin and Ribeiro 1996; Ferguson and Jeffers 1999; Jung et al. 2016; Prigigallo et al. 2015; Schwingle et al. 2007; Werres et al. 2001), and cause billions of dollars in losses to crops worldwide (Kamoun et al. 2015), including ornamentals.

Oomycete diseases are common in ornamental greenhouses where warm temperatures, high humidity, and densely grown plants represent favorable conditions for pathogen growth and sporulation (Donahoo and Lamour 2008). Moreover, irrigation sources and practices play a key role in pathogen spread (Hong and Moorman 2005; Yamak et al. 2002).

Surveys of nurseries and greenhouses for species of *Phytophthora* have been conducted in several states in the United States in the last 15 years (Donahoo and Lamour 2008; Hong et al. 2006; Hu et al. 2008; Hwang and Benson 2005; Olson and Benson 2011; Olson et al. 2013; Schwingle et al. 2007; Warfield et al. 2008; Yakabe et al. 2009).

Phytophthora nicotianae infects approximately 260 genera, including many herbaceous and woody plants, and is present in nurseries and greenhouses worldwide (Cline et al. 2008). This species is the most commonly encountered in bedding plants in the United States, where it causes extensive losses both in production and in the landscape (Olson et al. 2013). The pathogen causes root and crown rot on *G. jamesonii* and pansy, as well as aerial blight on annual vinca (*Catharanthus roseus*), calibrachoa (*Calibrachoa* × *hybrida*), and, most recently, on lobelia (*Lobelia erinus*) (Lin et al. 2018). The symptoms caused by *P. nicotianae* develop at temperatures higher than 28°C but the oomycete colonization occurs at lower temperatures (Chase et al. 2018).

P. tropicalis is a relevant pathogen on numerous bedding ornamental plants such as annual vinca (*C. roseus*) (Hao et al. 2010), cyclamen (*Cyclamen persicum*) (Gerlach and Schubert 2001), gloxinia (*Sinningia speciosa*), verbena (*Verbena* × *hybrida*) (Olson and Benson 2011), begonia (*Begonia* sp.), gerbera (*G. jamesonii*), lupine (*Lupinus albus*), and dusty miller (*Senecio bicolor*) (Hong et al. 2008).

In gerbera, *P. cryptogea* is responsible for severe root rot resulting in stunting, wilting, and death of plants (Garibaldi et al. 2003b). However, gerbera is grown in soilless mixes in both the United States and Europe (Tognoni and Incrocci 2003); thus, diseases observed in hydroponic production have not been revealed when this species is grown in peat-based mixes. Beyond gerbera, *P. cryptogea* is reported worldwide as causing root and crown rot on a number of ornamental species, including *Begonia* spp., *Chrysanthemum morifolium*, petunia, *Salvia officinalis*, and *Verbena hybrida* (Chase et al.

Fig. 2. Some bedding plant diseases caused by different groups of pathogens.

2018; Garibaldi et al. 2015c). *P. drechsleri* is another *Phytophthora* sp. causing crown rot on bedding plants such as *Calibrachoa* spp., *Celosia argentea*, *G. jamesonii*, and *Helichrysum bracteatum*, among others. Recently, the newly described species *P. chrysanthemi* was reported on chrysanthemum in the United States (Lin et al. 2017), Europe (Tomic and Ivic 2015), and Japan (Naher et al. 2011).

The genera *Pythium* and *Globisporangium* are often associated with damping-off and crown and root rot diseases during propagation, causing the death of seedlings and cuttings (Daughtrey and Benson 2005; Garzón et al. 2011; Moorman et al. 2002). Primary inoculum of *Pythium* and *Globisporangium* spp. is commonly present in the soil and usually is introduced into greenhouses on used

Fig. 3. Evaluation of different Asteraceae species for susceptibility to *Fusarium oxysporum* f. sp. *chrysanthemi*.

Table 1. Pathogenicity of *Fusarium oxysporum* strains isolated from orange coneflower on different bedding plant hosts in the family Asteraceae^x

Isolates	Trial 1 ^y				Trial 2 ^z		
	Cone	African	ChrysSQ	Paris	Cone	ChrysPC	Gerbera
<i>F. oxysporum</i> f. sp. <i>chrysanthemi</i>							
DB32	59.2 b (S)	55.0 b (S)	21.7 b (MS)	84.2 b (HS)	37.5 ab (MS)	35.0 ab (MS)	87.5 b (HS)
FC32	64.2 b (S)	80.0 b (S)	19.2 b (PR)	87.5 b (HS)	100.0 c (HS)	40.0 b (MS)	53.1 b (S)
DB23	60.0 b (S)	65.0 b (S)	39.2 b (MS)	95.8 b (HS)	47.5 a-c (MS)	35.0 ab (MS)	90.6 b (HS)
DB43	77.5 b (S)	69.2 b (S)	44.2 b (MS)	92.5 b (HS)	43.8 a-c (MS)	30.0 ab (MS)	68.8 b (HS)
<i>F. oxysporum</i> IT22	64.2 b (S)	51.7 b (S)	38.3 b (MS)	92.5 b (HS)	37.5 ab (MS)	55.0 b (S)	71.9 b (HS)
<i>F. oxysporum</i> IT23	nt	nt	nt	nt	56.3 a-c (S)	50.0 b (MS)	62.5 b (HS)
<i>F. oxysporum</i> IT24	63.3 b (S)	75.0 b (S)	30.0 b (MS)	93.3 b (HS)	56.3 a-c (S)	50.0 b (MS)	75.0 b (HS)
<i>F. oxysporum</i> IT25	66.7 b (S)	64.2 b (S)	36.7 b (MS)	96.7 b (HS)	68.8 bc (S)	45.0 b (MS)	62.5 b (HS)
Noninoculated	0.0 a	0.0 a	0.0 a	0.0 a	0.0 a	0.0 a	0.0 a

^x Data are expressed as disease severity (0 to 100) and compared with reference strains of *F. oxysporum* f. sp. *chrysanthemi* at the end of two trials (Matić et al. 2018). Values followed by the same letter are not significantly different ($P = 0.05$) according to Tukey's multiple range test; nt = not tested. R = resistant (disease index 0 to 5), PR = partially resistant (disease index 6 to 20), MS = moderately susceptible (disease index 21 to 50), S = susceptible (disease index 51 to 75), and HS = highly susceptible (disease index 76 to 100).

^y Trial 1: Cone = orange coneflower, African = African daisy (Variegata Bianca), ChrysSQ = chrysanthemum (SupraQuivre), and Paris = Paris daisy (Stella 2000).

^z Trial 2: Cone = orange coneflower, ChrysPC = chrysanthemum (Polochon Cognac), and Gerbera = Gerbera (Carambole).

containers. Moreover, seeds of bedding plants can be colonized by these oomycetes and contribute to inoculum transmission (Faust et al. 2017). *Pythium aphanidermatum*, *Globisporangium irregulare* (former *P. irregulare*), and *G. ultimum* (former *P. ultimum*) are the most common pathogens in the “Pythium root rot” group causing diseases of ornamental plants (Chase et al. 2018; Garibaldi et al. 2009c; Guarnaccia et al. 2015). Among the bedding plants most commonly affected by these oomycetes are *Pelargonium* spp., *Catharanthus roseus* (annual vinca), and *Impatiens* spp. In addition to root rot, affected plants may show stunting, yellowing, or wilting of the foliage; occasionally, stem or crown rot or canker may occur in absence of root rot (Chase et al. 2018). When conditions are wet and humid, rot from the roots can proceed up to the basal leaves, where presence of white hyphae can be observed (Moorman et al. 2002).

Downy mildews are also major threats: *Basidiophora*, *Hyaloperonospora*, *Peronospora*, and *Plasmopara* are the most common oomycete genera causing downy mildew in bedding plants. Patches of chlorosis and necrosis are typical symptoms produced on the upper leaf surface, while prolific white sporulation can be observed on the leaf underside. *Coleus*, *Impatiens*, *Matthiola*, and *Viola* spp. are affected by *Peronospora belbahrii*, *Plasmopara obducens*, *Peronospora parasitica*, and *Hyaloperonospora parasitica*, respectively (Daughtrey et al. 2006; Koike 2000; Palmateer et al. 2008; Rivera et al. 2016). Other common hosts in the United States include salvia (*Peronospora lamii*) (Choi et al. 2009), snapdragon (*P. antirrhini*; Byrne et al. 2005), and verbena (*P. verbenae*; Braun et al. 2009). *P. digitalis* and *P. arthurii* caused outbreaks on *Digitalis purpurea* (Fig. 5) and *Oenothera biennis* in different countries of United States and Europe, and in New Zealand (Belbahri et al. 2005; Garibaldi et al. 2013b, 2018a).

***Rhizoctonia solani*.** Diseases of ornamental plants caused by *Rhizoctonia* spp. commonly occur in the greenhouse, nursery, and landscape (Aiello et al. 2017; Daughtrey and Benson 2005). *Rhizoctonia solani* (teleomorph *Thanatephorus cucumeris*) is a necrotrophic fungus in the division Basidiomycota (Sneh et al. 1991) that causes root, crown, and foliar diseases on a wide range of agronomic crops (Couch 1995; Sneh et al. 1991). *Rhizoctonia* spp. are classified into three groups based on the number of nuclei per cell (Ogoshi 1996): multinucleate, binucleate, and uninucleate. Different anastomosis groups are present within each group. *R. solani* is highly widespread, with a host range that includes over 500 plant species (Farr and Rossman 2020). Root and stem rot, seedling damping-off, leaf spot, and foliar web blight are reported worldwide on economically important ornamental species such as *Begonia*, *Catharanthus*, *Chrysanthemum*, *Dianthus*, *Impatiens*, *Osteospermum*, *Pelargonium*, *Petunia*, *Rosa*, and *Verbena* spp., among others (Benson and Cartwright 1996; Chase 1991; Holcomb and Carling 2000; Hyakumachi et al. 2005; Rinehart et al. 2007). Symptom development is favored by warm temperatures and high humidity. Damping-off of young

seedlings or cuttings in bedding plants commonly develops in a circular pattern within a flat in production or within a bed in the landscape, where the mycelium appears whitish-brown and spiderweb-like on the soil or on the basal leaves (Chase et al. 2018). Brown to black lesions on the crown and base of stems are typical symptoms caused by *Rhizoctonia* spp. on *Begonia*, *Impatiens*, and *Pelargonium* spp., leading to stunting or death of infected plants. *R. solani* is a very limiting factor in ornamental nurseries in Italy, where several *Rhizoctonia* diseases have been reported in recent decades (Aiello et al. 2008, 2009a,b; Garibaldi et al. 2003a, 2006, 2009d,e, 2010b, 2013a). Moreover, several bedding plants cultivated in private gardens were reported as susceptible to *R. solani* infections: web blight and crown rot are reported on different *Campanula* and *Lychnis* spp. (Fig. 6) (Garibaldi et al. 2015a,d); crown and stem rot on *Abelmoschus manihot* and *Echinacea purpurea* (Garibaldi et al. 2019f, 2020e); and leaf blight on *Aquilegia*, *Digitalis*, *Hosta*, *Lupinus*, and *Salvia* spp. (Garibaldi et al. 2009a,d,e,f, 2010b).

***Alternaria* leaf spots.** Leaf spot caused by *Alternaria* spp. is one of the most frequent diseases of bedding plants propagated through seeds. *Alternaria* is a cosmopolitan genus that includes several saprophytic and pathogenic species, recently revised into 26 sections (Woudenberg et al. 2013). *Alternaria* sect. *Alternaria* includes major pathogens and contains species that are differentiated based on morphological characteristics (i.e., small and concatenated conidia).

Alternaria alternata is normally a saprophyte; however, it is also able to cause disease on several ornamental plants (Thomma 2003). *A. nobilis*, *A. dianthicola*, *A. saponariae*, *A. tagetica*, and *A. zinniae* are other *Alternaria* spp. reported on bedding plants such as dianthus, *Saponaria* spp., zinnia, and marigold (Garibaldi et al. 2013c; Chase et al. 2018). These species present a gray to olivaceous-black mycelium on artificial media, with typical muriform conidia produced in chains.

Extensive research has been conducted on this disease related to particular hosts such as *Zinnia* and *Pelargonium* spp. reported from throughout the world. However, *Alternaria* leaf spots are also observed in a high number of new ornamental plant hosts (Matić et al. 2020). Outbreaks of leaf spots on new ornamental, medicinal, and aromatic plants caused by *Alternaria* spp. massively occurred during the last two decades in various areas of Italy, Greece, Serbia, the United States, Canada, Brazil, Uruguay, Mexico, South Africa, Asia (China, South Korea, India, Iraq, Iran, and Pakistan), and Australia (Farr and Rossman 2020). Common symptoms consist of small, circular, brown to purple (depending on the host species), water-soaked leaf spots which progressively become larger and necrotic. On severely infected plants, defoliation is observed and, occasionally, plant death. *Alternaria* spp. can be transmitted by seeds, and the recent globalization of the seed market is obviously favoring this process (Gullino et al. 2014). Plant debris, weeds, and seeds colonized

Fig. 4. Symptoms of Fusarium wilt on cyclamen.

Fig. 5. Symptoms of downy mildew caused by *Peronospora digitalidis* on *Digitalis purpurea*.

by *Alternaria* spp. are the sources of overwintering inoculum (Laemmle 2001).

Other hosts include ornamental and medicinal plants such as *Rudbeckia fulgida* (Matić et al. 2020), *Mentha × piperita*, *Salvia elegans*, and *Echinacea purpurea* (Garibaldi et al. 2018b,e,f; 2020d). More recently, symptoms caused by *A. alternata* occurred on *Plectranthus scutellarioides*, *Ceratostigma willmottianum*, *D. purpurea*, *Phlox maculata* (Fig. 7), and *Alcea rosea* (Fig. 8) (Garibaldi et al. 2019d,e, 2020a,b,d). *Alternaria arborescens* was reported as a pathogen of *Symphyotrichum novi-belgii* (Garibaldi et al. 2020c). *Campanula medium* and *C. rapunculoides* recently displayed *Alternaria* leaf spot as small, light-brown, circular spots, becoming irregular and dark brown and enlarging to cover the whole leaf surface, leading to severe plant defoliation (Garibaldi et al. 2019b). Similarly, *Salvia dorisiana*, *S. elegans*, and *S. involucrata* were recently affected by *Alternaria* leaf spot in Northern Italy. Chlorosis and irregular brown necrosis were observed on the margins and blades of infected leaves along with leaf drop (Garibaldi et al. 2018b, 2019c, 2020f; Matic et al. 2020). *A. tenuissima* is a further emerging species recently reported on several bedding plants such as *Begonia semperflorens*, *Coreopsis lanceolata*, and *Iris* spp. in China (Li and Liu 2019; Li et al. 2010; Sun et al. 2019; Zhang et al. 2020) and *Echeveria* spp. in Korea.

Anthrachnose. The genus *Colletotrichum* is considered one of the 10 most economically important plant pathogens in the world (Dean et al. 2012). *Colletotrichum* spp. are present in a broad range of environments including open fields, greenhouses, and postharvest warehouses (Cannon et al. 2012).

Until recently, anthracnose was not considered a major disease problem on bedding plants. However, in the last few years, several disease reports have been published highlighting an increased incidence of *Colletotrichum* spp. on ornamental hosts (Chase et al. 2018; Guarnaccia et al. 2019c; McMillan and Graves 1996; Polizzi et al. 2011). Among these, *Campanula*, *Cyclamen*, and *Salvia* spp. are more severely affected by *Colletotrichum* spp. (Garibaldi et al. 2016c). Round, brown, lightly depressed leaf spots are observed on infected cyclamen leaves. The spots can be numerous and small or may coalesce to form large necrotic patches. Leaf pedicels and flowers can also be affected (Liu et al. 2011). Brown to black, necrotic, circular lesions on leaves of *Campanula trachelium* and *C. rapunculoides* were detected in Italy, and strains of *Colletotrichum lineola* and *C. nymphaeae* were found associated with the disease (Guarnaccia et al. in press). On *Salvia*, *C. fiorinae*, *C. fruticola* or *C. nigrum*, and *C. bryoniicola* are responsible for leaf anthracnose on *Salvia leucantha*, *S. greggii*, and *S. nemorosa*, respectively. Although these three species of bedding plants are members of the same genus, the *Colletotrichum* spp. found to infect them belong to four different SCs, demonstrating high intraspecific variability associated with this disease (Guarnaccia et al. 2019c). Furthermore, *C. fuscum*

has been recently found in Italy as causing leaf anthracnose on *Coreopsis lanceolata* (Fig. 9) (Guarnaccia et al. in press).

Phoma-like leaf spots. The term “Phoma-like” is a general concept used to identify a broad group of fungal species commonly found on herbaceous stems and leaves (Aveskamp et al. 2010; Chen et al. 2015), including species with no adequate characteristics to be assigned to a genus level (Crous et al. 2004). A recent revision resolved 36 genera, including plant pathogens causing plant infection and seeds infestation.

R. fulgida is one of the bedding plants known as a host of Phoma-like diseases: infected leaves show extensive and irregular, dark-

Fig. 7. Severe attack of *Alternaria alternata* causing leaf spot on *Phlox maculata*.

Fig. 6. Collapse of *Lychnis coronaria* plant caused by *Rhizoctonia solani*.

Fig. 8. Leaf spot caused by *Alternaria alternata* on *Alcea rosea*.

brown, necrotic lesions, slightly sunken and with a well-defined border. Lesions measure 0.5 to 3 mm in diameter and occasionally coalesce to cover the entire leaf, with consequent leaf drop. Stems can also be affected, leading to death of the plant (Garibaldi et al. 2010a). Similarly, *S. nemorosa* and *S. greggii* can present leaf spots caused by *Phoma herbarum* (Fig. 8) and *P. exigua* (Garibaldi et al. 2016b; Gilardi et al. 2017). *P. novae-verbascicola*, a major species in terms of number of bedding plant hosts, causes typical leaf spots on two *Verbascum* spp., *Verbascum blattaria* and *V. nigrum*, where it was reported also as colonizing seeds (Bertetti et al. 2016; Garibaldi et al. 2013d, 2014a). The genus *Campanula* includes several species reported as hosts of Phoma-like fungi, such as *Campanula trachelium* and *C. medium*, infected by *Stagonosporopsis trachelii* (Garibaldi et al. 2015b, 2017a). In India, *P. costarricensis* has been reported as responsible of leaf spot on *Delphinium malabaricum* (Patil et al. 2012).

Powdery mildews. Many different genera of powdery mildew fungi occur on bedding plants, among which *Erysiphe*, *Golovinomyces*, and *Podosphaera* are prevalent (Chase et al. 2018). Primary infections start from overwintering hyphae in dormant buds, or from ascospores landing on plant tissues (Bélanger et al. 2002). Although the pathogens are not usually seen in greenhouses, they are commonly found in landscapes on bedding plants planted outdoors at the end of the growing season (Chase et al. 2018). The pathogens produce hyphae on the surface of infected tissues that give rise to conidiophores on which asexual propagules (i.e., conidia) are produced (Braun and Cook 2012) and dispersed via air currents and splashing water. Extensive periods of dry conditions and free moisture are both detrimental to conidia germination (Bélanger et al. 2002).

Although powdery mildew fungi do not usually kill their hosts, infected plants show reduced development and photosynthetic ability (McGrath and Shishkoff 2001) and can become unsightly and unmarketable (Chase et al. 2018). In some crops such as *Phlox* and *Dahlia*, severely affected leaves become distorted, dry out, and fall off prematurely (Chase et al. 2018; Farinas et al. 2019; Garibaldi et al. 2011a). Bedding plants frequently affected by powdery mildew include *Begonia*, *Campanula*, *Dahlia*, *Gerbera*, *Petunia*, *Phlox*, *Salvia*, *Verbena*, *Viola*, and *Zinnia* spp. (Chase et al. 2018).

Erysiphe begoniicola (formerly *Microsphaera begoniae*) is a common powdery mildew of begonia on five continents (Amano 1986; Cho et al. 2017; Crous et al. 2000; Pennycook 1989). Although not officially reported, the pathogen is also probably present in North America (Chase et al. 2018). *E. cruciferarum* was reported on *Cleome hassleriana* in Italy (Garibaldi et al. 2009f). *Podosphaera xanthii* is the major powdery mildew species infecting petunia in both the United States and Europe (Kiss et al. 2008), where *Golovinomyces orontii* is also sometimes found associated with this host (Garibaldi et al. 2007). Another species recently reported on petunia in the United States and also present in Europe is *Euoidium longipes* (syn. *Oidium longipes*) (Kiss et al. 2008; Kiss and Bereczky 2011). *G. orontii* was also reported on *Abelmoschus manihot*, *Campanula glomerata*, and *C. rapunculoides* (Fig. 10), while *G. magnicellulatus* was reported associated with powdery mildew on *Phlox paniculata* in Italy (Garibaldi et al. 2012a, 2016a, 2018c, 2019a) and the United States (Baysal-Gurel et al. 2020; Farinas et al. 2020). The most common powdery mildew reported on zinnia worldwide is *G. cichoracearum* (Chase et al. 2018). This species has a very wide host range; it infects more than 2,500 hosts, including many ornamentals in the Asteraceae family such as gerbera, orange coneflower, purple coneflower, and Paris daisy (Garibaldi et al. 2008a,b, 2018d; Troisi et al. 2010a).

Rust and smut diseases. Several bedding plants species are affected by rust fungi, which are obligate parasites and often host specific. *Puccinia lagenophorae* is present in both the United States and Europe, appearing as raised bumps with orange aeciospores and causing leaf wilt and plant death. This rust species is pathogen of hundreds of plants, such as *Bellis*, *Calendula*, and *Gazania* spp. (Hernández and Hennen 2003). *P. chrysanthemi* and *P. horiana* affect *Chrysanthemum* spp. while *P. pelargonii-zonalis* affects *Pelargonium* spp. (Chase et al. 2018). *Pucciniastrum circaeae* and *P. epilobii* are known as pathogen of *Fuchsia* spp. in Italy, Chile, and

the United States as well as Indonesia (Ferrada et al. 2020; Garibaldi et al. 2012b; Wahyuno et al. 2012), while *Coleosporium campanulae* is a pathogen of *Campanula rapunculoides* (Fig. 11) (Garibaldi et al. 2017c).

White smut diseases, often considered of minor importance, have a great impact on the cultivation of particular bedding plant species. This is the case of leaf smut of *Gaillardia* spp. (Fig. 12) that was found in California caused by *Entyloma gaillardianum* (Glawe et al. 2010) and in Virginia caused by *E. polysporum* (Hong and Banko 2003). The pathogen *E. gaillardianum* causes round, flat, white to tan leaf spots with indistinct margins up to 1 cm in diameter, that turn to brown and necrotic, followed by necrosis of the entire leaf, also reported also in Europe (Italy) (Garibaldi et al. 2018g). Moreover, *E. dahlia* and *E. calendulae* have been reported on dahlia and cosmos, respectively (Chase et al. 2018).

Sustainable Disease Management

It is well known that unique features of the ornamentals industry such as the high unit value strongly influence disease management (Baker and Linderman 1979). This, of course, also applies to the bedding plants sector (Daughtrey and Benson 2005; Daughtrey and Buitenhuis 2020; Gullino and Garibaldi 2007). Seeds and cuttings contaminated by several pathogens are often introduced into the production system. When cuttings originate from facilities located in third-world countries, the risk of introducing new invasive pathogens into new areas increases (Gullino and Garibaldi 2007).

Plant health problems must be addressed at the production level without negatively affecting the environment. Moreover, environmental regulations and water use restrictions are often very limiting factors for both producers and consumers. Effective management tools that have low environmental impact and are relatively inexpensive are the preferred choice. This is particularly challenging in the case of crops having such an important aesthetic value. This

Fig. 9. Leaf anthracnose caused by *Colletotrichum fuscum* on *Coreopsis lanceolata*.

challenge has led to a comprehensive revision of disease management strategies, which today are almost completely based on prevention (Fig. 13). That was considered a dream 40 years ago; however, now this brings the ornamental industry to the forefront of the development of innovative disease management strategies.

Preventive Measures

Disease management approaches have shifted in recent years from curative to preventive, not just by adopting sanitation practices but also enhancing crop resilience (Fig. 13) (Gullino and Garibaldi 2016; Kruidhof and Elmer 2020). Disease management is more and more coupled with greenhouse energy saving and technologies for sensing, monitoring, and aiding in decision making (Kruidhof and Elmer 2020).

Clean stock and diagnostic tools. Propagation material can be a source of pathogens and must be pathogen free. Pathogens can be exterior contaminants (i.e., *Puccinia anthirrii* on seed of snapdragon or *Fusarium oxysporum* f. sp. *papaveris* on *Papaver nudicaule*) or may be internal (i.e., *F. oxysporum* f. sp. *matthiolae* on matthiola, *Alternaria zinniae* on zinnia, or *Heterosporium tropeoli* on nasturtium) (Baker 1972; Bertetti et al. 2015; Wu et al. 2006). Oospores of *Plasmopara obducens* have been observed in seeds of balsam impatiens; such seeds, when planted in disinfested soil, produced infected plants (Shishkoff 2019).

Seed can now be checked by using quick and reliable diagnostic tools and, if needed, treated with chemical, physical, or other means (Munkvold and Gullino 2020). Molecular diagnostic tools help with early pathogen identification, and work best when used by the seed industry to check the phytosanitary status of the material (Spadaro et al. 2020). For example, diagnostic tools were developed to rapidly detect *Phytophthora cryptogea* on naturally infected gerbera plants (Minerdi et al. 2008).

Still, at the beginning of this century, only a few centers had strong expertise in modern diagnostic methods (Crous 2005; Schaad et al. 2003). Increased international biosecurity and biodefense investments have helped to keep phytosanitary diagnostics abreast of new developments. The intensive development of new technologies provides new, more effective, quicker, and less labor-intensive methods for both validating the origin and tracing the movements of plant consignments.

There is a continued drive to provide rapid diagnostic methods for use at points of entry and inspection and to extension services for many important pathogens (Bonants 2014; J. E. Thomas et al. 2017; Verrier et al. 2017). Advances in molecular-based diagnostics should be very useful in the ornamentals industry. These new methods will have a strong impact on plant disease management (Munkvold and Gullino 2020).

Contaminated seeds do not need to be discarded but can be treated by the use of fungicides, heat, resistance inducers, antagonistic microorganisms, and plant extracts (Gullino et al. 2014). Due to regulatory constraints, at present there is a tendency to reduce the use of chemicals, replacing them with other means (Munkvold and Gullino 2020). However, when the pathogen is present inside the seed, only systemic fungicides and heat are effective for its eradication. In the case of bulb crops such as gladiolus and iris, hot-water treatments provided promising results for disease control (Gullino 2012).

Cultural practices. *Monitoring and control of environmental parameters.* In the case of greenhouse crops such as bedding plants, growers have the unique possibility to exert some control over the

Fig. 11. Rust on *Campanula rapunculoides* caused by *Coleosporium campanulae*.

Fig. 10. Powdery mildew caused by *Golovinomyces orontii* on *Campanula rapunculoides*.

Fig. 12. Round, white leaf spots on *Gaillardia aristata* caused by *Entyloma gaillardianum*.

environmental conditions in which crops are grown, both for optimizing plant growth and for managing pathogens. In the most sophisticated production facilities, disease management is greatly enhanced by monitoring and controlling temperature, light, humidity, water, ventilation, carbon dioxide, and crop nutrition with high precision. Manipulating the interactions of temperature and humidity is important in the control of foliar diseases, while rhizosphere moisture and temperature are relevant for root and stem diseases.

Dew deposition in the greenhouse, which is favorable for downy mildews, rusts, and *Botrytis* blight, is common on cool nights

following warm, humid days. Regulating day and night atmospheres is important for disease control and also helps in reducing the total amount of chemicals sprayed (Gullino and Garibaldi 2016; Hausbeck and Moorman 1996). *Botrytis cinerea* on fuchsia was managed with specific climate or ventilation conditions (Friedrich et al. 2005). For gray mold (*B. cinerea*) management, ventilation alone helps, minimizing the risk of epidemics by permitting good air movement.

Heating and ventilating at the same time or using a dehumidification device allows greenhouse growers to reduce relative humidity (Cámara-Zapata et al. 2019).

Fig. 13. Smart prevention measures for sustainable disease management.

Avoiding temperatures conducive to *Phytophthora infestans* and reducing moisture reduced late blight on petunia (Becktell et al. 2005). On rose, *Peronospora sparsa* is now much less important than in the past, due to the possibility of better control of greenhouse relative humidity, as well as the availability of effective chemicals (Gullino and Garibaldi 2016).

Root-zone or soil heating can be accomplished with either floor or bench heating systems. Altered greenhouse and bench design can improve air movement, thus reducing the risk of diseases. Through-the-bench air movement is, perhaps, the most neglected and simplest mean of reducing seedling rots in high-density systems.

Development of Fusarium root and crown rot incited by *F. hostae* in container-grown hostas is affected by the type of wounding that occurs during propagation and by container mix content, watering schedule, and temperature. Peat or peat-bark mixes reduced disease incidence and severity, and disease was higher on plants growing in container mix and at moderate (20 to 25°C) temperatures (Wang and Jeffers 2002). The application of lime to the medium increases the pH and suppresses Fusarium wilts (Elmer 2012). This has been observed in crops such as chrysanthemum (Engelhard and Woltz 1973) and gerbera (Gullino and Garibaldi 2016).

Shading is often used by growers when crops are wilting but the effect of this practice is mostly on temperature rather than on light intensity. In the case of root rot of lavender caused by *Phytophthora nicotianae* var. *parasitica* with a high level of rot incidence, a $\geq 50\%$ reduction in rot incidence was achieved by growing plants under shade. Maintaining plants under shade provided sufficient root rot control for practical cultivation conditions. The effect was attributed to reduced heating of the root zone in black pots used by growers; white containers might also have benefits. Shading strongly increases the severity of powdery mildew caused by *Golovinomyces cichoracearum* on aster (Gullino and Garibaldi 2016)—each host-pathogen system may have its own unique response to light.

Soilless cultivation and treatments. If bedding plants are grown with recirculating irrigation systems, the spread of diseases in soilless cultivation systems can be counteracted by adopting proper disinfection methods such as slow sand filtration for the recirculating solution. Closed recirculating soilless systems represent an interesting environment for exploiting innovative disease management options. Increasing the value of electrical conductivity of the nutrient solution and the use of potassium silicate amendments have proved effective against a number of foliar and soilborne diseases such as powdery mildews, downy mildews, leaf spots, and Fusarium wilts (Garibaldi et al. 2014b; Gullino et al. 2015b). When associated with a higher level of electrical conductivity, the deposition of amorphous silica in the cell wall could result as an increase in the production of lignin that could contribute to limiting the penetration of the pathogen within the plant cell (Gullino and Garibaldi 2016; Gullino et al. 2015b). The effects of electrical conductivity to induce resistance against *F. oxysporum* f. sp. *cyclaminis* on cyclamen have been documented by Elmer (2012) using chloride salts.

Soilless systems used for bedding plant production allow for a better manipulation of the microbiological conditions. This is achieved

through the application of microorganisms able to colonize the root system of plants grown under a strictly controlled environment. Suppressiveness of reused substrates adopted in soilless cultivation represents a very interesting feature, with many possible practical applications (Clematis et al. 2009). Antagonistic microorganisms isolated from soilless hydroponic systems have already been selected and proven effective against Fusarium wilt of rocket (Srinivasan et al. 2009) and *Pythium ultimum* on cucumber (Liu et al. 2009). Although utilized for field crops and vegetables, the widespread use of organic amendments for disease control is still not being largely practiced in bedding plant production, mainly because bedding plants are not usually produced in soil. In the future, organic amendments may have a role in garden plantings of bedding plants.

Resistance inducers are providing effective and long-term management of several soilborne pathogens on vegetable crops (Gilardi et al. 2019). Their ability to strengthen plant defense and contribute to reduced use of synthetic fungicides (Shoresh et al. 2010; Walters et al. 2013) is currently being investigated.

Resistance inducers are in continuous development. However, several aspects such as pathogen lifestyle, plant developmental stage, environmental and climatic conditions (temperature, relative humidity, and disease pressure), timing, and formulation and type of application, contribute to highly variable efficacy results (Walters and Fountaine 2009). A better understanding of the mode of action is needed; thus, the use of resistance inducers should be further investigated to improve efficacy in applications to bedding plants.

Substrate suppressiveness. Suppressiveness is a phenomenon very well described for soils. The phenomenon also occurs in the soilless substrates used in floriculture (Hadar 2011; Pugliese et al. 2015). It has been exploited for practical use against Fusarium wilts on carnation, cyclamen, and bulb crops, among others (Garibaldi and Gullino 1990; Gullino et al. 2015a).

In the past, much research has been carried out on peat, alone or mixed with other substrates, investigating its natural suppressiveness. Sphagnum peat mixes did naturally suppress diseases caused by soilborne pathogens but, within a few weeks after potting, they became conducive to diseases (Hoitink and Boehm 1999). Bacteria and fungi contributed to the suppression of root rots and wilts in peat mixes (Tahvonen and Kuuluvainen 1993). Because light peat decomposes in pots during production and the disease suppression effect is lost, amendments of decomposed peat mixes with microorganisms should consider the decomposition state (Hoitink and Locke 2012). Mixing light peat with more decomposed peat, at a 1:1 ratio, resulted in commercial sphagnum potting mixes capable of providing, through suppressiveness, control of soilborne pathogens for most greenhouse crops (Hoitink and Boehm 1999).

In the case of substrates, when hardwood bark (composted or not) is used, improved plant growth is generally observed, especially in potted plants. Suppressiveness and improved vigor of plants in such bark substrates result from the physical characteristics of bark composts and from higher levels of antagonists supported by these composts (Hoitink and Boehm 1999). Peat mixes also support well the introduction of biocontrol agents or the addition of composts

Table 2. Effect of municipal compost against *Phytophthora* spp. on ornamentals (Pugliese et al. 2012)^z

Substrate	Path	Compost (%)	<i>Phytophthora cinnamomi</i> on azalea		<i>P. nicotianae</i> on <i>Skimmia japonica</i>			
			Trial 1	Trial 2	Trial 1		Trial 2	
			Healthy (%)	Healthy (%)	Healthy (%)	Biomass (g)	Healthy (%)	Biomass (g)
Peat + Compost	Yes	10	92 a	56 bc	42 d	3.8 c	53 c	11.4 c
Peat + Compost	Yes	20	79 ab	10 d	58 bc	5.3 abc	87 bc	16.2 bc
Peat + Compost	Yes	40	89 ab	44 cd	75 abc	7.1 ab	90 ab	21.8 b
Peat + Compost	No	20	96 a	93 a	100 a	8.1 a	100 a	22 b
Peat	No	0	100 a	100 a	100 a	7.2 ab	100 a	29.2 a
Peat	Yes	0	69 b	6 d	45 d	3.4 c	50 c	9.5 c
Peat + Met	Yes	0	92 a	80 ab	89 ab	7.7 a	100 a	25.8 ab

^z Values followed by the same letter are not significantly different according to Tukey's honestly significant difference test ($P < 0.05$). Path = presence or absence of the pathogen, Compost = percentage of compost (vol/vol), Healthy = percentage of healthy plants, and Peat + Met = peat + metalaxyl-M (25 ml/m³).

(Hoitink and Locke 2012). Growing mixes fortified with *Trichoderma hamatum* 382 controlled Botrytis blight on geranium (Olson and Benson 2007) and begonia (Horst et al. 2005). The same biocontrol agents added to potting mixes controlled Fusarium wilt of cyclamen (Hoitink and Locke 2012).

During the past 25 years, recycling of composted organic wastes has been adopted for environmental, economic, and production reasons. From an environmental point of view, compost is considered an attractive peat substitute, after increasing concerns about the impact of peat extraction and the damage to peat land natural habitats by the horticulture industry (Silva et al. 2007). The cost of composts can be lower than that of peat. Composted materials can suppress soilborne pathogens (Noble and Coventry 2005; Pane et al. 2011; Pugliese et al. 2015; Termorshuizen et al. 2006). The consistency of disease control is improved when compost is enriched with selected microorganisms (Pugliese et al. 2011).

The suppressive capacity of compost against soilborne pathogens has been demonstrated in several studies leading to greater production efficiency and reduced nontarget effects (Garibaldi 1988; Hadar 2011; Hoitink and Boehm 1999; Noble 2011; Noble and Coventry 2005). Low rates of compost in growing media are generally indicated in order to avoid negative growth effects and phytotoxicity caused by high pH and electrical conductivity, and other phytotoxic compounds present in composts (Sullivan and Miller 2001). However, it is generally necessary to include at least 20% (vol/vol) of compost in containers in order to observe a suppressive effect (Table 2). Cases of increase of disease severity caused by composts used in containers have also been reported. A 50% spruce bark compost increased black root rot caused by *Thielaviopsis basicola* in poinsettias and Fusarium wilt of cyclamen, compared with a peat substrate (Krebs 1990). Highly saline composts were reported to enhance Pythium and Phytophthora diseases, while composts with higher nitrogen or ammonium content enhance Fusarium wilts (Hoitink et al. 2001). Success or failure of compost for disease control depends on the nature of the raw materials from which the compost was prepared, on the composting process used, and on the maturity and quality of the compost (Termorshuizen et al. 2006).

Control of soilborne diseases with organic amendments must be viewed as part of a systems approach where several aspects of the impact of crop production practices on resident soil microbial communities are addressed. New approaches to monitor how microbial community structures in soil change as a result of organic amendments may lead to a better understanding of which changes in microbial communities are responsible for conferring the disease suppressive effects (Cucu et al. 2019, 2020). This may eventually lead to improved, reliable disease controls for bedding plants resulting from organic amendment of growing media in greenhouse production.

Regulatory control. Quarantine restrictions and eradication measures are sometimes necessary in order to avoid the spread of pathogens that can severely affect the production of certain economically important crops (Ebbels 2003; Gamliel and Fletcher 2017).

State, regional, and international laws and regulations govern the production, sale, and transportation of ornamental plants. Measures designed to control the introduction of foreign pests are enforced in several countries. Domestic and international quarantines restrict the movement of specific plant materials at risk of carrying pests in order to prevent or delay their introduction. These quarantines often require cultural practices or chemical treatments to satisfy movement requirements (Guarnaccia et al. 2019b). Quarantine regulated pest lists are available in most regions (Ebbels 2003; Stebbins and Johnson 2001). Many pathogens are not detected because they are present as latent infections in plants that look healthy; missed detections in routine inspection are highly possible (Slippers and Wingfield 2007). Border controls are less effective than they should be because local quarantine lists of pathogens are often found only in inaccessible national databases or government publications; thus, the quarantines may not be supported by relevant data and nomenclature of the species in question may not be up to date (Crous et al. 2016).

Quarantines have been used, for instance, to limit movement of rust pathogens of geranium, chrysanthemum, daylily, and gladiolus

in several countries. Quarantines proved effective in the United States as well as in Australia in the case of chrysanthemum white rust, incited by *Puccinia horiana* (Bonde and Rizvi 1995), and gladiolus rust, incited by *Uromyces transversalis* (Beilharz et al. 2001). The same happened in Colombia where, in the early 1980s, a strict eradication and control campaign was implemented to keep all chrysanthemum exports free of *P. horiana* (Ortega 1999). If white rust was detected on plant material imported from Colombia, United States imports would be halted. In other cases, quarantine measures were ineffective. For instance, the containment of *P. hemerocallis*, the causal agent of daylily rust, in the United States was not possible due to the widespread movement of plants by hobbyists and nurseries (Williams-Woodward et al. 2001). Also, geranium rust, caused by *P. pelargonii-zonalis*, despite quarantine restrictions and the destruction of infected plants, became endemic in Europe and the United States (Wise et al. 2004).

Quarantine restriction and eradication efforts can be costly and have a significant economic impact on floriculture production and, as noted, are not always effective. However, the fact that ornamental plants can vector new pathogens, potentially causing losses on other crops, must be considered.

Research needs and future outlook. Many important problems of the bedding plant industry are unresolved and new ones are emerging as the industry undergoes more changes in production, marketing, and shipping procedures. Major changes will include more widely adopted mechanization and automation systems for improved crop management and the continuous introduction of new species, according to consumer taste. Some of those changes will affect the severity of diseases, thus challenging plant pathologists.

The phytopathology of bedding plants indeed includes a wide range of diseases on a variety of crops, with great opportunity for imaginative research and development of new methods for disease management, mostly based on preventive measures. The relatively low cost of the final product as compared with other floriculture crops is a strong constraint and forces growers to avoid too many external inputs.

The energy crisis, along with increasing restrictions on the use of pesticides, and the effect of climate change, along with the constant introduction of new crops from new areas around the world, pose new and continuous challenges to researchers and growers. In the future, interventions taking place at the production level, particularly at the seedling and plug production stage—those able to increase plant resilience—will be the most useful.

Plant pathologists will have to work more closely with horticulturists to ensure that new management practices have beneficial or, at worse, neutral effects on plant health.

The global movement of bedding plants is leading to alarming introductions of new plant pathogens into areas where they were previously not present. The detection of these organisms will lean on fast and accurate molecular techniques for quarantine and screening in the future (McTaggart et al. 2016). Methods to improve the accuracy and speed of field and laboratory diagnosis have been developed and need to be implemented more extensively (Spadaro et al. 2020; S. Thomas et al. 2017). The use of pathogen-free propagation material obtained through sanitation, clean growing media, pots, containers, or benches, disinfected mostly with steam, will continue to be important in the management of soilborne diseases. There is a demand for more effective disease control agents (biological microbials and natural materials). More research in the field of biological control is certainly needed in order to realize its potential. New approaches to monitor how microbial community structures in soil change as a result of amendments may lead to a better understanding of which changes in microbial communities are responsible for conferring the disease-suppressive effects of compost (Cucu et al. 2019). This may eventually lead to improved, more reliable disease control resulting from compost amendment of soil, sand, or peat, both in container crops in greenhouses and in the field (Noble 2011; Noble and Coventry 2005). Research on etiology, monitoring, modeling, and breeding for resistance is also needed.

Finally, there is a strong need for improved information flow among researchers, extension personnel, growers, and consumers; for

maintaining close links between the production and marketing sectors of the industry; and for giving more attention to consumers' needs. Due to the increased attention paid by consumers to environmental issues, efforts should be made to educate them to make wise choices. Unfortunately, few funding programs are designed specifically for ornamental pathology research and most agencies still consider ornamentals as a niche sector. This is very unfortunate, because such an attitude does not consider the fact that a number of techniques developed specifically for the ornamental industry (i.e., culture-indexing cuttings, apical meristem culture, improvements in tissue culture methods, virus indexing by grafting on indicator plants, soil steaming, and soilless cultivation) have also proved useful for controlling diseases on other crops (Baker and Linderman 1979; Gullino and Garibaldi 2007).

The very peculiar nature of the ornamental industry also requires very well-trained extension specialists. Unfortunately, during the past decade, most of the courses devoted to diseases of ornamentals, which were offered as part of the curriculum in many universities, have disappeared (Fletcher et al. 2020). The ornamental industry also needs highly qualified support by extension specialists, which is often missing.

The high economic and aesthetic value of bedding plants to a large portion of the population make them deserving of more attention by plant pathologists and, more generally, by researchers, while their beauty will continue attracting consumers. Thus, diseases of bedding plants and other ornamentals will continue to provide a fruitful and stimulating field of study for plant pathologists, and research in this field will hopefully no longer be considered as focused on the outer fringes of agriculture.

Acknowledgments

We thank A. Masino for drawing the figures, the anonymous reviewers and the Senior Editor for their precious contributions to improving the manuscript, and the many coworkers and students who, throughout the years, have collaborated in this field.

Literature Cited

- Aiello, D., Caiazzo, R., Carella, A., Lahoz, E., Nicoletti, R., Vitale, A., and Polizzi, G. 2009a. Characterisation of *Rhizoctonia* species isolated from ornamental nurseries. *J. Plant Pathol.* 91:45.
- Aiello, D., Castello, I., Vitale, A., Lahoz, E., Nicoletti, R., and Polizzi, G. 2008. First report of damping-off on African daisy caused by *Rhizoctonia solani* AG-4 in Italy. *Plant Dis.* 92:1367.
- Aiello, D., Guarnaccia, V., Formica, P. T., Hyakumachi, M., and Polizzi, G. 2017. Occurrence and characterisation of *Rhizoctonia* species causing diseases of ornamental plants in Italy. *Eur. J. Plant Pathol.* 148:967-982.
- Aiello, D., Hansen, Z., Smart, C., Polizzi, G., and Guarnaccia, V. 2018. Characterisation and mefenoxam sensitivity of *Phytophthora* spp. from ornamental plants in Italian nurseries. *Phytopathol. Mediterr.* 57:245-256.
- Aiello, D., Vitale, A., Lahoz, E., Nicoletti, R., and Polizzi, G. 2009b. First report of crown and root rot caused by *Rhizoctonia solani* AG-4 on orange jessamine in Italy. *Plant Dis.* 93:204.
- Amano, K. 1986. Host Range and Geographical Distribution of the Powdery Mildew Fungi. Japan Scientific Society Press, Tokyo, Japan.
- Armstrong, G. M., Armstrong, J. K., and Littrell, R. H. 1970. Wilt of chrysanthemum caused by *Fusarium oxysporum* f. sp. *chrysanthemi*, forma specialis nov. *Phytopathology* 60:496-498.
- Aveskamp, M. M., De Gruyter, J., Woudenberg, J. H. C., Verkley, G. J. M., and Crous, P. W. 2010. Highlights of the Didymellaceae: A polyphasic approach to characterise *Phoma* and related pleosporalean genera. *Stud. Mycol.* 65:1-60.
- Baker, K. F. 1972. Seed pathology. Pages 317-416 in: *Seed Biology*. T. T. Kozlowski, ed. Academic Press, New York, London.
- Baker, K. F., and Linderman, R. G. 1979. Unique features of the pathology of ornamental plants. *Annu. Rev. Phytopathol.* 17:253-277.
- Baysal-Gurel, F., Farinas, C., Peduto Hand, F., and Avin, F. 2020. First report of powdery mildew of *Phlox* caused by *Golovinomyces magnicellulatus* in Tennessee. *Plant Dis.* 104:2294.
- Becktell, M. C., Daughtrey, M. L., and Fry, W. E. 2005. Epidemiology and management of petunia and tomato late blight in the greenhouse. *Plant Dis.* 89:1000-1008.
- Beilharz, V., Parberry, D. G., and Pascoe, I. G. 2001. Gladiolus rust (caused by *Uromyces transversalis*) in eastern Australia. *Australas. Plant Pathol.* 30:267-270.
- Bélanger, R. R., Bushnell, W. R., Dik, A. J., and Carver, T. L. W. 2002. The Powdery Mildews: A Comprehensive Treatise. American Phytopathological Society, St. Paul, MN, U.S.A.
- Belbahri, L., Calmin, G., Pawlowski, J., and Lefort, F. 2005. Phylogenetic analysis and Real Time PCR detection of a presumably undescribed *Peronospora* species on sweet basil and sage. *Mycol. Res.* 109:1276-1287.
- Benson, D. M., and Cartwright, D. K. 1996. Ornamental diseases incited by *Rhizoctonia* spp. Pages 303-314 in: *Rhizoctonia* Species: Taxonomy, Molecular, Biology, Ecology, Pathology, and Disease Control. S. B. Sneh, S. Jabaji-Hare, G. Neate, and G. Dijst, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Bertetti, D., Ortu, G., Gullino, M. L., and Garibaldi, A. 2015. Contamination of seeds of Iceland poppy (*Papaver nudicaule* L.) by *Fusarium oxysporum*. *Phytoparasitica* 43:189-196.
- Bertetti, D., Ortu, G., Gullino, M. L., and Garibaldi, A. 2016. Contamination of moth mullein (*Verbascum blattaria* L.) seeds by *Phoma novae-verbascicola*. *J. Plant Pathol.* 98:361-363.
- Bertoldo, C., Gilardi, G., Spadaro, D., Gullino, M. L., and Garibaldi, A. 2015. Genetic diversity and virulence of Italian strains of *Fusarium oxysporum* isolated from *Eustoma grandiflorum*. *Eur. J. Plant Pathol.* 141:83-97.
- Bonanomi, G., Antignani, V., Pane, C., and Scala, F. 2007. Suppression of soilborne fungal diseases with organic amendments. *J. Plant Pathol.* 89:311-324.
- Bonants, P. J. M. 2014. Results of the EU Project QBOL, focusing on DNA barcoding of quarantine organisms, added to an International database (Q-Bank) on identification of quarantine pathogens and relatives. Pages 119-133 in: *Detection and Diagnostics of Plant Pathogens*. M. L. Gullino and P. J. M. Bonants, eds. Springer, Dordrecht, The Netherlands.
- Bonde, M. R., and Rizvi, S. A. 1995. Myclobutanil as a curative agent for chrysanthemum white rust. *Plant Dis.* 79:500-505.
- Booth, C. 1984. The *Fusarium* problem: Historical, economic and taxonomic aspects. Pages 1-13 in: *The Applied Mycology of Fusarium*. M. O. Moss and J. E. Smith, eds. Symp. Br. Mycol. Soc. Held at Queen Mary College, London, September 1982.
- Braun, U., and Cook, R. T. A. 2012. Taxonomy Manual of the Erysiphales (Powdery Mildews). Page 703 in: *CBS Biodiversity Series 11. Centraalbureau voor Schimmelcultures*, Utrecht, The Netherlands.
- Braun, U., Jage, H., Richter, U., and Zimmerman, H. 2009. *Peronospora verbenae* sp. nov.—A new downy mildew on *Verbena officinalis*. *Schlechtendalia* (Halle) 19:77-80.
- Byrne, J. M., Hausbeck, M. K., and Sconyers, L. E. 2005. Influence of environment on atmospheric concentrations of *Peronospora antirrhini* sporangia in field-grown snapdragon. *Plant Dis.* 89:1060-1066.
- Cámara-Zapata, J. M., Sánchez-Molina, J. A., Rodríguez, F., and López, J. C. 2019. Evaluation of a dehumidifier in a mild weather greenhouse. *Appl. Therm. Eng.* 146:92-103.
- Cannon, P. F., Damm, U., Johnston, P. R., and Weir, B. S. 2012. *Colletotrichum*—current status and future directions. *Stud. Mycol.* 73:181-213.
- Chase, A. R. 1991. Characterization of *Rhizoctonia* species isolated from ornamentals in Florida. *Plant Dis.* 75:234-238.
- Chase, A. R., Daughtrey, M. L., and Cloyd, R. A. 2018. *Compendium of Bedding Plant Diseases and Pests*. American Phytopathological Society, St. Paul, MN, U.S.A.
- Cho, S. E., Choi, I. Y., Hong, S. H., Lee, Y. H., and Shin, H. D. 2017. First report of powdery mildew caused by *Erysiphe begoniicola* on begonia in Korea. *Plant Dis.* 101:1681.
- Choi, Y.-J., Shin, H.-D., and Thines, M. 2009. Two novel *Peronospora* species are associated with recent reports of downy mildew on sages. *Mycol. Res.* 113:1340-1350.
- Clematis, F., Minuto, A., Gullino, M. L., and Garibaldi, A. 2009. Suppressiveness to *Fusarium oxysporum* f. sp. *radicis-lycopersici* in re-used perlite and perlite-peat substrates in soilless tomatoes. *Biol. Control* 48:108-114.
- Cline, E. T., Farr, D. F., and Rossman, A. Y. 2008. A synopsis of *Phytophthora* with accurate scientific names, host range, and geographic distribution. *Plant Health Prog.* 9 <https://apsjournals.apsnet.org/doi/10.1094/PHP-2008-0318-01-RV>.
- Couch, H. B. 1995. *Diseases of Turfgrasses*, 3rd ed. Krieger Publishing Company, Malabar, FL, U.S.A.
- Crous, P. W. 2005. Impact of molecular phylogenetics on the taxonomy and diagnostics of fungi. *Bull. OEPP* 35:47-51.
- Crous, P. W., Gams, W., Stalpers, J. A., Robert, V., and Stegehuis, G. 2004. MycoBank: An online initiative to launch mycology into the 21st century. *Stud. Mycol.* 50:19-22.
- Crous, P. W., Groenewald, J. Z., Slippers, B., and Wingfield, M. J. 2016. Global food and fibre security threatened by current inefficiencies in fungal identification. *Philos. Trans. R. Soc. B* 371:20160024.
- Crous, P. W., Phillips, A. J. L., and Baxter, A. P. 2000. *Phytopathogenic Fungi from South Africa*. University of Stellenbosch, Department of Plant Pathology Press, Stellenbosch, South Africa.
- Cucu, M. A., Gilardi, G., Pugliese, M., Gullino, M. L., and Garibaldi, A. 2020. An assessment of the modulation of the population dynamics of pathogenic *Fusarium oxysporum* f. sp. *lycopersici* in the tomato rhizosphere by means of the application of *Bacillus subtilis* QST 713, *Trichoderma* sp. TW2 and two composts. *Biol. Control* 142:104158.

Cucu, M. A., Gilardi, G., Pugliese, M., Matic, S., Gisi, U., Gullino, M. L., and Garibaldi, A. 2019. Influence of different biological control agents and compost on total and nitrification-driven microbial communities at rhizosphere and soil level in a lettuce-*Fusarium oxysporum* f. sp. *lactucae* pathosystem. *J. Appl. Microbiol.* 126:905-918.

Daughtrey, M., and Buitenhuis, R. 2020. Integrated pest and disease management in greenhouse ornamentals. Pages 625-679 in: *Integrated Pest*

and Disease Management in Greenhouse Crops. M. L. Gullino, R. Albajes, and P. Nicot, eds. Springer, Dordrecht, The Netherlands.

Daughtrey, M. L., and Benson, D. M. 2005. Principles of plant health management for ornamental plants. *Annu. Rev. Phytopathol.* 43:141-169.

Daughtrey, M. L., Holcomb, G. E., Eshenaur, B., Palm, M. E., Belbahri, L., and Lefort, F. 2006. First report of downy mildew on greenhouse

Dr. Vladimiro Guarnaccia has been Assistant Professor at the Department of Agricultural, Forest, and Food Sciences at the University of Torino since 2019, actively collaborating with the Centre of Competence Agroinnova. In 2013, he obtained his Ph.D. in Plant Health Technologies and Protection of Agro-ecosystems at University of Catania, Italy, working on fungal pathogens of ornamental, subtropical, and tropical plants. As a researcher at Westerdijk Institute (formerly CBS) in Utrecht, The Netherlands (2015–2018), he explored the fungal diversity of wood diseases of citrus and other fruit crops. He also worked at the University of Stellenbosch, South Africa (2018) on the characterization and biological control of fungi causing root disease of citrus trees. His current research at the University of Torino includes berry fruit, apple, and other economically important fruit crops. He is currently working on disease etiology, epidemiology, and diagnostics of these crops. He has authored or coauthored over 70 peer-reviewed articles in scientific journals as well as technical extension articles. He is member of the editorial boards of the journals *Phytopathologia Mediterranea*, *FUSE*, and *CABI Agriculture and Bioscience*.

Dr. Francesca Peduto Hand is an associate professor and extension specialist in the Department of Plant Pathology at The Ohio State University. Previously, she received an M.S. degree in Agricultural Science and Technology (2003) and a Ph.D. in Plant Pathology (2008), both from the University of Florence, Italy, and conducted postdoctoral research at the University of California Davis (2009–2012). Her background and interests lay in the fundamental and applied aspects of pathogen biology, disease epidemiology, diagnostics, and integrated disease management. She has authored or coauthored over 40 peer-reviewed articles and multiple book chapters. She has an active research and extension program focused on management of diseases of ornamental plants in greenhouse and nursery production. In her work, she uses a combination of traditional, metabolomic, and genomic approaches to study recurrent and emergent infectious diseases that are limiting the profitability of Ohio's ornamental industry, with the ultimate goal of modifying grower production practices based on improved knowledge of pathogen biology, epidemiology, and genomics. She is a strong advocate for women leadership in STEM.

Angelo Garibaldi is an Emeritus Professor and President of the Centre of Competence Agroinnova at the University of Torino. He is still very much involved in research in the field of plant disease management, diseases of ornamental crops, and biology and epidemiology of emerging diseases on horticultural crops. In particular, he has been and is very much involved in studying the biology and epidemiology of diseases affecting ornamental plants, with much attention to the new trends of this sector. On the occasion of his eightieth birthday, a cartoon book, *Angelo, the Flower's Doctor*, was dedicated to him. He has been President of the Associazione Fitopatologica Italiana (1986–1990), Dean of the Faculty of Agriculture (1990–1996), and Vice-Chancellor of the University of Torino (1996–2004). He managed many international and national research projects funded by the European Union and different ministries and agencies, and has always been strongly involved in technology transfer. He has mentored hundreds of students during his long academic career and is the author or coauthor of eight books and over 1,000 scientific articles. He is a founder and currently president of AgriNewTech S.r.l., an academic spin-off company.

Maria Lodovica Gullino is a Professor in Plant Pathology, Director of the Center of Competence Agroinnova, and Vice Rector for the Valorization of Human and Cultural Resources at the University of Torino. She has been President of the Italian Association for Crop Protection (2003–2006), Italian Association of the Agrarian Societies (2004–2006), International Society for Plant Pathology (2008–2013), and Italian Society for Plant Pathology (2017–2019). Her research interests cover plant disease management, food safety and security, plant and food biosecurity, and effects of climate change on plant diseases. She is coauthor of 25 books and over 900 scientific papers. She is a Fellow of the American Phytopathological Society and of the International Society for Plant Pathology. She has been and is responsible for many research projects from the European Union, Ministry of University and Research, Ministry of Agriculture, Ministry of Environment, regional governments, and private industry, and is the founder of AgriNewTech S.r.l., an academic spin-off company. As a professional journalist and writer of popular books aimed at attracting people at large and children to science, she is very much involved in communication and public engagement activities.

- and landscape *Coleus* caused by a *Peronospora* sp. in Louisiana and New York. Plant Dis. 90:1111.
- Dean, R., Van Kan, J. A. L., Pretorius, Z. A., Hammond-Kosack, K. E., Di Pietro, A., Spanu, P. D., Rudd, J. J., Dickman, M., Kahmann, R., Ellis, J., and Foster, G. D. 2012. The top 10 fungal pathogens in molecular plant pathology. Mol. Plant Pathol. 13:414-430.
- de Gruyter, J., Aveskamp, M. M., Woudenberg, J. H. C., Verkley, G. J. M., Groenewald, J. Z., and Crous, P. W. 2009. Molecular phylogeny of *Phoma* and allied anamorph genera: Towards a reclassification of the *Phoma* complex. Mycol. Res. 113:508-519.
- Donahoo, R. S., and Lamour, K. H. 2008. Characterization of *Phytophthora* species from leaves of nursery woody ornamentals in Tennessee. HortScience 43:1833-1837.
- Ebbels, D. L. 2003. Principles of Plant Health and Quarantine. CABI Publishing, Wallingford, U.K. doi.org/10.1079/9780851996806.0000
- Elmer, W. H. 2002. Influence of inoculum density of *Fusarium oxysporum* f. sp. *cyclaminis* and sodium chloride on cyclamen and the development of Fusarium wilt. Plant Dis. 86:389-393.
- Elmer, W. H. 2008. Preventing the spread of Fusarium wilt of *Hiemalis* begonias in the greenhouse. Crop Prot. 27:1078-1083.
- Elmer, W. H. 2012. Cultural practices. Pages 67-74 in: Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Elmer, W. H., and Daughtrey, M. 2012. Fusarium wilt of cyclamen. Pages 213-219 in: Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Engelhard, A. W., and Woltz, S. S. 1973. Fusarium wilt of chrysanthemum: Complete control of symptoms with an integrated fungicide-lime-nitrate regime. Phytopathology 63:1256-1259.
- Erwin, D. C., and Ribeiro, O. K. 1996. Phytophthora Diseases Worldwide. American Phytopathological Society, St. Paul, MN, U.S.A.
- European Commission. 2020. Flowers and ornamental plants – Production. Statistics 2010 – 2019. https://ec.europa.eu/info/sites/info/files/food-farming-fisheries/plants_and_plant_products/documents/flowers-ornamental-plants-statistics_en.pdf
- Farinas, C., Jourdan, P., Paul, P. A., and Peduto Hand, F. 2019. Development and evaluation of laboratory bioassays to study powdery mildew pathogens of *Phlox in vitro*. Plant Dis. 103:1536-1543.
- Farinas, C., Jourdan, P., Paul, P. A., Slot, J. C., Daughtrey, M. L., Devi Ganesan, V., Baysal-Gurel, F., and Peduto Hand, F. 2020. *Phlox* species show quantitative and qualitative resistance to a population of powdery mildew isolates from the Eastern United States. Phytopathology 110:1410-1418
- Farr, D. F., and Rossman, A. Y. 2020. Fungal Databases U.S. National Fungus Collections. United States Department of Agriculture–Agricultural Research Service <https://nt.ars-grin.gov/fungal-databases/>
- Faust, J. E., Dole, J. M., and Lopez, R. G. 2017. The floriculture vegetative cutting industry. Pages 121-172 in: Horticultural Reviews, Vol. 44, 1st ed. J. Janick, ed. Wiley-Blackwell, Oxford, U.K.
- Ferguson, A. J., and Jeffers, S. N. 1999. Detecting multiple species of *Phytophthora* in container mixes from ornamental crop nurseries. Plant Dis. 83:1129-1136.
- Ferrada, E. E., Böhm, L., Montenegro, O., Llancao, M., Romero, C., Briceno, E. X., Morales, R. A., and Díaz, G. A. 2020. First report of leaf rust of *Fuchsia magellanica* caused by *Pucciniastrum circaeae* in Valdivia, Chile. Plant Dis. 104:1548.
- Fletcher, J., Gamliel, A., Gullino, M. L., McKirdy, S. J., Smith, G. R., and Stack, J. P. 2020. A fresh look at graduate education in plant pathology in a changing world: Global needs and perspectives. J. Plant Pathol. 102:609-618.
- Fletcher, J., and Martin, J. A. 1972. Spread and control of Fusarium wilt of carnation. Plant Pathol. 21:182-187.
- Friedrich, S., Gebelein, D., and Boyle, C. 2005. Control of *Botrytis cinerea* in glasshouse fuchsia by specific climate management. Eur. J. Plant Pathol. 111:249-262.
- Gamliel, A., and Fletcher, J. 2017. Containment and eradication of invasive pathogens. Pages 243-256 in: Practical Tools for Plant and Food Biosecurity. M. L. Gullino, J. P. Stack, J. Fletcher, and J. D. Mumford, eds. Springer, Dordrecht, The Netherlands.
- Garibaldi, A. 1988. Research on substrates suppressive to *Fusarium oxysporum* and *Rhizoctonia solani*. Acta Hort. 221:271-278.
- Garibaldi, A., Bertetti, D., Amatulli, M. T., and Gullino, M. L. 2010a. First report of leaf spot of orange coneflower (*Rudbeckia fulgida*) caused by a *Phoma* sp. in Italy. Plant Dis. 94:788.
- Garibaldi, A., Bertetti, D., Frati, S., and Gullino, M. L. 2008a. First report of powdery mildew caused by *Golovinomyces cichoracearum* on orange coneflower (*Rudbeckia fulgida*) in Italy. Plant Dis. 92:975.
- Garibaldi, A., Bertetti, D., Frati, S., Minuto, A., and Gullino, M. L. 2008b. Powdery mildew caused by *Golovinomyces cichoracearum* on Paris daisy (*Argyranthemum frutescens*) in Italy. Plant Dis. 92:1135.
- Garibaldi, A., Bertetti, D., and Gullino, M. L. 2006. First report of leaf blight caused by *Rhizoctonia solani* AG 1B on Madagascar periwinkle (*Catharanthus roseus*) in Italy. Plant Dis. 90:1361.
- Garibaldi, A., Bertetti, D., and Gullino, M. L. 2007. First report of powdery mildew caused by *Golovinomyces orontii* (*Erysiphe orontii*) on *Petunia hybrid* in Italy. Plant Dis. 91:632.
- Garibaldi, A., Bertetti, D., and Gullino, M. L. 2009a. First report of leaf blight on *Hosta fortunei* caused by *Rhizoctonia solani* AG 4 in Italy. Plant Dis. 93:432.
- Garibaldi, A., Bertetti, D., and Gullino, M. L. 2009b. Outbreak of powdery mildew caused by *Erysiphe cruciferarum* on spider flower (*Cleome hassleriana*) in Italy. Plant Dis. 93:963.
- Garibaldi, A., Bertetti, D., Matic, S., and Gullino, M. L. 2017a. A leaf spot caused by *Stagonosporopsis trachelii* on *Campanula trachelium* in Italy. J. Plant Pathol. 99:540.
- Garibaldi, A., Bertetti, D., Matic, S., and Gullino, M. L. 2018a. First report of downy mildew caused by *Peronospora arthurii* on common evening primrose (*Oenothera biennis*) in Italy. Plant Dis. 102:688.
- Garibaldi, A., Bertetti, D., Matic, S., and Gullino, M. L. 2018b. First report of leaf spot of *Salvia elegans* caused by *Alternaria alternata* in Italy. Plant Dis. 102:1034.
- Garibaldi, A., Bertetti, D., Matic, S., and Gullino, M. L. 2018c. First report of powdery mildew caused by *Golovinomyces orontii* on *Campanula glomerata* in Italy. Plant Dis. 102:2041.
- Garibaldi, A., Bertetti, D., Matic, S., and Gullino, M. L. 2018d. Powdery mildew caused by *Golovinomyces cichoracearum* on eastern purple coneflower (*Echinacea purpurea*) in Italy. Plant Dis. 102:1028.
- Garibaldi, A., Bertetti, D., Matic, S., and Gullino, M. L. 2019a. First report of powdery mildew caused by *Golovinomyces orontii* on *Abelmoschus manihot* in Italy. Plant Dis. 103:154.
- Garibaldi, A., Bertetti, D., Matic, S., Luongo, I., and Gullino, M. L. 2019b. First report of *Alternaria alternata* on *Campanula rapunculoides* in Italy. Plant Dis. 103:368.
- Garibaldi, A., Bertetti, D., Matic, S., Luongo, I., and Gullino, M. L. 2019c. First report of leaf necrosis of *Salvia dorisiana* caused by *Alternaria alternata* in Italy. Plant Dis. 103:1025.
- Garibaldi, A., Bertetti, D., Matic, S., Luongo, I., and Gullino, M. L. 2020a. First report of leaf necrosis caused by *Alternaria alternata* on *Plectranthus scutellarioides* in Italy. Plant Dis. 104:590.
- Garibaldi, A., Bertetti, D., Matic, S., Luongo, I., and Gullino, M. L. 2020b. First report of leaf spot caused by *Alternaria alternata* on *Phlox maculata* in Italy. Plant Dis. 104:2519.
- Garibaldi, A., Bertetti, D., Matic, S., Luongo, I., and Gullino, M. L. 2020c. First report of leaf spots caused by *Alternaria arborescens* on *Symphyotrichum novi-belgii* in Italy. Plant Dis. 104:281.
- Garibaldi, A., Bertetti, D., Ortega, S. F., and Gullino, M. L. 2016a. Powdery mildew caused by *Golovinomyces magnicellulatus* on *Phlox paniculata* in Italy. J. Plant Pathol. 98:176.
- Garibaldi, A., Bertetti, D., Ortu, G., and Gullino, M. L. 2013a. First report of web blight on *Nigella damascena* caused by *Rhizoctonia solani* AG 1-B in Italy. J. Plant Pathol. 95:S4.75.
- Garibaldi, A., Bertetti, D., Ortu, G., and Gullino, M. L. 2014a. First report of *Phoma nova-verbascicola* on moth mullein (*Verbascum blattaria*) in Italy. J. Plant Pathol. 96:436.
- Garibaldi, A., Bertetti, D., Ortu, G., and Gullino, M. L. 2015a. First report of web blight on rose campion (*Lychnis coronaria*) caused by *Rhizoctonia solani* AG 1-IB in Italy. Plant Dis. 99:162.
- Garibaldi, A., Bertetti, D., Ortu, G., and Gullino, M. L. 2015b. A leaf spot caused by *Stagonosporopsis trachelii* on *Campanula medium* in Italy. J. Plant Pathol. 97:546.
- Garibaldi, A., Bertetti, D., Pensa, P., and Gullino, M. L. 2009c. First report of collar and stem rot caused by *Pythium aphanidermatum* on fig-marigold (*Lampranthus* sp.) in Italy. Plant Dis. 93:672.
- Garibaldi, A., Bertetti, D., Pensa, P., Ortu, G., and Gullino, M. L. 2015c. *Phytophthora cryptogea* on common sage (*Salvia officinalis* L.) in Italy. Plant Dis. 99:161.
- Garibaldi, A., Bertetti, D., Poli, A., and Gullino, M. L. 2011a. Severe outbreaks of powdery mildew caused by *Podosphaera* sp. on *Phlox drummondii* in Italy. J. Plant Pathol. 93:S4.84.
- Garibaldi, A., Bertetti, D., Poli, A., and Gullino, M. L. 2012a. Powdery mildew caused by *Golovinomyces orontii* on creeping bellflower (*Campanula rapunculoides*) in Italy. Plant Dis. 96:291.
- Garibaldi, A., Bertetti, D., Poli, A., and Gullino, M. L. 2013b. Outbreak of downy mildew caused by *Peronospora digitalidis* on common foxglove (*Digitalis purpurea*) in Italy. J. Plant Pathol. 95:660.
- Garibaldi, A., Bertetti, D., Poli, A., and Gullino, M. L. 2013c. First report of leaf spot of *Saponaria officinalis* caused by *Alternaria nobilis* in Italy. Plant Dis. 97:424.
- Garibaldi, A., Bertetti, D., Poli, A., and Gullino, M. L. 2013d. A leaf spot caused by *Phoma novae-verbascicola* on black mullein (*Verbascum nigrum* L.) in Italy. Plant Dis. 97:1660.
- Garibaldi, A., Bertetti, D., Rapetti, S., and Gullino, M. L. 2011b. First report of a new *Fusarium oxysporum* causing carnation wilt in Colombia. J. Plant Pathol. 93:S4.74.

- Garibaldi, A., Gilardi, G., Bertetti, D., and Gullino, M. L. 2009d. First report of leaf blight on Fan columbine (*Aquilegia flabellata*) caused by *Rhizoctonia solani* AG 4 in Italy. Plant Dis. 93:433.
- Garibaldi, A., Gilardi, G., Bertetti, D., and Gullino, M. L. 2009e. Report of leaf blight on Washington Lupine (*Lupinus polyphyllus*) caused by *Rhizoctonia solani* AG 4 in Italy. Plant Dis. 93:429.
- Garibaldi, A., Gilardi, G., Bertetti, D., and Gullino, M. L. 2009f. First report of leaf blight on foxglove (*Digitalis purpurea*) caused by *Rhizoctonia solani* AG-1-IA in Italy. Plant Dis. 93:318.
- Garibaldi, A., Gilardi, G., Bertetti, D., and Gullino, M. L. 2010b. First report of leaf blight on woodland sage caused by *Rhizoctonia solani* AG 1 in Italy. Plant Dis. 94:1071.
- Garibaldi, A., Gilardi, G., Franco-Ortega, S., and Gullino, M. L. 2016b. First report of a leaf spot caused by *Boeremia exigua* var. *linicola* on autumn sage (*Salvia greggii*) in Italy and worldwide. Plant Dis. 100:1777.
- Garibaldi, A., Gilardi, G., Franco-Ortega, S., and Gullino, M. L. 2016c. First report of leaf spot caused by *Colletotrichum fiorinae* on Mexican bush sage (*Salvia leucantha*) in Italy. Plant Dis. 100:654.
- Garibaldi, A., Gilardi, G., and Gullino, M. L. 2005. Fusarium wilt of bit-terroot (*Lewisia cotyledon*) in Italy caused by *Fusarium oxysporum*. Plant Dis. 89:684.
- Garibaldi, A., Gilardi, G., and Gullino, M. L. 2014b. Critical aspects in disease management as a consequence of the evolution of soil-borne pathogens. Acta Hortic. 1044:43-50.
- Garibaldi, A., Gilardi, G., Matic, S., and Gullino, M. L. 2017b. First report of Fusarium wilt on orange coneflower (*Rudbeckia fulgida*) in northern Italy. Plant Dis. 101:1546.
- Garibaldi, A., Gilardi, G., Matic, S., and Gullino, M. L. 2017c. First report of rust caused by *Coleosporium campanulae* on *Campanula rapunculoides* in Italy. J. Plant Pathol. 99:287.
- Garibaldi, A., Gilardi, G., Matic, S., and Gullino, M. L. 2018e. First report of leaf spot of peppermint (*Mentha × piperita*) caused by *Alternaria alternata* in Italy. Plant Dis. 102:1041.
- Garibaldi, A., Gilardi, G., Matic, S., and Gullino, M. L. 2018f. First report of leaf spot caused by *Alternaria alternata* on *Echinacea purpurea* in Italy. Plant Dis. 102:1450.
- Garibaldi, A., Gilardi, G., Matic, S., and Gullino, M. L. 2018g. First report of leaf smut caused by *Entyloma gaillardianum* on *Gaillardia aristata* in Italy. Plant Dis. 102:678.
- Garibaldi, A., Gilardi, G., Matic, S., and Gullino, M. L. 2019d. First report of *Alternaria alternata* causing leaf spot on *Digitalis purpurea* in Italy. Plant Dis. 103:1770.
- Garibaldi, A., Gilardi, G., Matic, S., Luongo, I., and Gullino, M. L. 2019e. First report of leaf necrosis caused by *Alternaria alternata* on *Ceratostigma willmottianum* in Italy. Plant Dis. 103:1412.
- Garibaldi, A., Gilardi, G., Matic, S., Luongo, I., and Gullino, M. L. 2020d. First report of leaf spot of *Alcea rosea* caused by *Alternaria alternata* in Italy. Plant Dis. 104:588.
- Garibaldi, A., Gilardi, G., Ortu, G., and Gullino, M. L. 2012b. First report of rust caused by *Pucciniastrum circaeae* on *Fuchsia × hybrida* in Italy. Plant Dis. 96:588.
- Garibaldi, A., and Gullino, M. L. 1985. Wilt of *Ranunculus asiaticus* caused by *Fusarium oxysporum* f. sp. *ranunculi*, forma specialis nova. Phytopathol. Mediterr. 24:213-214.
- Garibaldi, A., and Gullino, M. L. 1990. Disease management of ornamental plants: A never ending challenge. Meded. Fac. Landbouwwet. Rijksuniv. Gent 55:189-201.
- Garibaldi, A., and Gullino, M. L. 2010. Emerging soilborne diseases of horticultural crops and new trends in their management. Acta Hortic. 883:37-47.
- Garibaldi, A., and Gullino, M. L. 2012a. Fusarium wilt of some ornamental compositae. Pages 205-211 in: Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Garibaldi, A., and Gullino, M. L. 2012b. Fusarium wilt of carnation. Pages 191-198 in: Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Garibaldi, A., Minuto, A., Bertetti, D., and Gullino, M. L. 2004. Fusarium wilt of gerbera in soil and soilless crops in Italy. Plant Dis. 88:311.
- Garibaldi, A., Minuto, A., Bertetti, D., Nicoletti, R., and Gullino, M. L. 2003a. First report of web blight on yellow-sage (*Lantana camara*) caused by *Rhizoctonia solani* in Europe. Plant Dis. 87:875.
- Garibaldi, A., Minuto, A., Grasso, V., and Gullino, M. L. 2003b. Application of selected antagonistic strains against *Phytophthora cryptogea* on gerbera in closed soilless systems with disinfection by slow sand filtration. Crop Prot. 22:1053-1061.
- Garibaldi, A., Minuto, A., Uchimura, M. S., and Gullino, M. L. 2008c. Fusarium wilt of gerbera caused by a *Fusarium* sp. in Brazil. Plant Dis. 92:655.
- Garibaldi, A., Ortu, G., Bertetti, D., and Gullino, M. L. 2015d. First report of web blight on Creeping Bellflower (*Campanula rapunculoides*) caused by *Rhizoctonia solani* AG 1-IB in Italy. Plant Dis. 99:1037.
- Garibaldi, A., Tabone, G., Bertetti, D., and Gullino, M. L. 2020e. First report of *Rhizoctonia solani* AG-4 HG-I causing crown rot on *Abelmoschus manihot* in Italy. J. Plant Pathol. 102:1303.
- Garibaldi, A., Tabone, G., Matic, S., Luongo, I., and Gullino, M. L. 2020f. First report of *Alternaria alternata* causing leaf spot on *Salvia involucrata* in Italy. J. Plant Pathol. 102:1279.
- Garzón, C. D., Molineros, J. E., Yáñez, J. M., Flores, F. J., Jiménez-Gasco, M. M., and Moorman, G. W. 2011. Sublethal doses of mefenoxam enhance Pythium damping-off of geranium. Plant Dis. 95:1233-1238.
- Gerlach, W. W. P., and Schubert, R. 2001. A new wilt of *Cyclamen* caused by *Phytophthora tropicalis* in Germany and the Netherlands. Plant Dis. 85:334.
- Gilardi, G., Matic, S., Gullino, M. L., and Garibaldi, A. 2017. First report of *Phoma herbarum* causing leaf spot of woodland sage (*Salvia nemorosa*) in northern Italy. Plant Dis. 101:1824.
- Gilardi, G., Pugliese, M., Gullino, M. L., and Garibaldi, A. 2019. Nursery treatments with resistant inducers, soil amendments and biocontrol agents for the management of the Fusarium wilt of lettuce under glasshouse and field conditions. J. Phytopathol. 167:98-110.
- Glawe, D. A. 2008. The powdery mildews: A review of the world's most familiar (yet poorly known) plant pathogens. Annu. Rev. Phytopathol. 46:27-51.
- Glawe, D. A., Barlow, T., and Koike, S. T. 2010. First report of leaf smut of *Gaillardia × grandiflora* caused by *Entyloma gaillardianum* in North America. Plant Health Prog. 11. <https://apsjournals.apsnet.org/doi/10.1094/PHP-2010-0428-01-BR>
- Guarnaccia, V., Aiello, D., Polizzi, G., Crous, P. W., and Sandoval-Denis, M. 2019a. Soilborne diseases caused by *Fusarium* and *Neocosmospora* spp. on ornamental plants in Italy. Phytopathol. Mediterr. 58:127-137.
- Guarnaccia, V., Gehrmann, T., Silva-Junior, G. J., Fourie, P. H., Haridas, S., Vu, D., Spatafora, J., Martin, F. M., Robert, V., Grigoriev, I. V., Groenewald, J. Z., and Crous, P. W. 2019b. *Phyllosticta citricarpa* and sister species of global importance to *Citrus*. Mol. Plant Pathol. 20:1619-1635.
- Guarnaccia, V., Gilardi, G., Martino, I., Garibaldi, A., and Gullino, M. L. 2019c. Species diversity in *Colletotrichum* causing anthracnose of aromatic and ornamental *Lamiaceae* in Italy. Agronomy (Basel) 9:613.
- Guarnaccia, V., Hansen, Z. R., Aiello, D., Smart, C. D., and Polizzi, G. 2015. First detection of root rot and foliar blight on pittosporum (*Pittosporum tenuifolium*) caused by *Pythium irregulare* in Italy. J. Phytopathol. 163:411-414.
- Guarnaccia, V., Martino, I., Gilardi, G., Garibaldi, A., and Gullino, M. L. 2019d. *Colletotrichum* spp. causing anthracnose on ornamental plants in Northern Italy. J. Plant Pathol. In press. doi.org/10.1007/s42161-020-00684-2
- Guarnaccia, V., Sandoval-Denis, M., Aiello, D., Polizzi, G., and Crous, P. W. 2018. *Neocosmospora perseae* sp. nov., causing trunk cankers on avocado in Italy. Fungal Syst. Evol. 1:131-140.
- Gullino, M. L. 2012. Fusarium wilts of bulb crops. Pages 199-204 in: Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Gullino, M. L., Daughtrey, M. L., Garibaldi, A., and Elmer, W. H. 2015a. Fusarium wilts of ornamental crops and their management. Crop Prot. 73:50-59.
- Gullino, M. L., and Garibaldi, A. 2006. Evolution of fungal diseases of ornamental plants and main implications for their management. Pages 464-471 in: Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues. J. A. Teixeira da Silva, ed. Global Science Books, London, U.K.
- Gullino, M. L., and Garibaldi, A. 2007. Critical aspects in management of fungal diseases of ornamental plants and directions in research. Phytopathol. Mediterr. 46:133-149.
- Gullino, M. L., and Garibaldi, A. 2016. Environmental modification for disease management. Pages 1-18 in: Handbook of Florists' Crops Diseases. Handbook of Plant Disease Management. R. J. McGovern and W. H. Elmer, eds. Springer International Publishing, Cham, Switzerland.
- Gullino, M. L., Gilardi, G., and Garibaldi, A. 2014. Seed-borne fungal pathogens of leafy vegetable crops. Pages 47-56 in: Global Perspectives on the Health of Seeds and Plant Propagation Material. M. L. Gullino and G. Munkvold, eds. Springer, Dordrecht, The Netherlands.
- Gullino, M. L., Pugliese, M., and Garibaldi, A. 2015b. Use of silicon amendments against foliar and vascular diseases of vegetables grown soilless. Pages 293-306 in: Sustainable Crop Disease Management using Natural Products. G. Sangeetha, V. Kurucheva, and J. Jayaraj, eds. CABI, Oxfordshire, U.K.
- Hadar, Y. 2011. Suppressive compost: When plant pathology met microbiological ecology. Phytoparasitica 39:311-314.
- Hao, W., Richardson, P. A., and Hong, C. X. 2010. Foliar blight of annual Vinca (*Catharanthus roseus*) caused by *Phytophthora tropicalis* in Virginia. Plant Dis. 94:274.
- Hausbeck, M. K., and Moorman, G. W. 1996. Managing *Botrytis* in greenhouse-grown flower crops. Plant Dis. 80:1212-1219.

- Hernández, J. R., and Hennen, J. F. 2003. Rust fungi causing galls, witches' brooms, and other abnormal plant growths in northwestern Argentina. *Mycologia* 95:728-755.
- Hoitink, H. A. J., and Boehm, M. J. 1999. Biocontrol within the context of soil microbial communities: A soil-dependent phenomenon. *Annu. Rev. Phytopathol.* 37:427-446.
- Hoitink, H. A. J., Krause, M. S., and Han, Y. 2001. Spectrum and mechanisms of plant disease control with composts. Pages 263-274 in: *Compost Utilization in Horticultural Cropping Systems*. H. Hoitink and H. Keener, eds. CRC Press LLC, Boca Raton, FL, U.S.A.
- Hoitink, H. A. J., and Locke, J. C. 2012. An integrated approach to biological control of *Fusarium* species in containerized crops. Pages 109-115 in: *Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops*. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Holcomb, G. E., and Carling, D. E. 2000. First report of web blight of verbena caused by *Rhizoctonia solani*. *Plant Dis.* 84:492.
- Hong, C., Richardson, P. A., and Kong, P. 2008. Pathogenicity to ornamental plants of some existing species and new taxa of *Phytophthora* from irrigation water. *Plant Dis.* 92:1201-1207.
- Hong, C. X., and Banko, T. J. 2003. First report on white smut of *Gaillardia* × *grandiflora* caused by *Entyloma polysporum* in Virginia. *Plant Dis.* 87:313.
- Hong, C. X., and Moorman, G. W. 2005. Plant pathogens in irrigation water: Challenges and opportunities. *CRC Crit. Rev. Plant Sci.* 24:189-208.
- Hong, C. X., Richardson, P. A., Kong, P., Jeffers, S. N., and Oak, S. W. 2006. *Phytophthora tropicalis* isolated from diseased leaves of *Pieris japonica* and *Rhododendron catawbiense* and found in irrigation water and soil in Virginia. *Plant Dis.* 90:525.
- Horst, L. E., Locke, J., Krause, C. R., McMahon, R. W., Madden, L. V., and Hoitink, H. A. J. 2005. Suppression of Botrytis blight of begonia by *Trichoderma hamatum* 382 in peat and compost amended potting mixes. *Plant Dis.* 89:1195-1200.
- Horst, R. K., and Nelson, P. E. 1997. Compendium of Chrysanthemum Diseases. American Phytopathological Society, St. Paul, MN, U.S.A.
- Hou, L., Hernández-Restrepo, M., Groenewald, J. Z., Cai, L., and Crous, P. W. 2020. Citizen science project reveals high diversity in *Didymellaceae* (Pleosporales, Dothideomycetes). *MycKeys* 65:49-99.
- Hu, J. H., Hong, C. X., Stromberg, E. L., and Moorman, G. W. 2008. Mefenoxam sensitivity and fitness analysis of *Phytophthora nicotianae* isolates from nurseries in Virginia, USA. *Plant Pathol.* 57:728-736.
- Hwang, J., and Benson, D. M. 2005. Identification, mefenoxam sensitivity, and compatibility type of *Phytophthora* spp. attacking floriculture crops in North Carolina. *Plant Dis.* 89:185-190.
- Hyakumachi, M., Priyatomo, A., Kubota, M., and Fukui, H. 2005. New anastomosis groups, AG-T and AG-U of binucleate *Rhizoctonia* spp. causing root and stem rot of cut-flower and miniature roses. *Phytopathology* 95:784-792.
- Jackson, C. R., and McFadden, L. A. 1961. Chrysanthemum diseases in Florida. Page 3 in: *Fla. Agric. Exp. Sta. Bull.* No. 635, vol. 47.
- Jung, T., Orlikowski, L., Henricot, B., Abad-Campos, P., Aday, A. G., Aguín Casal, O., Bakonyi, J., Cacciola, S. O., Cech, T., Chavarriaga, D., Corcobado, T., Cravador, A., Decourcelle, T., Denton, G., Diamandis, S., Dogmus-Lehtijärvi, H. T., Franceschini, A., Ginetti, B., Glavendekic, M., Hantula, J., Hartmann, G., Herrero, M., Ivic, D., Horta Jung, M., Lilja, A., Keca, N., Kramarets, V., Lyubenova, A., Machado, H., Magnano di San Lio, G., Mansilla Vázquez, P. J., Marçais, B., Matsiakh, I., Milenkovic, I., Moricca, S., Nagy, Z. A., Nechwatal, J., Olsson, C., Oszako, T., Pane, A., Paplomatas, E. J., Pintos Varela, C., Prospero, S., Rial Martínez, C., Rigling, D., Robin, C., Rytönen, A., Sánchez, M. E., Scanu, B., Schlenzig, A., Schumacher, J., Slavov, S., Solla, A., Sousa, E., Stenlid, J., Talgø, V., Tomic, Z., Tsopeles, P., Vannini, A., Vetrano, A. M., Wenneker, M., Woodward, S., and Pérez-Sierra, A. 2016. Widespread *Phytophthora* infestations in European nurseries put forest, seminatural and horticultural ecosystems at high risk of *Phytophthora* diseases. *For. Pathol.* 46:134-163.
- Kamoun, S., Furzer, O., Jones, J. D. G., Judelson, H. S., Ali, G. S., Dalio, R. J. D., Roy, S. G., Schena, L., Zambounis, A., Panabieres, F., Cahill, D., Ruocco, M., Figueiredo, A., Chen, X.-R., Hulvey, J., Stam, R., Lamour, K., Gijzen, M., Tyler, B. M., Grünwald, N. J., Mukhtar, M. S., Tomé, D. F. A., Tör, M., Van den Ackerveken, G., McDowell, J., Daayf, F., Fry, W. E., Lindqvist-Kreuzer, H., Meijer, H. J. G., Petre, B., Ristaino, J., Yoshida, K., Birch, P. R. J., and Govers, F. 2015. The top 10 oomycete pathogens in molecular plant pathology. *Mol. Plant Pathol.* 16:413-434.
- Kiss, L., and Berczyk, Z. 2011. First report of *Oidium longipes* as the causal agent of Petunia powdery mildew in the United Kingdom. *Plant Dis.* 95:361.
- Kiss, L., Jankovics, T., Kovacs, G. M., and Daughtrey, M. L. 2008. *Oidium longipes*, New Powdery mildew fungus on petunia in the USA: A potential threat to ornamental and vegetable solanaceous crops. *Plant Dis.* 92:818-825.
- Koike, S. T. 2000. Downy mildew of stock, caused by *Peronospora parasitica*, in California. *Plant Dis.* 84:103.
- Krebs, E. 1990. Rinden-Kultursubstrate und Schadpilze. *Deutscher Gartenbau* 44:2874-2877.
- Kruidhof, H. M., and Elmer, W. H. 2020. Cultural methods for greenhouse pest and disease management. Pages 285-330 in: *Integrated Pest and Disease Management in Greenhouse Crops*. M. L. Gullino, R. Albajes, and P. Nicot, eds. Springer, Dordrecht, The Netherlands.
- Laemmlen, F. 2001. *Alternaria* Diseases. Publ. no. 8040. Regents of University of California, Division of Agriculture and Natural Resources, Oakland, CA, U.S.A.
- Li, P., and Liu, D. 2019. First report of leaf spot caused by *Alternaria tenuissima* on *Coreopsis lanceolata* in Anqing, China. *Plant Dis.* 103:2131.
- Li, Y., Garibaldi, A., and Gullino, M. L. 2010. Molecular detection of *Fusarium oxysporum* f. sp. *chrysanthemi* on three host plants: *Gerbera jamesonii*, *Osteospermum* sp. and *Argyranthemum frutescens*. *J. Plant Pathol.* 92:525-530.
- Lin, S., Martin, D., Taylor, N. J., Gabriel, C., Devi Ganeshan, V., and Peduto Hand, F. 2018. First report of *Phytophthora* aerial blight caused by *Phytophthora nicotianae* on vinca, lobelia and calibrachoa in Ohio. *Plant Dis.* 102:456.
- Lin, S., Martin, D., Taylor, N. J., Gabriel, C., and Peduto Hand, F. 2017. Occurrence of *Phytophthora chrysanthemi* causing root and stem rot of garden mums in the United States. *Plant Dis.* 101:1060.
- Liu, B., Munster, M., Johnson, C., and Louws, F. J. 2011. First report of anthracnose caused by *Colletotrichum fragariae* on cyclamen in North Carolina. *Plant Dis.* 95:1480.
- Liu, J. B., Gilardi, G., Gullino, M. L., and Garibaldi, A. 2009. Effectiveness of *Trichoderma* spp. obtained from re-used soilless substrates against *Pythium ultimum* on cucumber seedlings [Wirksamkeit von *Trichoderma* spp. aus wiederverwendeten erdlosen Trägersubstraten gegenüber *Pythium ultimum* an Gurkensämlingen]. *J. Plant Dis. Prot.* 116:156-163.
- Lombard, L., Sandoval-Denis, M., Lamprecht, S. C., and Crous, P. W. 2019. Epitypification of *Fusarium oxysporum*—Clearing the taxonomic chaos. *Persoonia* 43:1-47.
- Lombard, L., Van der Merwe, N. A., Groenewald, J. Z., and Crous, P. W. 2015. Generic concepts in *Nectriaceae*. *Stud. Mycol.* 80:189-245.
- Matić, S., Gilardi, G., Gullino, M. L., and Garibaldi, A. 2018. Evidence for an expanded host range of *Fusarium oxysporum* f. sp. *chrysanthemi*. *J. Plant Pathol.* 100:97-104.
- Matić, S., Tabone, G., Garibaldi, A., and Gullino, M. L. 2020. *Alternaria* leaf spot caused by *Alternaria* species: An emerging problem on ornamental plants in Italy. *Plant Dis.* 104:2275-2287.
- McGrath, M. T., and Shishkoff, N. 2001. Resistance to triadimefon and benomyl: Dynamics and impact on managing cucurbit powdery mildew. *Plant Dis.* 85:147-154.
- McMillan, R. T., and Graves, W. R. 1996. Periwinkle twig blight caused by *Colletotrichum dematium* on *Catharanthus roseus*. *Proc. Fla. State Hortic. Soc.* 109:19-20.
- McTaggart, A. R., Van der Nest, M. A., Steenkamp, E. T., Roux, J., Slipers, B., Shuey, L. S., Wingfield, M. J., and Drenth, A. 2016. Fungal genomics challenges the dogma of name-based biosecurity. *PLoS Pathog.* 12:e1005475.
- Minerdi, D., Moretti, M., Li, Y. L., Garibaldi, A., and Gullino, M. L. 2008. Conventional PCR and real time quantitative PCR detection of *Phytophthora cryptogea* on *Gerbera jamesonii*. *Eur. J. Plant Pathol.* 122:227-237.
- Minuto, A., Gullino, M. L., and Garibaldi, A. 2007. *Gerbera jamesonii*, *Osteospermum* sp. and *Argyranthemum frutescens*: New hosts of *Fusarium oxysporum* f. sp. *chrysanthemi*. *J. Phytopathol.* 155:373-376.
- Moorman, G. W., Kang, S., Geiser, D. M., and Kim, S. H. 2002. Identification and characterization of *Pythium* species associated with greenhouse floral crops in Pennsylvania. *Plant Dis.* 86:1227-1231.
- Munkvold, G. P. 2009. Seed pathology progress in the academia and industry. *Annu. Rev. Phytopathol.* 47:285-311.
- Munkvold, G. P., and Gullino, M. L. 2020. Seed and propagative material. Pages 331-354 in: *Integrated Pest and Disease Management in Greenhouse Crops*. M. L. Gullino, R. Albajes, and P. Nicot, eds. Springer, Dordrecht, The Netherlands.
- Naher, M., Motohash, K., Watanabe, H., Chikuo, Y., Senda, M., Suga, H., Brasier, C., and Kageyama, K. 2011. *Phytophthora chrysanthemi* sp. nov., a new species causing root rot of chrysanthemum in Japan. *Mycol. Prog.* 10:21-31.
- Noble, R. 2011. Risks and benefits of soil amendment with composts in relation to plant pathogens. *Australas. Plant Pathol.* 40:157-167.
- Noble, R., and Coventry, E. 2005. Suppression of soil-borne plant diseases with composts: A review. *Biocontrol Sci. Technol.* 15:3-20.
- O'Donnell, K., Sutton, D. A., Fothergill, A., McCarthy, D., Rinaldi, M. G., Brandt, M. E., Zhang, N., and Geiser, D. M. 2008. Molecular phylogenetic diversity, multilocus haplotype nomenclature, and in vitro antifungal resistance within the *Fusarium solani* species complex. *J. Clin. Microbiol.* 46:2477-2490.

- Ogoshi, A. 1996. Introduction – the genus *Rhizoctonia*. Pages 1-9 in: *Rhizoctonia* Species: Taxonomy, Molecular Biology, Ecology, Pathology, and Disease Control. B. Sneh, S. Jabaji-Hare, S. Neate, and G. Dijst, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Olson, H. A., and Benson, D. M. 2007. Induced systemic resistance and the role of binucleate *Rhizoctonia* and *Trichoderma hamatum* 382 in biocontrol of Botrytis blight in geranium. *Biol. Control* 42:233-241.
- Olson, H. A., and Benson, D. M. 2011. Characterization of *Phytophthora* spp. on floriculture crops in North Carolina. *Plant Dis.* 95:1013-1020.
- Olson, H. A., Jeffers, S. N., Ivors, K. L., Steddom, K. C., Williams-Woodward, J. L., Mmbaga, M. T., Benson, D. M., and Hong, C. X. 2013. Diversity and mefenoxam sensitivity of *Phytophthora* spp. associated with the ornamental horticulture industry in the southeastern United States. *Plant Dis.* 97:86-92.
- Ortega, L. 1999. Prevention and eradication of white rust (*Puccinia horiana*) in Colombia. *Acta Hortic.* 482:187-196.
- Palmateer, A. J., Harmon, P. F., and Schubert, T. S. 2008. Downy mildew of coleus (*Solenostemon scutellarioides*) caused by *Peronospora* sp. in Florida. *Plant Pathol.* 57:372.
- Pane, C., Spaccini, R., Piccolo, A., Scala, F., and Bonanomi, G. 2011. Compost amendments enhance peat suppressiveness to *Pythium ultimum*, *Rhizoctonia solani* and *Sclerotinia minor*. *Biol. Control* 56:115-124.
- Patil, V. B., Mali, A. M., Mahamuni, R. J., Chavan, N. S., and Kamble, S. S. 2012. First report of leaf spot caused by *Phoma costarricensis* on *Delphinium malabaricum* in Western Ghats of India. *Plant Dis.* 96:1074.
- Pennycook, S. R. 1989. Plant Diseases Recorded in New Zealand, Vol. 3. Plant Diseases Division, DSIR, Auckland, New Zealand.
- Polizzi, G., Aiello, D., Guarnaccia, V., Perrone, G., and Stea, G. 2011. First report of damping-off on strawberry tree caused by *Colletotrichum acutatum* and *C. simmondsii* in Italy. *Plant Dis.* 95:1588.
- Prigigallo, M. I., Mosca, S., Cacciola, S. O., Cooke, D. E. L., and Schena, L. 2015. Molecular analysis of *Phytophthora* diversity in nursery-grown ornamental and fruit plants. *Plant Pathol.* 64:1308-1319.
- Pugliese, M., Gilardi, G., Garibaldi, A., and Gullino, M. L. 2015. Organic amendments and soil suppressiveness: Results with vegetable and ornamental crops. Pages 495-509 in: *Organic Amendments and Soil Suppressiveness in Plant Disease Management*. Soil Biology, Vol. 46. M. K. Meghvansi and A. Varma, eds. Springer International Publishing, Cham, Switzerland.
- Pugliese, M., Gullino, M. L., and Garibaldi, A. 2012. Compost suppressiveness against *Phytophthora* spp. on *Skimmia japonica* and azalea. *Commun. Agric. Appl. Biol. Sci.* 77:237.
- Pugliese, M., Liu, B. P., Gullino, M. L., and Garibaldi, A. 2008. Selection of antagonists from compost to control soil-borne pathogens [Selektion von Antagonisten aus Kompost zur Kontrolle bodenbürtiger Pathogene]. *J. Plant Dis. Prot.* 115:220-228.
- Pugliese, M., Liu, B. P., Gullino, M. L., and Garibaldi, A. 2011. Microbial enrichment of compost with biological control agents to enhance suppressiveness to four soil-borne diseases in greenhouse. *J. Plant Dis. Prot.* 118:45-50.
- Rinehart, T. A., Copes, W. E., Toda, T., and Cubeta, M. A. 2007. Genetic characterization of binucleate *Rhizoctonia* species causing web blight on azalea in Mississippi and Alabama. *Plant Dis.* 91:616-623.
- Rivera, Y., Salgado-Salazar, C., Windham, A. S., and Crouch, J. A. 2016. Downy mildew on coleus (*Plectranthus scutellarioides*) caused by *Peronospora belbahrii* sensu lato in Tennessee. *Plant Dis.* 100:655.
- Schaad, N. W., Frederick, R. D., Shaw, J., Schneider, W. L., Hickson, R., Petrillo, M. D., and Luster, D. G. 2003. Advances in molecular-based diagnostic in meeting crop biosecurity and phytosanitary issues. *Annu. Rev. Phytopathol.* 41:305-324.
- Scheuerell, S. J., Sullivan, D. M., and Mahaffee, W. F. 2005. Suppression of seedling damping-off caused by *Pythium ultimum*, *P. irregulare*, and *Rhizoctonia solani* in container media amended with a diverse range of Pacific Northwest compost sources. *Phytopathology* 95:306-315.
- Schwingle, B. W., Smith, J. A., and Blanchette, R. A. 2007. *Phytophthora* species associated with diseased woody ornamentals in Minnesota nurseries. *Plant Dis.* 91:97-102.
- Shishkoff, N. 2019. The oospore stage of *Plasmopara obducens*, impatiens downy mildew. *Mycologia* 111:632-646.
- Shoresh, M., Harman, G. E., and Mastouri, F. 2010. Induced systemic resistance and plant responses to fungal biocontrol agents. *Annu. Rev. Phytopathol.* 48:21-43.
- Silva, M. T. B., Mendoça, A. M., Seijo, Y. C., and Viquiera, F. D. F. 2007. Assessment of municipal solid waste compost quality using standardized methods before preparation of plant growth media. *Waste Manage. Res.* 25:99-108.
- Sinclair, W. A., and Lyon, H. H. 2005. Page 660 in: *Diseases of Trees and Shrubs*. Cornell University Press, Ithaca, NY, U.S.A.
- Slippers, B., and Wingfield, M. J. 2007. Botryosphaeriaceae as endophytes and latent pathogens of woody plants: Diversity, ecology and impact. *Fungal Biol. Rev.* 21:90-106.
- Sneh, B., Burpee, L., and Ogoshi, A. 1991. Identification of *Rhizoctonia* Species. American Phytopathological Society, St. Paul, MN, U.S.A.
- Spadaro, D., Agustí, N., Ortega, S. F., and Hurtado Ruiz, M. A. 2020. Diagnostics and identification of diseases, insects and mites. Pages 231-258 in: *Integrated Pest and Disease Management in Greenhouse Crops*. M. L. Gullino, R. Albajes, and P. Nicot, eds. Springer, Dordrecht, The Netherlands.
- Srinivasan, K., Gilardi, G., Garibaldi, A., and Gullino, M. L. 2009. Bacterial antagonists from used rockwool soilless substrates suppress Fusarium wilt of tomato. *J. Plant Pathol.* 91:147-154.
- Stebbins, T., and Johnson, D. 2001. Regulatory control. Pages 457-458 in: *Diseases of Woody Ornamentals and Trees in Nurseries*. R. K. Jones and D. M. Benson, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Sullivan, D. M., and Miller, R. O. 2001. Compost quality attributes, measurements, and variability. Pages 95-120 in: *Compost Utilization in Horticultural Cropping Systems*. P. J. Stofella and B. A. Kahn, eds. Lewis Publishers, New York, NY, U.S.A.
- Sun, H., Song, C., Mubeen, I., and Huang, J. 2019. First report of *Alternaria tenuissima* causing leaf spot disease on *Iris tectorum* in China. *Plant Dis.* 103:1786.
- Tahvonen, O., and Kuuluvainen, J. 1993. Economic growth, pollution, and renewable resources. *J. Environ. Econ. Manage.* 24:101-118.
- Termorshuizen, A. J., van Rijn, E., van der Gaag, D. J., Alabouvette, C., Chen, Y., Lagerlöf, J., Malandrakis, A. A., Paplomatas, E. J., Rämert, B., Ryckeboer, J., Steinberg, C., and Zmora-Nahum, S. 2006. Suppressiveness of 18 composts against 7 pathosystems: Variability in pathogen response. *Soil Biol. Biochem.* 38:2461-2477.
- Thomas, J. E., Wood, T. A., Gullino, M. L., and Ortu, G. 2017. Diagnostic tools for plant biosecurity. Pages 209-226 in: *Practical Tools for Plant and Food Biosecurity*. M. L. Gullino, J. P. Stack, J. Fletcher, and J. D. Mumford, eds. Springer, Dordrecht, The Netherlands.
- Thomas, S., Wahabzada, M., Kuska, M. T., Rascher, U., and Mahlein, A. K. 2017. Observation of plant-pathogen interaction by simultaneous hyperspectral imaging reflection and transmission measurements. *Funct. Plant Biol.* 44:23-34.
- Thomma, B. P. 2003. *Alternaria* spp.: From general saprophyte to specific parasite. *Mol. Plant Pathol.* 4:225-236.
- Tognoni, F., and Incrocci, L. 2003. Le colture fuori suolo: Situazione in Italia e prospettive per il futuro. *Inf. Fitopatol.* 53:7-12.
- Tomic, Z., and Ivic, D. 2015. *Phytophthora chrysanthemi* Naher, Motohash, Watanabe, Chikuo, Senda, Suga, Brasier & Kageyama—New cause of chrysanthemum disease in Croatia. *Glas. Biljne Zast.* 15:291-300.
- Tompkins, R. M., and Snyder, W. C. 1972. Cyclamen wilt in California and its control. *Plant Dis.* 56:493-497.
- Troisi, M., Bertetti, D., Garibaldi, A., and Gullino, M. L. 2010a. First report of powdery mildew caused by *Golovinomyces cichoracearum* on Gerbera (*Gerbera jamesonii*) in Italy. *Plant Dis.* 94:130.
- Troisi, M., Bertetti, D., Gullino, M. L., and Garibaldi, A. 2013. Race differentiation in *Fusarium oxysporum* f. sp. *chrysanthemi*. *J. Phytopathol.* 161:675-688.
- Troisi, M., Gullino, M. L., and Garibaldi, A. 2010b. *Gerbera jamesonii*, a new host of *Fusarium oxysporum* f. sp. *tracheiphilum*. *J. Phytopathol.* 158:8-14.
- USDA. 2019. Floriculture Crops: 2018 Summary. United States Department of Agriculture—National Agricultural Statistics Service. https://www.nass.usda.gov/Publications/Todays_Reports/reports/floran19.pdf
- Van Os, E. A., Hofland-Zijlstra, J. D., and Postma, J. 2012. Management of irrigation water as a tool for disease management. Pages 75-82 in: *Fusarium Wilts of Greenhouse Vegetable and Ornamental Crops*. M. L. Gullino, J. Katan, and A. Garibaldi, eds. American Phytopathological Society, St. Paul, MN, U.S.A.
- Verrier, P. J., Thomas, J. E., and Stack, J. P. 2017. The European Union Plant Diagnostic Information System (EUPDIS): A platform for collaborative diagnostics and a tool for early detection of plant pathogens. Pages 227-241 in: *Practical Tools for Plant and Food Biosecurity*. M. L. Gullino, J. P. Stack, J. Fletcher, and J. D. Mumford, eds. Springer, Dordrecht, The Netherlands.
- Wahyuno, D., Susan, D., and Kakishima, M. 2012. Occurrence of a rust disease of *Fuchsia* caused by *Pucciniastrum epilobii* in Indonesia. *Australas. Plant Dis. Notes* 7:143-146.
- Walters, D. R., and Fountaine, J. M. 2009. Practical application of induced resistance to plant diseases: An appraisal of effectiveness under field conditions. *J. Agric. Sci.* 147:523-535.
- Walters, D. R., Ratsep, J., and Havis, N. D. 2013. Controlling crop diseases using induced resistance: Challenges for the future. *J. Exp. Bot.* 64:1263-1280.
- Wang, B., and Jeffers, S. N. 2002. Effects of cultural practices and temperature on Fusarium root and crown rot of container-grown hostas. *Plant Dis.* 86:225-231.
- Warfield, C. Y., Hwang, J., and Benson, D. M. 2008. *Phytophthora* blight and dieback in North Carolina nurseries during a 2003 survey. *Plant Dis.* 92:474-481.

- Werres, S., Marwitz, R., Man In't Veld, W. A., De Cock, A. W. A. M., Bonants, P. J. M., De Weerd, M., Themann, K., Ilieva, E., and Baayen, R. P. 2001. *Phytophthora ramorum* sp. nov., a new pathogen on *Rhododendron* and *Viburnum*. Mycol. Res. 105:1155-1165.
- Williams-Woodward, J. L., Hennen, J. F., Parda, K. W., and Fowler, J. M. 2001. First report of daylily rust in the United States. Plant Dis. 85:1121.
- Wise, K. A., Mueller, D. S., and Buck, J. W. 2004. Quarantines and ornamental rusts. <https://www.apsnet.org/edcenter/apsnetfeatures/Pages/OrnamentalRusts.aspx>
- Woudenberg, J. H. C., Groenewald, J. Z., Binder, M., and Crous, P. W. 2013. *Alternaria* redefined. Stud. Mycol. 75:171-212.
- Woudt, L. P. 1995. Identification of *Fusarium oxysporum* f. sp. *cyclaminis* by DNA fingerprinting and PCR 1. Bull. OEPP 25:109-112.
- Wu, W. S., Li, Y. L., and Wu, H. C. 2006. Seed-borne fungi of ornamental flower plants. Plant Pathol. 35:373-375.
- Yakabe, L. E., Blomquist, C. L., Thomas, S. L., and MacDonald, J. D. 2009. Identification and frequency of *Phytophthora* species associated with foliar diseases in California ornamental nurseries. Plant Dis. 93: 883-890.
- Yamak F., Peever, T. L., Grove, G. G., and Boal, R. J. 2002. Occurrence and identification of *Phytophthora* spp. pathogenic to pear fruit in irrigation water in the Wenatchee River valley of Washington state. Phytopathology 92:1210-1217.
- Zhang, X., Jiang, B., Wang, R., Li, Y., and Sun, L. 2020. Occurrence of leaf spot of *Begonia semperflorens* caused by *Alternaria tenuissima* in China. Plant Dis. 104:2292.