Resistance in *Capsicum pubescens* to *Xanthomonas campestris* pv. *vesicatoria* Pepper Race 6

F. Sahin and S. A. Miller, Department of Plant Pathology, The Ohio State University, Ohio Agricultural Research and Development Center, Wooster 44691

ABSTRACT

Sahin, F., and Miller, S. A. 1998. Resistance in *Capsicum pubescens* to *Xanthomonas campestris* pv. *vesicatoria* pepper race 6. Plant Dis. 82:794-799.

One hundred seventy *Capsicum* spp. germplasm accessions were evaluated as potential sources of resistance to *X. campestris* pv. *vesicatoria* pepper race 6 (P6). This race has been identified recently in Ohio and overcomes all three known resistance genes in cultivated pepper. Only *C. pubescens* plant introduction (PI) 235047 was found to be resistant to *X. campestris* pv. *vesicatoria* P6 strains using hypersensitivity and pathogenicity tests. Further studies using PI 235047 as a differential line showed that strains classified as *X. campestris* pv. *vesicatoria* races P1 and P3 were heterogeneous in terms of virulence. Strains of *X. campestris* pv. *vesicatoria* P1 and P3 that were compatible with PI 235047 were reclassified as two new pepper races, designated *X. campestris* pv. *vesicatoria* P7 and P8, respectively. According to the new race classification, PI 235047 plants are incompatible with *X. campestris* pv. *vesicatoria* pepper races P0, P1, P3, P4, and P6, but compatible with races P2, P5, P7, and P8.

Bacterial spot of pepper (*Capsicum* spp.), caused by *Xanthomonas campestris* pv. *vesicatoria* (proposed to be renamed *Xanthomonas axonopodis* pv. *vesicatoria*; 37), is one of the most destructive pepper diseases both in the greenhouse and in the field when environmental conditions are favorable for the pathogen (8,24,25,33). The current management methods for bacterial spot, including sanitation, chemical application, cultural practices, and use of *X. campestris* pv. *vesicatoria*-resistant cultivars (10,14,17,23–25,27,29,33,36), are not always successful. Contaminated seeds are the main source of *X. campestris* pv. *vesicatoria* inoculum (3–5,17). However, seed treatment, which may not completely eliminate *X. campestris* pv. *vesicatoria* inoculum if internal seed infection occurs (5), is not always done, for economic reasons as well as potential reduction of seed viability. Cost, potential chemical residues on fruit, and resistance among *X. campestris* pv. *vesicatoria* strains are the main disadvantages of chemical applications (2,27,29,33,36). Therefore, use of pepper cultivars resistant to *X. campestris* pv.

vesicatoria is economically and technically the most practical method for bacterial spot management (23–25). There is an increasing interest in developing pepper cultivars with resistance to bacterial spot. However, development of disease-resistant cultivars requires the identification and incorporation of resistance genes into horticulturally superior and widely acceptable cultivars. Good progress has been made in identification of suitable sources of disease resistance by screening a number of accessions of cultivated pepper species and related wild species with great genetic diversity (11–13,15,16,20). So far, three genes for resistance to bacterial spot, designated *Bs1*, *Bs2*, and *Bs3*, have been identified in three different plant introduction (PI) lines: PI 163192 (*C. annuum*), PI 260435 (*C. chacoense*), and PI 271322 (*C. annuum*), respectively (11,13,20).

Resistance in pepper to strains of *X. campestris* pv. *vesicatoria* is associated with the hypersensitive response (HR; 1,6–9,11–13,15,16,18–21,28,30,34). Currently, seven pepper races (P0–P6) have been identified among *X. campestris* pv. *vesicatoria* strains worldwide (8,22,28,29,32,33). Pepper races are easily identified based on the presence or absence of HR in the *X. campestris* pv. *vesicatoria*-susceptible cultivar Early Calwonder (ECW) and its near-isogenic derivative lines, designated ECW-10R, ECW-20R, and ECW-30R, which contain the resistance genes *Bs1*, *Bs2*, and *Bs3*, respectively (28). Race P6 strains, first identified in Ohio in 1994 (32), are able to defeat all of the known resistance genes in pepper (24,25,32,33). The number of *X. campestris* pv. *vesicatoria* P6 strains is increasing in Ohio and

most of these are resistant to copper, streptomycin, or both (31). Therefore, new sources of resistance in pepper need to be identified.

The objectives of this study were to (i) evaluate *Capsicum* germplasm for sources of resistance to *X. campestris* pv. *vesicatoria* race P6 and (ii) utilize newly identified resistant pepper germplasm to determine the existence of additional variation in virulence among strains of other *X. campestris* pv. *vesicatoria* pepper races.

MATERIALS AND METHODS

Plant material and growing conditions. Seeds for all PI lines of *Capsicum* spp., including 119 accessions of *C. annuum* and 51 accessions of other *Capsicum* spp. (*C. anomallum*, *C. baccatum*, *C. chacoense*, *C. chinense*, *C. frutescens*, *C. galapagoense*, and *C. pubescens*) originating from 40 countries were obtained from the United States Department of Agriculture-Agriculture Research Station (USDA-ARS) Plant Genetic Resources Conservation Unit, University of Georgia, Griffin (Table 1).

A total of 20 seeds from each of the 170 accessions and the bell pepper cultivar Marengo (*X. campestris* pv. *vesicatoria*-susceptible) were sown in commercial peat mix (Baccto High Density Professional Planting Mix, Michigan Peat Co., Houston) in new 128-cell plastic flats (Landmark Plastics Corp., Akron, OH). The seedlings were grown on the greenhouse bench until they reached the second or third true-leaf stage (approximately 3 to 4 weeks). Then 15 seedlings of each accession were transplanted into five 25-cm-diameter plastic pots (three plants/pot) containing steamed soil (mixture of 0.43 m³ Wooster silty-loam, 0.43 m³ muck soil, 0.14 m³ peat moss, and 1.2 kg lime/m³). The plants were maintained for 7 days in the greenhouse (20 to 28°C) after transplanting. Plants were fertilized once a week with Peter's 20-20-20 (N-P-K; 454 mg/378 ml) and watered as required. Nine plants from each accession were used to test pathogenicity and six plants were used to test HR elicitation. This experiment was conducted twice. The remaining five seedlings of the PI accessions and Marengo were used as negative controls in the experiments.

Inoculum preparation. A highly aggressive strain of *X. campestris* pv. *vesicatoria* pepper race 6, *X. campestris* pv.

Corresponding author: S. A. Miller
E-mail address: miller.769@osu.edu

This research was supported by Ataturk University, Erzurum, Turkey, and State and Federal Funds appropriated to the Ohio Agricultural Research and Development Center (OARDC), The Ohio State University. OARDC Research Article Number 148-97.

Accepted for publication 24 March 1998.

Publication no. D-1998-0518-01R
© 1998 The American Phytopathological Society

vesicatoria 17b, isolated from pepper in 1994 in Ohio, was used. Strain *X. campestris* pv. *vesicatoria* 17b was stored frozen (−80°C) in 15% glycerol and recovered from frozen culture by restreaking onto yeast dextrose carbonate agar (YDC) medium (26) and incubating at 28°C for 48 h. The bacteria were washed from the plate with a sterile glass spreader and suspended in sterile distilled water. The bacterial concentration was adjusted to approximately 10⁸ CFU/ml ($A_{600nm} = 0.1$). This concentration was used for inoculation and infiltration of pepper seedlings.

Plant inoculation. Nine 5-week-old pepper seedlings (three plants/pot) of each accession and Marengo were dipped into 4 liters bacterial suspension in a 5-liter plastic container. Five Marengo seedlings, dipped into sterilized distilled water, were used as a negative control in each experiment. The inoculated plants were arranged in a completely randomized design and incubated in a mist chamber at 90 to 95% relative humidity (RH) for 3 days, then transferred to a greenhouse at 20 to 28°C for 7 days. The control plants were incubated under the same conditions, but in different greenhouses. The plants were observed for development of characteristic bacterial spot symptoms 10 days after inoculation. Plants were rated as positive or negative based on the presence or absence of disease symptoms. Pathogenicity was evaluated based on symptom development

on Marengo. To test for elicitation of an HR, a bacterial suspension (10⁸ CFU/ml), prepared as described above, was infiltrated into the intercostal area of the leaves of two plants of each PI line and Marengo by using a 3-cc syringe (Becton Dickinson and Co., Franklin Lakes, NJ) without a hypodermic needle. Sterilized distilled water was infiltrated into the leaves of plants of all PI lines and Marengo as negative controls. The inoculated plants were arranged in a completely randomized design on the greenhouse bench and maintained for 24 to 48 h at 20 to 28°C. The presence of rapid tissue necrosis at the inoculation site was recorded within 24 to 48 h after infiltration. This test was repeated at least three times.

Variation in host specificity, virulence and growth in plants. Seventy-two strains, including all of the known *X. campestris* pv. *vesicatoria* pepper races (P0 to P6), were used. Strains V-19 and V-205, and strains 1138, 1140, and 1141 were obtained from D. F. Ritchie and F. Louws, respectively (Department of Plant Pathology, North Carolina State University, Raleigh). The remaining 67 strains were isolated from pepper or tomato plants collected from commercial fields in Ohio between 1993 and 1996. All strains were tested for HR elicitation on *C. pubescens* PI 235047, ECW, ECW-10R, ECW-20R, and ECW-30R as described above. The cultivar Marengo was used as a negative

control. The results were recorded as positive or negative based on the presence or absence, respectively, of collapsed tissue at the inoculation site within 48 h after infiltration. This experiment was conducted four times.

Of the 72 *X. campestris* pv. *vesicatoria* strains tested for HR on PI 235047, 58 were also tested for pathogenicity on this accession and Marengo. Two plants of each accession were dip-inoculated with individual strains as described above. Two plants of each line were dipped in sterile water as a negative control. The inoculated plants were arranged in a completely randomized design with two replications of a single plant each per experiment. The inoculated plants were incubated in a mist chamber at 90 to 95% RH for 3 days, then transferred to the greenhouse at 20 to 28°C for 7 days. The plants were rated for disease severity 10 days after inoculation by using a scale of 1 to 5, in which 1 = no disease, 2 = a few water-soaked lesions, 3 = many spots with coalescence and slight plant wilting, 4 = severe leaf defoliation, and 5 = plants dead. Average disease severity values (DSVs) for each of the bacterial strains on PI 235047 and Marengo plants were calculated from two experiments. The data were evaluated with analysis of variance (ANOVA), and differences among the treatment means were determined by calculating Fisher's least significant difference at $P = 0.05$.

Table 1. Evaluation of *Capsicum* spp. accessions for resistance to *Xanthomonas campestris* pv. *vesicatoria* race P6

Host/PI	HR ^a	Disease ^b
<i>Capsicum annuum</i>		
102881, 102882, 102883, 103048, 105262, 105444, 105339, 105538, 109252, 109254, 109255, 109469, 123164, 123165, 123166, 123469, 123470, 123471, 124080, 124540, 125801, 125802, 125803, 127445, 131352, 135826, 135873, 135874, 138557, 138558, 138559, 138561, 143886, 148626, 153565, 155349, 159226, 159227, 159228, 159255, 159265, 159278, 162607, 162608, 164556, 164557, 164558, 164561, 164562, 164564, 164565, 164678, 164961, 167062, 167100, 167215, 169102, 169106, 169110, 169115, 169122, 169130, 169131, 169138, 176888, 181732, 181733, 181734, 182646, 184037, 184038, 184039, 187314, 187315, 187325, 188472, 188473, 188474, 249907, 249911, 251622, 257049, 257187, 262172, 262905, 263073, 263074, 263106, 263110, 264662, 267730, 267731, 267733, 275008, 275009, 275011, 281316, 281323, 281327, 281330, 281341, 281400, 281413, 281416, 286419, 289762, 291999, 294452, 297456, 338490, 338974, 385960, 385961, 390436, 390437, 390970, 390971, 401728, 401729	—	+
<i>C. annuum</i> var. <i>glabriusculum</i>		
511885, 511886, 511887	—	+
<i>C. anomallum</i>		
501532	—	+
<i>C. baccatum</i>		
260434, 273425, 281398, 3700004, 3700005, 413669, 439359, 439360, 439361, 439364, 439369	—	+
<i>C. baccatum</i> var. <i>baccatum</i>		
215699, 238061, 260533, 260567, 281306	—	+
<i>C. baccatum</i> var. <i>pendulum</i>		
152217, 152234, 199506, 203522, 213915, 224440, 238062, 241679, 257122, 241674, 159235, 188803	—	+
<i>C. baccatum</i> var. <i>praetermissum</i>		
260595	—	+
<i>C. chacoense</i>		
260426, 260427, 260430, 260431, 273429	—	+
<i>C. chinense</i>		
152222, 152225, 152452, 194879, 200729, 209589	—	+
<i>C. frutescens</i>		
123474, 159238, 181865, 188479	—	+
<i>C. galapagoense</i>		
G-1567	—	+
<i>C. pubescens</i>		
235047	+	—
585266	—	+

^a HR = hypersensitive response; — = HR not elicited; + = HR elicited.

^b Disease = presence (+) or absence (−) of symptoms.

Another experiment was conducted in the greenhouse to characterize disease resistance in *C. pubescens* PI 235047 plants to *X. campestris* pv. *vesicatoria* P6 strains based on bacterial population dynamics in inoculated leaves. Two plants each of PI 235047 and Marengo were inoculated separately with *X. campestris* pv. *vesicatoria* 17b (race P6) and *X. campestris* pv. *vesicatoria* 116 (race P1), a virulent strain that did not elicit an HR on PI 235047. Inoculation, incubation, and disease assessment were done as described above. All of the leaves on two inoculated plants of each cultivar were collected after scoring disease severity. Half of the leaves were surface sterilized by dipping into 95% EtOH, and the others were dipped into sterile distilled water. The leaves were

kept on paper towels for 30 min to allow the alcohol to evaporate. Then the leaf samples were macerated using a ball bearing tissue grinder (BioReba AG, Basel, Switzerland) in a separate plastic bag (Agdia Inc, Elkhorn, IN) containing 1 ml sterile distilled water/g leaf tissue (fresh weight). Serial 10-fold dilutions were performed in sterile distilled water and 0.1 ml of the appropriate dilutions was spread onto each of two plates of CKTM medium (35). Characteristic *X. campestris* pv. *vesicatoria* colonies were counted after the plates were incubated for 4 days at 28°C.

RESULTS

Characteristic symptoms of bacterial spot were observed on all Marengo plants, and all but one of the *Capsicum* spp. accessions inoculated with *X. campestris* pv. *vesicatoria* race P6, strain 17b. *C. pubescens* PI 235047 plants (Fig. 1) did not develop distinguishable symptoms of bacterial spot in either experiment (Table

1). The negative control Marengo plants inoculated with sterilized water also did not show any recognizable symptoms.

An HR to *X. campestris* pv. *vesicatoria* 17b was elicited in all infiltrated leaves of PI 235047 plants within 24 h after inoculation (Table 1, Fig. 2). None of the other accessions or Marengo developed an HR during the 24 to 48-h incubation period after infiltration. HR was not observed on leaves of *C. pubescens* PI 235047 or Marengo plants infiltrated with sterilized distilled water.

Variation in host specificity, virulence and growth in plants. Of the 72 *X. campestris* pv. *vesicatoria* strains tested, 46 (64%) were race P3 (Table 2). The remaining strains were distributed as follows: 1 P0, 10 P1, 2 P2, 3 P4, 1 P5, and 10 P6. Although 41 strains (60%) elicited an HR on *C. pubescens* PI 235047, the remaining 31 strains (40%) were compatible. All strains of *X. campestris* pv. *vesicatoria* races P0, P4, and P6 and almost half of the P1 and P3 strains tested were incompati-

Fig. 1. *Capsicum pubescens* PI 235047 plants. (A) Flower (purple), (B) fruit (yellow).

Fig. 2. Leaves of *Capsicum pubescens* PI 235047 plants infiltrated with a suspension of *Xanthomonas campestris* pv. *vesicatoria* (A) race P6 or (B) race P1. (A) Incompatible reaction, (B) compatible reaction.

Table 2. Hypersensitive response (HR) in *Capsicum pubescens* PI 235047 plants and average disease severity (DSV) on PI 235047 and the pepper cultivar Marengo inoculated with strains of different pepper races of *Xanthomonas campestris* pv. *vesicatoria*

Strain ^b	Race ^c	HR test ^d	DSV ^a	
			PI 235047	Marengo
V-19	P0	+	1.00	2.75
Xcv-63	P1	+	1.25	3.25
Xcv-118	P1	+	1.00	3.50
Xcv-119a	P1	+	1.00	3.75
Xcv-119b	P1	+	1.25	4.00
Xcv-120	P1	+	1.75	3.25
Xcv-106a	P3	+	1.75	3.75
Xcv-106b	P3	+	1.25	3.50
Xcv-107a	P3	+	1.00	2.75
Xcv-107b	P3	+	1.25	3.50
Xcv-108	P3	+	1.00	3.75
Xcv-109a	P3	+	1.50	3.25
Xcv-109b	P3	+	1.50	3.50
Xcv-112	P3	+	1.00	3.25
Xcv-113	P3	+	1.75	2.75
Xcv-121	P3	+	1.25	2.75
Xcv-122a	P3	+	1.00	3.50
Xcv-122b	P3	+	1.25	3.75
Xcv-122c	P3	+	1.25	3.50
Xcv-122d	P3	+	1.00	3.00
Xcv-127a	P3	+	1.25	3.75
Xcv-127b	P3	+	1.00	3.75
Xcv-127c	P3	+	1.25	3.25
Xcv-129a	P3	+	1.00	3.50
Xcv-129b	P3	+	1.00	4.00
Xcv-130	P3	+	1.00	3.50
Xcv-794a	P3	+	NT	NT
Xcv-794b	P3	+	NT	NT
1138	P4	+	NT	NT
1140	P4	+	NT	NT
1141	P4	+	NT	NT

(continued on next page)

^a Average disease severity values calculated from two experiments, each with two replications, after the plants were rated using a 1 to 5 scale where 1 = no disease, 2 = a few water-soaked lesions, 3 = many spots with coalescence and slight plant wilting, 4 = severe leaf defoliation, and 5 = plants dead; NT = not tested.

^b *Xanthomonas campestris* pv. *vesicatoria* strains isolated from pepper or tomato.

^c Race of the *X. campestris* pv. *vesicatoria* strains were determined by infiltration of bacterial suspension (10⁸ CFU/ml) into the leaves of pepper differential lines.

^d Strains of *X. campestris* pv. *vesicatoria* eliciting a hypersensitive response on PI 235047 plants; + = incompatible (resistant) reaction, - = compatible (susceptible) reaction.

^e ANOVA = analysis of variance; LSD = least significant difference.

ble. All of the P2 and P5 strains were compatible on this accession.

All of the strains that elicited an HR (incompatible) on *C. pubescens* PI 235047 caused virtually no symptoms on PI 235047 plants after dip-inoculation. All of the *X. campestris* pv. *vesicatoria* strains that did not elicit an HR (compatible) on PI 235047 caused serious disease on PI 235047 plants. The average DSVs for PI 235047 plants inoculated with *X. campestris* pv. *vesicatoria* strains in the incompatible group (DSVs = 1 to 1.75) were significantly ($P = 0.05$) less than those (DSVs = 2.25 to 3.75) for plants inoculated with compatible strains (Table 2). Strains of *X. campestris* pv. *vesicatoria* in both groups caused serious disease on cultivar Marengo. While there were some significant differences between individual strains of *X. campestris* pv. *vesicatoria* in virulence on PI 235047 and Marengo, there were no differences between groups in virulence on Marengo (average DSVs for the compatible group = 3.5, incompatible group = 3.42). However, the DSVs of PI 235047 plants inoculated with compatible strains of *X. campestris* pv. *vesicatoria*

(average DSV = 2.8) were significantly less than those of Marengo plants (average DSV = 3.5) inoculated with these strains.

Without surface sterilization, the average density of *X. campestris* pv. *vesicatoria* 17b (P6) in PI 235047 leaves was 6×10^5 CFU/g, significantly ($P = 0.05$) less than the bacterial density (3.9×10^9 CFU/g) in Marengo leaves. After surface sterilization, the average density of *X. campestris* pv. *vesicatoria* 17b in PI 235047 leaves was 3.55×10^3 CFU/g, also significantly less than in Marengo leaves (7.2×10^7 CFU/g). The population density of *X. campestris* pv. *vesicatoria* 116 (P1) in non-surface-disinfested PI 235047 and Marengo leaves was 1.4×10^8 and 5.9×10^8 CFU/g, respectively. After surface disinfection, the population density in PI 235047 leaves was 6.4×10^7 CFU/g and in Marengo leaves was 6.5×10^7 CFU/g. These population densities did not differ significantly ($P = 0.05$).

DISCUSSION

Of the 170 accessions from eight different *Capsicum* species evaluated as potential sources of resistance to *X. campestris*

pv. *vesicatoria* P6 strains, only *C. pubescens* PI 235047 was resistant. None of the *C. pubescens* PI 235047 plants tested showed leaf spot symptoms 10 days after inoculation, while other accessions and Marengo plants were highly susceptible. Furthermore, elicitation of HR was observed in *C. pubescens* PI 235047 plants within 24 h after infiltration, but not in other PI lines or Marengo plants. This supports evidence from previous studies (13,15,16,20) that there is a strong correlation between the incompatible interaction of *X. campestris* pv. *vesicatoria* strains on pepper and HR elicitation, which can be used as a method for screening *Capsicum* accessions for resistance to the bacterial spot pathogen. In addition, these results indicate that *C. pubescens* PI 235047 plants carry a resistance (R) gene or genes that presumably correspond to a novel (unknown) avirulence gene or genes in *X. campestris* pv. *vesicatoria* P6 strains. Although genetic analyses of resistance in *C. pubescens* PI 235047 and avirulence in *X. campestris* pv. *vesicatoria* P6 strains have not been done, it is likely that the resistance is governed by a single gene, which we propose to designate *Bs4*, with the corresponding avirulence gene in the pathogen designated *avrBs4*. Our results, demonstrating a qualitative response to the pathogen and the presence of an HR in PI 235047 plants infiltrated with *X. campestris* pv. *vesicatoria* P6 strains, provide indirect evidence for single-gene resistance to bacterial spot in this accession. Furthermore, all three previously identified bacterial spot resistance genes in pepper, *Bs1*, *Bs2*, and *Bs3*, are single dominant genes (11,13,15,16,20). Genetic studies are needed to determine the inheritance of resistance to bacterial spot in this new resistance source.

Further evaluation of variation in virulence among *X. campestris* pv. *vesicatoria* pepper races showed that *C. pubescens* PI 235047 plants are not only incompatible with *X. campestris* pv. *vesicatoria* P6 strains, but also with P0, P4, and half of the P1 and P3 strains tested. The diversity among *X. campestris* pv. *vesicatoria* P1 and P3 strains provided evidence that the strains within *X. campestris* pv. *vesicatoria* pepper races may not be homogeneous in terms of host specificity and virulence. Although pathological diversity among strains of *X. campestris* pv. *vesicatoria* pepper races P0, P2, P4, P5, and P6 was not observed, this does not necessarily indicate that the strains within these races were homogeneous, because the number of strains tested was limited. In order to discriminate different pathogenic groups within *X. campestris* pv. *vesicatoria* races, *C. pubescens* PI 235047 was used as an additional pepper differential line. Therefore, *X. campestris* pv. *vesicatoria* strains were reclassified into races based on HR on pepper differential lines including PI 235047 (Table 3).

Table 2. (continued from preceding page)

Strain ^b	Race ^c	HR test ^d	DSV ^a	
			PI 235047	Marengo
Xcv-131b	P6	+	NT	NT
Xcv-787	P6	+	NT	NT
Xcv-788	P6	+	NT	NT
Xcv-789	P6	+	NT	NT
Xcv-790	P6	+	NT	NT
Xcv-791	P6	+	NT	NT
Xcv-792	P6	+	NT	NT
Xcv-793	P6	+	NT	NT
Xcv-69	P1	–	3.25	3.50
Xcv-114	P1	–	3.25	4.00
Xcv-115	P1	–	3.25	3.50
Xcv-116	P1	–	3.75	3.75
Xcv-117	P1	–	2.75	3.25
Xcv-66	P2	–	2.25	4.00
Xcv-89	P2	–	2.50	3.25
Xcv-90	P3	–	2.50	3.25
Xcv-91	P3	–	2.75	3.75
Xcv-110a	P3	–	3.25	3.75
Xcv-110b	P3	–	2.75	3.75
Xcv-110c	P3	–	2.25	3.25
Xcv-111	P3	–	3.00	3.75
Xcv-123a	P3	–	2.75	3.50
Xcv-123b	P3	–	3.25	2.75
Xcv-123c	P3	–	2.75	3.25
Xcv-123d	P3	–	2.25	2.75
Xcv-124a	P3	–	2.75	3.25
Xcv-124b	P3	–	3.00	3.00
Xcv-124c	P3	–	3.00	3.75
Xcv-124d	P3	–	3.25	3.25
Xcv-124e	P3	–	2.75	3.00
Xcv-125a	P3	–	2.75	4.00
Xcv-126a	P3	–	2.25	3.75
Xcv-126b	P3	–	2.75	4.00
Xcv-128a	P3	–	2.50	4.00
Xcv-128b	P3	–	3.00	3.50
Xcv-128c	P3	–	2.25	3.25
Xcv-128d	P3	–	2.50	3.50
Xcv-767	P3	–	NT	NT
V-205	P5	–	3.25	3.75
ANOVA ^e ($P = 0.05$)			LSD = 0.17	LSD = 0.20

Two new *X. campestris* pv. *vesicatoria* pepper races, designated *X. campestris* pv. *vesicatoria* P7 and P8, were identified by splitting from *X. campestris* pv. *vesicatoria* race P1 and P3, the strains that were compatible with PI 235047. Based on the new race classification of *X. campestris* pv. *vesicatoria* strains, *C. pubescens* PI 235047 plants are incompatible with strains of *X. campestris* pv. *vesicatoria* pepper races P0, P1, P3, P4, and P6. However, they are compatible with strains of *X. campestris* pv. *vesicatoria* pepper races P2, P5, P7, and P8.

The disease severity data showed that *C. pubescens* PI 235047 plants are significantly less susceptible than Marengo plants to *X. campestris* pv. *vesicatoria* strains that do not elicit an HR in PI 235047. These results suggest that *C. pubescens* PI 235047 plants may also provide a source of quantitative resistance to *X. campestris* pv. *vesicatoria* races P2, P5, P7, and P8.

Surface sterilization of inoculated pepper leaves caused at least a 10-fold reduction in average bacterial population densities, suggesting that *X. campestris* pv. *vesicatoria* applied during inoculation may survive on leaf surfaces for a long time. Therefore, the bacterial counts determined directly from dilution plating without surface disinfection may not represent the internal bacterial densities or be correlated with the disease severity values of the inoculated leaves.

The finding of a new resistance source in *C. pubescens* PI 235047 to *X. campestris* pv. *vesicatoria* has several implications. The gene or genes responsible for resistance to race P6 may be incorporated into commercial pepper cultivars to provide protection against an emerging problem. It may also be possible to transfer apparent quantitative resistance as well. Although crossing barriers between purple-flowered (*C. pubescens*) and white-flowered (including *C. annuum*, *C. baccatum*, *C. chinense*, and *C. frutescens*) pepper species are known, barriers between *C. pubescens* and *C. annuum* can be overcome by using bridge pepper species such as *C. cardenasii* or *C. eximium* (38). Furthermore, this new resistance source may allow molecular

characterization of additional resistance genes in *Capsicum* spp. and avirulence genes in *X. campestris* pv. *vesicatoria*.

ACKNOWLEDGMENTS

We thank L. V. Madden and P. E. Lipps for providing pre-submission reviews, and M. S. Krause for preparation of the figures.

LITERATURE CITED

- Adamson, W. C., and Grover, S. Jr. 1983. Inheritance of bacterial spot resistance in pepper. HortScience 18:905-906.
- Adaskaveg, J. E., and Hine, R. B. 1985. Copper tolerance and zinc sensitivity of Mexican strains of *Xanthomonas campestris* pv. *vesicatoria*, causal agent of bacterial spot of pepper. Plant Dis. 69:993-996.
- Bashan, Y., Diab, S., and Okon, Y. 1982. Survival of *Xanthomonas campestris* pv. *vesicatoria* in pepper seeds and roots in symptomless and dry leaves in non-host plants and in the soil. Plant Soil 68:161-170.
- Bashan, Y., and Okon, Y. 1986. Internal and external infections of fruits and seeds of pepper by *Xanthomonas campestris* pv. *vesicatoria*. Can. J. Bot. 64:2865-2871.
- Bashan, Y., Okon, Y., and Henis, Y. 1982. Long-term survival of *Pseudomonas syringae* pv. *tomato* and *Xanthomonas campestris* pv. *vesicatoria* in tomato and pepper seeds. Phytopathology 72:1143-1144.
- Bonas U., Schulte, R., Fenselau, S., Minsavage, G. V., Staskawicz, B. J., and Stall, R. E. 1991. Isolation of a gene cluster from *Xanthomonas campestris* pv. *vesicatoria* that determines pathogenicity and the hypersensitive response on pepper and tomato. Mol. Plant-Microbe Interact. 4:81-88.
- Bonas, U., Stall, R. E., and Staskawicz, B. J. 1989. Genetic and structural characterization of the avirulence gene *avrBs3* from *Xanthomonas campestris* pv. *vesicatoria*. Mol. Gen. Genet. 218:127-136.
- Bouzar, H., Jones, J. B., Stall, R. E., Hodge, N. C., Minsavage, G. V., Benedict, A. A., and Alvarez, A. M. 1994. Physiological, chemical, serological, and pathogenic analyses of a worldwide collection of *Xanthomonas campestris* pv. *vesicatoria* strains. Phytopathology 84:663-671.
- Canteros, B., Minsavage, G., Bonas U., Pring, D., and Stall, R. 1991. A gene from *Xanthomonas campestris* pv. *vesicatoria* that determines avirulence in tomato is related to *avrBs3*. Mol. Plant-Microbe Interact. 4:628-632.
- Conover, R. A., and Gerhold, N. R. 1981. Mixtures of copper and maneb or mancozeb for control of bacterial spot of tomato and their compatibility for control of fungus diseases. Proc. Fla. State Hortic. Soc. 94:154-156.

- Cook, A. A., and Guevara, Y. G. 1984. Hypersensitivity in *Capsicum chacoense* to race 1 of the bacterial spot pathogen of pepper. Plant Dis. 68:329-330.
- Cook, A. A., and Stall, R. E. 1963. Inheritance of resistance in pepper to bacterial spot. Phytopathology 53:1060-1062.
- Cook, A. A., and Stall, R. E. 1982. Distribution of races of *Xanthomonas vesicatoria* pathogenic on pepper. Plant Dis. 66:388-389.
- Cox, R. S. 1982. Control of bacterial spot of tomato in southern Florida. Plant Dis. 66:870.
- Hibberd, A. M., Bassett, M. J., and Stall, R. E. 1987. Allelism tests of three dominant genes for hypersensitive resistance to bacterial spot of pepper. Phytopathology 77:1304-1307.
- Hibberd, A. M., Stall, R. E., and Bassett, M. J. 1987. Different phenotypes associated with incompatible races and resistance genes in bacterial spot disease of pepper. Plant Dis. 71:1075-1078.
- Jones, J. B., Pohronezny, K. L., Stall, R. E., and Jones, J. P. 1986. Survival of *Xanthomonas campestris* pv. *vesicatoria* in Florida on tomato crop residue, weeds, seeds, and volunteer tomato plants. Phytopathology 76:430-434.
- Kearney, B., Ronald, P. C., Dahlbeck, D., and Staskawicz, B. J. 1988. Molecular basis for evasion of plant host defence in bacterial spot disease of pepper. Nature 332:541-543.
- Keen, N. T. 1990. Gene-for-gene complementarity in plant-pathogen interactions. Annu. Rev. Genet. 24:447-463.
- Kim, B.-S., and Hartmann, R. W. 1985. Inheritance of a gene (*Bs3*) conferring hypersensitive resistance to *Xanthomonas campestris* pv. *vesicatoria* in pepper (*Capsicum annuum*). Plant Dis. 69:233-235.
- Knoop, V., Staskawicz, B., and Bonas, U. 1991. Expression of avirulence gene *avrBs3* from *Xanthomonas campestris* pv. *vesicatoria* is not under the control of *hrp* genes and is independent of plant factors. J. Bacteriol. 173:7142-7150.
- Kousik, C. S., and Ritchie, D. F. 1995. Isolation of pepper races 4 and 5 of *Xanthomonas campestris* pv. *vesicatoria* from diseased peppers in southern U. S. fields. Plant Dis. 79:540.
- Kousik, C. S., and Ritchie, D. F. 1996. Evaluation of bell pepper varieties against races 1-4 of the bacterial spot pathogen, 1995. Biol. Cult. Tests Plant Dis. Control 11:97.
- Kousik, C. S., and Ritchie, D. F. 1996. Disease potential of pepper bacterial spot pathogen races that overcome the *Bs2* gene for resistance. Phytopathology 86:1336-1343.
- Kousik, C. S., and Ritchie, D. F. 1996. Mixed genotypes combined with copper sprays to manage bacterial spot of bell peppers. Phytopathology 86:502-508.
- Lelliott, R. A., and Stead, D. E. 1987. Methods for the diagnosis of bacterial diseases of plants. Blackwell Scientific Publications, Oxford.
- Marco, G. M., and Stall, R. E. 1983. Control of bacterial spot of pepper initiated by strains of *Xanthomonas campestris* pv. *vesicatoria* that differ in sensitivity to copper. Plant Dis. 67:779-781.
- Minsavage, G. V., Dahlbeck, D., Whalen, M. C., Kearney, B., Bonas, U., Staskawicz, B. J., and Stall, R. E. 1990. Gene for gene relationships specifying disease resistance in *Xanthomonas campestris* pv. *vesicatoria*-pepper interactions. Mol. Plant-Microbe Interact. 3:41-47.
- Ritchie, D. F., and Dittapongpich, V. 1991. Copper- and streptomycin-resistant strains and host differentiated races of *Xanthomonas*

Table 3. Proposed race classification of *Xanthomonas campestris* pv. *vesicatoria* pepper strains based on hypersensitive reaction (HR) on pepper differential lines and *Capsicum pubescens* PI 235047

Races	Pepper differential lines				
	ECW	ECW-10R <i>Bs1</i>	ECW-20R <i>Bs2</i>	ECW-30R <i>Bs3</i>	PI 235047 <i>Bs4</i> ^a
P0	— ^b	HR	HR	HR	HR
P1	—	—	HR	HR	HR
P2	—	HR	HR	—	—
P3	—	—	HR	—	HR
P4	—	—	—	HR	HR
P5	—	HR	—	—	—
P6	—	—	—	—	HR
P7	—	—	HR	HR	—
P8	—	—	HR	—	—

^a Proposed designation.

^b Compatible interaction.

- campestris* pv. *vesicatoria* in North Carolina. Plant Dis. 75:733-736.
30. Ronald, P. C., and Staskawicz, B. J. 1988. The avirulence gene *avrBs1* from *Xanthomonas campestris* pv. *vesicatoria* encodes a 50-kd protein. Mol. Plant-Microbe Interact. 1:191-198.
 31. Sahin, F. 1997. Detection, identification and characterization of strains of *Xanthomonas campestris* pv. *vesicatoria* by traditional and molecular methods, and resistance in *Capsicum* species to *Xanthomonas campestris* pv. *vesicatoria* pepper race 6. Ph.D. thesis. The Ohio State University, Columbus.
 32. Sahin, F., and Miller, S. A. 1995. First report of pepper race 6 of *Xanthomonas campestris* pv. *vesicatoria*, causal agent of bacterial spot of pepper. Plant Dis. 79:1188.
 33. Sahin, F., and Miller, S. A. 1995. Characterization of Ohio strains of *Xanthomonas campestris* pv. *vesicatoria*, causal agent of bacterial spot of pepper. Plant Dis. 80:773-778.
 34. Schulte, R., and Bonas, U. 1992. Expression of the *Xanthomonas campestris* pv. *vesicatoria* *hrp* gene cluster, which determines pathogenicity and hypersensitivity on pepper and tomato, is plant inducible. J. Bacteriol. 174:815-823.
 35. Sijam, K., Chang, C. J., and Gitaitis, R. D. 1991. An agar medium for the isolation and identification of *Xanthomonas campestris* pv. *vesicatoria* from seed. Phytopathology 81:831-834.
 36. Stall, R. E., and Thayer, P. L. 1962. Streptomycin resistance of the bacterial spot pathogen and control with streptomycin. Plant Dis. Rep. 46:389-392.
 37. Vauterin, L., Hoste, B., Kersters, K., and Swings, J. 1995. Reclassification of *Xanthomonas*. Int. J. Syst. Bacteriol. 45:472-489.
 38. Zijlstra, S., Purimahua, C., and Lindhout, P. 1991. Pollen tube growth in interspecific crosses between *Capsicum* species. Hort-Science 26:585-586.