AFLP Comparisons Among *Claviceps africana* Isolates from the United States, Mexico, Africa, Australia, India, and Japan

Paul W. Tooley, Erin D. Goley, and Marie M. Carras, USDA-ARS, Foreign Disease-Weed Science Research Unit, 1301 Ditto Ave., Ft. Detrick, MD 21702-5023; and Nichole R. O'Neill, USDA-ARS, Molecular Plant Pathology Laboratory, Beltsville, MD 20705

ABSTRACT

Tooley, P. W., Goley, E. D., Carras, M. M., and O'Neill, N. R. 2002. AFLP comparisons among *Claviceps africana* isolates from the United States, Mexico, Africa, Australia, India, and Japan. Plant Dis. 86:1247-1252.

Eighty-seven isolates of the sorghum ergot pathogen, *Claviceps africana*, from diverse geographic locations were analyzed using four different amplified fragment length polymorphism (AFLP) primer combinations to determine genetic relationships among isolates. Most isolates showed unique AFLP haplotypes, indicating that substantial genetic variation is present within *C. africana* populations. Two major groupings of isolates were observable, with ca. 70% similarity between the two groups. One group consisted of Australian, Indian, and Japanese isolates and the other of U.S., Mexican, and African isolates. In spite of overall high levels of genetic diversity observed in *C. africana*, isolates within the two major groups were between 75 and 100% similar. The observed associations of *C. africana* isolates from worldwide sources could be the result of intercontinental trade and/or movement of seed. The data indicate that Africa was the likely source of *C. africana* that has become established in the Americas since 1996. Analysis of additional isolates in future studies will reveal whether these groupings are being maintained or whether population subdivision or reshuffling may occur.

Additional keywords: population genetics, *Sphacelia sorghi*, sugary disease

Sorghum ergot first appeared in the Western Hemisphere in Brazil in 1995 (1,2,21), and by the end of the 1997 growing season it had spread to Mexico and the United States, reaching as far north as the state of Nebraska (11). Since then, sporadic outbreaks have occurred in the United States, with the disease spreading to additional states such as Georgia. While major devastating epidemics of sorghum ergot have generally not occurred in the United States since 1997, small outbreaks have occurred. Plant pathologists in sorghum-growing states have remained vigilant in monitoring the occurrence of ergot on volunteer sorghum and alternate hosts such as Johnson grass (*Sorghum halepense*). It is now widely believed that, in light of existing levels of inoculum observed on various host plants throughout states in which ergot is known to exist, the occurrence of environmental conditions that favor infection and disease development at the

right time in the sorghum-growing season could result in damaging ergot epidemics.

Sorghum ergot is caused by three distinct *Claviceps* species (*C. africana*, *C. sorghi*, and *C. sorghicola*) which are differentiated based on morphology and other characters (7,26). Molecular comparisons using sequences of β -tubulin and EF-1 α gene introns (24) have helped confirm the identity of the sorghum ergot pathogen present in the United States as *C. africana*, as distinct from other sorghum-infecting *Claviceps* species such as *C. sorghi* and *C. sorghicola*. In Japan (26) and India (2), two of the three sorghum-infecting *Claviceps* species coexist and may cause sorghum ergot in the same growing season.

Whether *Claviceps* species affecting sorghum can cross-hybridize in nature is not known, and sclerotia that give rise to the sexual stage of the pathogen have not been observed to germinate in nature. Currently, only a single species (*C. africana*) has been found in the United States causing sorghum ergot. However, global migrations such as the one that brought sorghum ergot to the Western Hemisphere in 1995 doubtless continue to occur through movement of sorghum seed and/or direct movement of conidia in wind currents and/or associated with insect or avian vectors. This could make it possible for additional species to enter and become established in the United States at any time.

In light of the ever-present potential for pathogen migration to occur and alter the composition of *Claviceps* species and genotypes in a particular region, it is important to document the genetic makeup of existing populations using the most reliable methods available. In particular, knowledge of levels of genotypic diversity is important in order to determine whether diversity is increasing due to influx of additional genotypes. With the potential for additional genetic diversity to be added to the U.S. population through future pathogen migrations, the likelihood of occurrence of new genotypes with enhanced pathogenic potential is increased. Sorghum breeders in states such as Texas and Kansas are attempting to incorporate ergot resistance into their breeding lines through screening and selection using existing pathogen genotypes. If new genotypes or pathogen species arrive in the United States, the more diverse pathogen population may render such resistance ineffective.

Knowledge of levels of pathogen diversity found elsewhere in the world can allow workers to better evaluate the epiphytotic potential of U.S. populations, as well as identify regions of the world where more stringent field screening of sorghum genotypes may be carried out. In a previous study, we used random amplified microsatellite (RAM) and amplified fragment length polymorphism (AFLP) analyses to characterize a small number of *C. africana* isolates from several continents; 20 isolates were analyzed using AFLPs with primers *MseI*+C and *EcoRI*+AG, and high levels of similarity were observed among isolates from the United States, Puerto Rico, and South Africa (25).

While the previous work allowed an initial look at genotypic diversity within *C. africana*, too few isolates were examined to give a clear overall picture of population relatedness on a global scale. The AFLP method detected intraspecific variation more readily than RAM (25) and was therefore the method of choice for this study. We have extended AFLP analysis to a larger group of isolates from worldwide sources and used three additional primer combinations to provide a deeper understanding of the genetic relationships among isolates from different geographic regions. The relationships and groupings we have identified among ergot strains from diverse continents provide basic knowledge about

Corresponding author: P. W. Tooley
E-mail: tooley@ncifcrf.gov

Accepted for publication 28 June 2002.

Publication no. D-2002-0821-01R

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 2002.

intercontinental populations of *C. africana* and will provide baseline data with which future population data may be compared. This will allow analysis of future migrations of *C. africana* and assessment of the degree to which levels of genetic variation may be shifting.

MATERIALS AND METHODS

Single conidial *C. africana* isolates were produced from ergot-infected sorghum provided by colleagues (Table 1). Cultures were established by diluting honeydew from infected sorghum seed heads ca. 100-fold in sterile distilled water, then streaking them onto plates of 2% water agar containing 100 µg of ampicillin and 100 µg of streptomycin sulfate per ml. Plates were

incubated at 22°C in the dark, and individual germinated conidia were transferred to plates of potato dextrose agar (PDA) where they grew into colonies. When samples had low conidial viability, a conidial suspension was made in sterile distilled water and used to inoculate flowering heads of susceptible sorghum plants (male sterile line aTX 623 or aTX 2752, obtained from G. Odvody and D. Frederiksen, Texas A&M University) grown in the USDA-ARS containment greenhouse complex at Ft. Detrick, MD. Newly flowering sorghum heads were immersed in a conidial suspension made by combining several infected sorghum florets containing fresh or dried sphacelia and/or honeydew with 150 ml of distilled water in a 1-qt plastic zipper-seal

freezer bag. After gentle agitation, numerous conidia were released from the infected florets, making a dense conidial suspension. Following immersion in the suspension, heads were bagged with 1-qt plastic zipper-seal freezer bags for 24 to 48 h to preserve high relative humidity. Symptoms of infection and initial honeydew production could be observed on inoculated heads 5 to 7 days after removal of the bags. Fresh honeydew produced on inoculated heads was then used to establish single-conidial cultures as described above.

DNA was extracted from 5 to 10 mg of lyophilized *C. africana* mycelium grown at 22°C in the dark in yeast-malt extract (YM) broth (Difco Laboratories, Detroit,

Table 1. Sources of the 87 *Claviceps africana* isolates used in this study^a

Species and isolate ^b	Geographic source	Year of isolation	Species and isolate	Geographic source	Year of isolation
Cls-1	Patancheru, India	1997	Cla-83	Miyakonojo, Japan	1988
Cls-2	Akola, India	1997	Cla-88	Miyakonojo, Japan	1990
Cls-3	Dharwar, India	1997	Cla-89	Lusaka, Zambia ^c	1999
Cls-4	Patancheru, India	1997	Cla-91	Roma, Queensland, Australia	1999
Cla-1	Norwin, Queensland, Australia	1996	Cla-92	Oakey, Queensland, Australia	1999
Cla-2	Norwin, Queensland, Australia	1996	Cla-93	Queensland, Australia	1997
Cla-3	Monto, Queensland, Australia	1997	Cla-94	Capella, Queensland, Australia	1999
Cla-4	College Station, Texas, USA	1997	Cla-95	Warwick, Queensland, Australia	1999
Cla-5	College Station, Texas, USA	1997	Cla-96	Roma, Queensland, Australia	1999
Cla-7	Isabela, Puerto Rico	1997	Cla-97	Emerald, Queensland, Australia	1999
Cla-8	Isabela, Puerto Rico	1997	Cla-99	Sinton, Texas, USA	1999
Cla-9	Weslaco, Texas, USA	1998	Cla-100	Sinton, Texas, USA	1999
Cla-10	Weslaco, Texas, USA	1998	Cla-101	Sinton, Texas, USA	1999
Cla-12	Isabela, Puerto Rico	1998	Cla-102	Kiowa County, Kansas, USA	1997
Cla-14	Isabela, Puerto Rico	1998	Cla-103	Kiowa County, Kansas, USA	1997
Cla-17	Isabela, Puerto Rico	1998	Cla-104	Kiowa County, Kansas, USA	1997
Cla-25	Isabela, Puerto Rico	1998	Cla-105	Allen County, Kansas, USA	1997
Cla-32	Guayanilla, Puerto Rico	1998	Cla-107	Allen County, Kansas, USA	1997
Cla-34	Guayanilla, Puerto Rico	1998	Cla-111	Isabela, Puerto Rico	1999
Cla-38	Virginia, Nebraska, USA	1997	Cla-112	Isabela, Puerto Rico	1999
Cla-42	Virginia, Nebraska, USA	1997	Cla-114	Nazareth, Texas, USA	1999
Cla-46	Grantham, Queensland, Australia	1997	Cla-116	Nazareth, Texas, USA	1999
Cla-47	Mutdapilly, Queensland, Australia	1997	Cla-117	Lusaka, Zambia ^c	1999
Cla-48	Gindie, Queensland, Australia	1997	Cla-120	Harare, Zimbabwe	1999
Cla-49	Hale Center, Texas, USA	1998	Cla-121	Harare, Zimbabwe	1999
Cla-51	Vega, Texas, USA	1998	Cla-122	Bulawayo, Zimbabwe	1999
Cla-52	Corpus Christi, Texas, USA	1998	Cla-123	Lusaka, Zambia ^c	1999
Cla-53	Corpus Christi, Texas, USA	1998	Cla-124	Lusaka, Zambia ^c	1999
Cla-54	Corpus Christi, Texas, USA	1998	Cla-128	Panmure, Zimbabwe	1999
Cla-56	Potchefstroom, South Africa	1998	Cla-129	Ft. Pierce, Florida, USA	1999
Cla-59	Ennis, Texas, USA	1998	Cla-130	Ft. Pierce, Florida, USA	1999
Cla-60	Ennis, Texas, USA	1998	Cla-133	Ft. Pierce, Florida, USA	1999
Cla-61	Ennis, Texas, USA	1998	Cla-134	Ft. Pierce, Florida, USA	1999
Cla-62	Comonfort, Mexico	1998	Cla-135	Nazareth, Texas, USA	1999
Cla-63	Comonfort, Mexico	1998	Cla-137	Nazareth, Texas, USA	1999
Cla-66	Celaya, Mexico	1998	Cla-138	Seek, Kansas	1999
Cla-68	Comonfort, Mexico	1998	Cla-142	Keese, Kansas	1999
Cla-69	Cortazu, Mexico	1998	Cla-143	Kiowa County, Kansas, USA	1997
Cla-70	Tifton, Georgia, USA	1998	Cla-IN-1	Patancheru, India	1999
Cla-74	Tifton, Georgia, USA	1998	Cla-IN-2	Maachinenapally, India	1999
Cla-77	Tifton, Georgia, USA	1998	Cla-IN-7	Chakalisherupally, India	1999
Cla-78	Ennis, Texas, USA	1998	Cla-IN-8	Injamura, India	1999
Cla-79	Ennis, Texas, USA	1998	Cla-IN-29	Mana, India	1999
Cla-80	Miyakonojo, Japan	1988			

^a *Sorghum bicolor* was the host for all *C. africana* isolates listed except for Cla-1 and Cla-2, whose host was forage sorghum (*Sorghum* sp.), and Cla-68, whose host was Johnson grass (*Sorghum halepense*).

^b Pure cultures of isolates Cls-1 to Cls-4 and Cla-IN-1, -2, -7, -8, and -29 were provided by R. Bandyopadhyay, ICRISAT, Patancheru, India, and pure cultures of isolates Cla-2 and Cla-3 were provided by Malcolm Ryley, Queensland Dept. of Primary Industries, Australia. Pure cultures of Cla-91 through Cla-97 were obtained from Birte Komolung, University of Queensland, Australia. All other cultures were established from single conidia in our containment laboratory as described in Materials and Methods.

^c Sample from which isolate was obtained was collected at the Golden Valley Research Farm, 25 miles north of Lusaka, Zambia, by G. Odvody and N. McLaren.

MI) using a modified CTAB DNA extraction protocol (23). AFLPs were generated with the AFLP Analysis System II (Invitrogen, Carlsbad, CA) using a method previously described by Tooley et al. (25). Primer sets used for analysis in the selective amplification included *MseI*+C combined with *EcoRI*+AG, *EcoRI*+AC, *EcoRI*+TA, and *EcoRI*+TT. Data were combined for the analysis. Initially, two additional primer combinations were used (*MseI*+C combined with *EcoRI*+AA and *EcoRI*+AT), but observed levels of polymorphism among isolates were very low and the data were not included in the analysis.

AFLP profiles were manually scored for presence or absence of bands and entered into a binary matrix. Only polymorphisms that could be scored unequivocally for presence or absence of bands were included in the analyses; faint bands were never scored as absent. Bands were assumed to be independent. Binary matrices were analyzed to obtain simple matching coefficients among the isolates using NTSYS-pc, version 2.0 (Exeter Software, Setauken, NY). Simple matching coefficients were clustered to generate similarity trees using SAHN clustering using the unweighted pair-group method with arithmetic average (UP-GMA) algorithm in NTSYS-pc. *Claviceps purpurea* was used as an outgroup species, since it represents on biological grounds a related species that is less closely related to each of the taxa under consideration (isolates of *C. africana*) than any are to one other (24,25).

Fig. 1. Section of autoradiograph (primer *EcoRI*+AG) showing amplified fragment length polymorphism (AFLP) differences between U.S. and Indian isolates of *Claviceps africana*. Arrows represent bands present in U.S. isolates and absent in Indian isolates (111 bp) or vice versa (120 bp, 83 bp).

Measures of genetic variability for isolates within and among regions were calculated using POPGENE (34) and tools for population genetic analysis (TFPGA) (17), which are Windows-based freeware programs for the analysis of population genetic data. Genetic diversity within each *C. africana* population was assessed by using the above programs to calculate the observed and effective number of alleles (13), Nei's gene diversity (18), the Shannon information index (22), the estimated (unbiased) average heterozygosity (18), and the proportion of polymorphic loci (9).

Analysis of molecular variance (AMOVA) was performed as described by Excoffier et al. (5) using the program Genetic Studio 2.0 (4) and using the methods of Weir and Cockerham (30). AMOVA produces estimates of variance components and *F* statistic analogs (Φ statistics) that reflect genetic diversity at different levels of hierarchical subdivision. It was developed to estimate and evaluate variance components of genetic markers within and between populations, and has been used previously to assess genetic relatedness in plant-pathogenic fungi (21,29,31) as well as in plants (10). The fixation index (or *F* statistic analog) calculated in AMOVA is equivalent to the reduction in heterozygosity expected with random mating at any one level of population hierarchy relative to another more inclusive level of the hierarchy (33). The genetic distance between any two populations is represented by its Φ_{st} value, which equals the proportion of the total variance that is partitioned between the two populations (5).

RESULTS

A total of 130 loci were scored for the four AFLP primer combinations used; 35 loci were scored for primer set *MseI*+C/*EcoRI*+AG, 30 loci for *MseI*+C/*EcoRI*+AC, 36 loci for *MseI*+C/*EcoRI*+TA, and 29 loci for *MseI*+C/*EcoRI*+TT. Figure 1 shows AFLP results for primer set *MseI*+C/*EcoRI*+AG, showing some prominent differences between U.S. and Indian isolates. Primer sets varied in percentage of polymorphic loci, gene diversity, and Shannon Index, with primer TT the lowest (Table 2). Primer TA showed the highest gene diversity and percentage of polymorphic loci.

The total number of AFLP bands observed over all four primer combinations varied little among isolate groups, ranging from 98 bands for African isolates to 105 bands for U.S. isolates (Table 3). However, nearly every isolate examined was unique in terms of its AFLP haplotype, indicating a high degree of overall genetic diversity within *C. africana* (Table 3). Genetic diversity in terms of percent polymorphic loci was lowest for Japanese isolates (10%) and highest for U.S. (45.4%) and African (43.8%) isolates (Table 3).

When a dendrogram was constructed from the combined AFLP data from all four primer combinations, two major groupings were observable. One group consisted of Australian, Indian, and Japanese isolates, and the other consisted of U.S., Mexican, and African isolates, with ca. 70% similarity between the groups (Fig. 2). Isolates within groups ranged from ca. 75 to 100% similar. Outgroup *C.*

Table 2. Gene diversity measures for four amplified fragment length polymorphism (AFLP) primer pair combinations used to analyze *Claviceps africana* from worldwide sources

Primer pair ^a	na ^b	ne ^c	h ^d	I ^e	ppl (%) ^f
AG	1.7222	1.3383	0.1983	0.3013	72.2
AC	1.7667	1.3674	0.2090	0.3139	76.7
TA	1.8611	1.5076	0.2872	0.4237	86.1
TT	1.4828	1.2988	0.1726	0.2537	48.3
Combined	1.3130	1.2099	0.1180	0.1740	71.8

^a AFLP primer pairs were *MseI*+C paired with, respectively, *EcoRI*+AG, *EcoRI*+AC, *EcoRI*+TA, and *EcoRI*+TT.

^b Observed number of alleles.

^c Effective number of alleles, which is the reciprocal of homozygosity (13).

^d Nei's gene diversity (18).

^e Shannon's information index as a measure of gene diversity (22).

^f Percentage of polymorphic loci (34).

Table 3. Estimates of genetic diversity within groups of *Claviceps africana* from different geographic regions^a

Geographic source	Isolates	Unique haplotypes	Total AFLP bands	Polymorphic loci (%)
India	9	8	101	31.5
Australia	13	12	100	38.5
Japan	3	3	99	10.0
U.S.	48	47	105	45.4
Africa	9	9	98	43.8
Mexico	5	5	102	35.4

^a Data are combined from four amplified fragment length polymorphism (AFLP) primer pair combinations representing 130 total loci.

purpurea isolate Clp-1 (ATCC 34699, Montana, USA, host *Hordeum vulgare*) showed ca. 30% similarity with *C. africana* isolates (data not shown). Table 4 and

Figure 3 show genetic similarity and Nei's genetic distance among six geographic collections of *C. africana*. Distances ranged from ca. 0.04 for the most closely

related groups to ca. 0.20 for the most distantly related groups. U.S. and African isolates were the most closely related, followed by Indian and Australian isolates. Japanese and African isolates were the most distantly related (Table 4, Fig. 3).

Hierarchical AMOVA was conducted for the combined AFLP data based on groups (United States/Africa/Mexico versus India/Australia/Japan) and collections within groups (United States, Africa, and Mexico within the first group and India, Australia, and Japan within the second group) (Table 5) identified from dendrograms (Fig. 2). Most of the AFLP haplotype genetic diversity (50.2%) was found within each collection, but a large amount (38.5%) also separated groups. The differences among collections within groups are relatively small (11.4%). The Φ statistic also reflected these trends (Table 5).

Pairwise comparisons of Φ_{st} values between collections of *C. africana* from six different regions (Table 6) show that the six populations were all significantly different from one another (at the 5% level), with the exception of Africa-Mexico and United States-Mexico. United States-Africa were different from one another, even though their genetic differentiation was low ($\Phi_{st} = 0.137$).

DISCUSSION

We used AFLP markers to analyze the genetic diversity among 87 isolates of the sorghum ergot pathogen, *C. africana*, from global sources. We found a high level of genetic diversity in all groups analyzed, with almost as many unique AFLP genotypes as isolates. Most of the AFLP haplotype genetic diversity was found within collections made in individual countries, but a large amount also separated the major geographical groups (United States/Africa/Mexico versus India/Australia/Japan). The differences among collections (countries) within the major geographical groups are relatively small. Wright (33) suggested that fixation indices (analogous to the Φ values in Table 5) ranging from 0.15 to 0.25 indicate great genetic differentiation, while values above 0.25 indicate very great genetic differentiation. Thus, we consider that the *C. africana* we sampled and analyzed shows very high levels of genetic differentiation.

In terms of AFLP analysis, we observed major associations of Indian, Australian, and Japanese isolates, and U.S., Mexican, and African isolates. Major groupings shared ca. 70% similarity, while within groups, isolates ranged from 75 to 100% similar. These associations support the results of other workers (20), who found similar associations of isolates based on random amplified polymorphic DNA (RAPD) analysis. They also support the results of our previous study (25), which examined AFLP variation in a smaller geographical representation of isolates.

Fig. 2. Dendrogram showing similarities among isolates of *Claviceps africana* based on amplified fragment length polymorphism (AFLP) analysis (data from four primer combinations combined) using simple matching coefficients. The tree was generated using the SAHN clustering program with the unweighted pair-group method with arithmetic average (UPGMA) algorithm in NTSYS-pc, version 2.0.

A limitation of our study was the low number of isolates collected from Japan and Mexico. Numbers collected from these locations were too low to have high confidence in diversity measures, pointing to the need for analysis of additional isolates from these locations in the future.

Langdon (16) believed that ecological and physiological adaptation were key processes involved in differentiation of *Claviceps* species. Genetic mechanisms which could explain such diversity include simple mutation, mitotic recombination, heterokaryosis, parasexual and/or sexual recombination (6), as well as more recently recognized genetic mechanisms such as chromosomal polymorphism, the presence of accessory chromosomes, cytoplasmically transmitted elements, and transposons (14). For the most studied *Claviceps* species, *C. purpurea*, a high degree of heterokaryosis observed among field isolates was theorized by Tudzynski (27) to be due to a high mutation rate. Also in *C. purpurea*, parasexual processes have been induced (3) and mitochondrial plasmids observed (28). White et al. (32) theorized that *C. purpurea* would have ample opportunity for genetic exchange because of the presence of the "honeydew stage" in which conidia are exposed to insects that visit florets, allowing mixing of sphaerial conidia from different populations and subsequent parasexual recombination. Similar mechanisms could occur with *C. africana*, as it also contains a honeydew stage.

Greater knowledge of the role of the sexual cycle, and whether it occurs in areas of recent introduction such as the United States, could shed light on the pathogen's propensity to generate novel genetic variation, which may complicate the deployment of effective long-term control measures. To date, sclerotia, which give rise to the sexual stage of the fungus, have not been observed in the United States, and it has been theorized that U.S. environmental conditions do not favor sclerotial formation. However, even with no evidence of the presence of an active sexual stage, *C. africana* genotypes appear to be highly diverse, as evidenced by the high numbers of unique AFLP genotypes observed in all populations. These results seem to agree with those observed for 29 *C. purpurea* isolates assayed using RAPD markers, in which high numbers of unique RAPD profiles were obtained (12).

The practical implications of our AFLP results include the finding of substantial genetic diversity within U.S. *C. africana* populations, facilitating the development of races or strains in the future. The diversity is surprising, given the very recent introduction of *C. africana* into the United States, and indicates that genetic mechanisms exist in *C. africana* to generate substantial variation in the absence of an active sexual stage.

The geographic associations of *C. africana* isolates observed from AFLP and RAPD analysis are reasonable in terms of the seed trade among the various countries

and continents. Australia, India, and Japan are linked as trading partners, as are the United States and Mexico. Based on RAPD analysis, Pažoutová (20) suggested

Table 4. Nei's unbiased measures of genetic identity and genetic distance^a for *Claviceps africana* amplified fragment length polymorphism (AFLP) data based on four primer combinations

Population	Population					
	India	Australia	U.S.	Africa	Mexico	Japan
India	...	0.9654	0.8242	0.8228	0.8259	0.9146
Australia	0.0352	...	0.8478	0.8640	0.8327	0.8556
U.S.	0.1934	0.1651	...	0.9666	0.9619	0.7906
Africa	0.1951	0.1461	0.0340	...	0.9187	0.7809
Mexico	0.1913	0.1830	0.0388	0.0848	...	0.8214
Japan	0.0893	0.1560	0.2349	0.2474	0.1967	...

^a Nei's genetic identity (above diagonal) and genetic distance (below diagonal) (18).

Fig. 3. Graphical representation of genetic distance relationships among six collections of *Claviceps africana* isolates from different geographic regions.

Table 5. Analysis of molecular variance (AMOVA) table for groups and collections of *Claviceps africana* from different geographic regions

	df	SS ^a	MS ^b	σ^c	% ^d	Φ^e	P value ^f
Among groups	2	226.417	113.209	4.957	38.5	0.385	0.010
Among coll. (groups)	4	81.064	20.266	1.462	11.4	0.185	0.010
Within collections	81	523.110	6.46	6.458	50.2	0.498	0.010
Total	87	830.591		12.877			

^a Sum of squares.

^b Mean square.

^c Variance component.

^d Percentage of total haplotype amplified fragment length polymorphism (AFLP) diversity present among groups, among populations within groups, and within populations.

^e Φ statistic as described by Excoffier et al. (5).

^f Probability of having a more extreme Φ statistic than the observed values by chance alone.

Table 6. Pairwise Φ_{st} values^a between six collections of *Claviceps africana* (above the diagonal) and their probability P values^b (below the diagonal)

Collection	Africa	Mexico	U.S.	Japan	Australia	India
Africa	0	0.147	0.137	0.536	0.377	0.466
Mexico	0.091	0	0.114	0.499	0.395	0.456
U.S.	<0.001	0.061	0	0.580	0.467	0.521
Japan	<0.001	0.010	<0.001	0	0.447	0.341
Australia	<0.001	<0.001	<0.001	<0.001	0	0.156
India	<0.001	<0.001	<0.001	<0.001	<0.001	0

^a F statistic analog (5).

^b Probability that a random distance calculated by analysis of molecular variance (AMOVA) with 1,000 permutations is greater than the observed distance, Φ_{st} .

that India or a Southeast Asian country such as Thailand may have been the likely origin of ergot introduction into Australia and also suggested that Africa is the source of *C. africana* introduction into the Americas. Our data from AFLP analysis lends weight to the hypothesis that Africa may have been the source of *C. africana* that reached South America in 1996 and the United States in 1997 via the seed trade or other mechanism.

The stability of isolate groupings we observed from AFLP analysis is not known. Recent evidence from studies of Australian *C. africana* using radiolabeled amplified fragment (RAF) analysis indicates that genetic diversity may be increasing in some populations since the collection of our isolates. Komolongo et al. (15) showed a higher level of recent genetic diversity than found in original populations. Perhaps movement of the pathogen from other regions either directly or via the sorghum seed trade is an ongoing process that has contributed to an increase in diversity in Australian *C. africana* populations.

Further analysis of greater numbers of *C. africana* isolates from diverse regions including Africa, India, Australia, South and Central America, and the United States will help determine the potential source of sorghum ergot in the United States and provide more data for assessing levels of genotypic diversity in this rapidly emerging pathogen. Greater sampling of isolates from Southeast Asian countries would be desirable to determine their similarity with Indian and Australian isolates. More detailed analysis of populations within India and Africa would also be desirable, as these are considered the regions in which *C. africana* has evolved for the longest period of time. Finally, other molecular tools such as microsatellites (19), dispersed repetitive DNA sequences (8), and/or transposon analysis (29) could be used to validate findings from AFLP analysis.

ACKNOWLEDGMENTS

We thank the colleagues who provided infected sorghum or pure cultures of *C. africana* for these studies, including Ranajit Bandyopadhyay, Sukumar Chakroborty, Larry Claflin, Jeffrey Dahlberg, Richard Frederiksen, Tom Isakeit, Stanley Jensen, Birte Komolongo, Neal McClaren, Gary Odvody, Malcolm Ryley, and Takao Tsukiboshi. Thanks also to S. Pažoutová for critical reading of the manuscript and many helpful suggestions.

LITERATURE CITED

1. Bandyopadhyay, R., Frederickson, D. E., McLaren, N. W., and Odvody, G. N. 1996.

- Ergot-A global threat to sorghum. *Int. Sorghum Millet Newsl.* 37:1-32.
2. Bandyopadhyay, R., Frederickson, D. E., McLaren, N. W., Odvody, G. N., and Ryley, M. J. 1998. Ergot: A new disease threat to sorghum in the Americas and Australia. *Plant Dis.* 82:356-367.
3. Brauer, K. L., and Robbers, J. E. 1987. Induced parasexual processes in *Claviceps* sp. strain SD58. *Appl. Environ. Microbiol.* 53:70-73.
4. Dyer, R. 1998. Genetic Studio 2.0. Shareware. Rodney Dyer, Dept. of Botany, Iowa State University, Ames.
5. Excoffier, L., Smouse, P. E., and Quattro, J. M. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics* 131:479-491.
6. Fincham, J. R. S., Day, P. R., and Radford, A. 1979. *Fungal Genetics*. University of California, Los Angeles.
7. Frederickson, D. E., Mantle, P. G., and De-Milliano, W. A. J. 1991. *Claviceps africana* sp. nov.; the distinctive ergot pathogen of sorghum in Africa. *Mycol. Res.* 95:1101-1107.
8. Hamer, J. E., Farrall, L., Orbach, M. J., Valent, B., and Chumley, F. G. 1989. Host species-specific conservation of a family of repeated DNA sequences in the genome of a fungal plant pathogen. *Proc. Natl. Acad. Sci. USA* 86:9981-9985.
9. Hartl, D. L., and Clark, A. G. 1997. *Principles of Population Genetics*. Sinauer Associates Inc., Sunderland, MA.
10. Huff, D. R., Peakall, R., and Smouse, P. E. 1993. RAPD variation within and among natural populations of outcrossing buffalograss [*Buchloe dactyoloides* (Nutt.) Engelm.]. *Theor. Appl. Genet.* 86:927-934.
11. Isakeit, T., Odvody, G. N., and Shelby, R. A. 1998. First report of sorghum ergot caused by *Claviceps africana* in the United States. *Plant Dis.* 82:592.
12. Jungehülsing, U., and Tudzynski, P. 1997. Analysis of genetic diversity in *Claviceps purpurea* by RAPD. *Mycol. Res.* 101:1-6.
13. Kimura, M., and Crow, J. F. 1964. The number of alleles that can be maintained in a finite population. *Genetics* 49:725-738.
14. Kistler, H. C., and Miao, V. P. W. 1992. New modes of genetic change in filamentous fungi. *Annu. Rev. Phytopathol.* 30:131-152.
15. Komolongo, B., Chakroborty, S., Ryley, M., and Yates, D. 2001. Genetic diversity of *Claviceps africana* in Australia. (Abstr.) *Phytopathology* 91:S50.
16. Langdon, R. F. N. 1954. The origin and differentiation of *Claviceps* species. *Trans. Br. Mycol. Soc.* 47:205-213.
17. Miller, M. P. 1997. Tools for population genetic analyses (TFPGA) 1.3: A Windows program for the analysis of allozyme and molecular population genetic data. Computer software. Mark P. Miller, Dept. of Fisheries and Wildlife, Utah State University, Logan.
18. Nei, M. 1978. Estimation of average heterozygosity and genetic distance from a small

- number of individuals. *Genetics* 89:583-590.
19. Neu, C., Kaemmer, D., Kahl, G., Fischer, D., and Weising, K. 1999. Polymorphic microsatellite markers for the banana pathogen *Mycosphaerella fijiensis*. *Mol. Ecol.* 8:513-525.
20. Pažoutová, S., Bandyopadhyay, R., Frederickson, D. E., Mantle, P. G., and Frederiksen, R. A. 2000. Relations among sorghum ergot isolates from the Americas, Africa, India, and Australia. *Plant Dis.* 84:437-442.
21. Pimentel, G., Peever, T. L., and Carris, L. M. 2000. Genetic variation among natural populations of *Tilletia controversa* and *T. bromi*. *Phytopathology* 90:376-383.
22. Shannon, C. E., and Weaver, W. 1949. *The Mathematical Theory of Communication*. University of Illinois, Urbana.
23. Taylor, B. H., and Powell, A. 1982. Isolation of plant DNA and RNA. *Focus* 4:4-6.
24. Tooley, P. W., Goley, E. D., Carras, M. M., Frederick, R. D., and Weber, E. L. 2001. Characterization of *Claviceps* species pathogenic on sorghum by sequence analysis of the β -tubulin gene intron 3 region and EF-1 α gene intron 4. *Mycologia* 93:541-551.
25. Tooley, P. W., O'Neill, N. R., Goley, E. D., and Carras, M. M. 2000. Assessment of diversity in *Claviceps africana* and other *Claviceps* species by RAM and AFLP analyses. *Phytopathology* 90:1126-1130.
26. Tsukiboshi, T., Shimanuki, T., and Uematsu, T. 1999. *Claviceps sorghicola* sp. nov., a destructive ergot pathogen of sorghum in Japan. *Mycol. Res.* 103:1403-1408.
27. Tudzynski, P. 1999. Genetics of *Claviceps purpurea*. Pages 79-93 in: *Ergot: The Genus Claviceps*. V. Kren and L. Cvak, eds. Harwood Academic Publishers, Amsterdam.
28. Tudzynski, P., Dubell, A., and Esser, K. 1983. Extrachromosomal genetics of *Claviceps purpurea*. I. Mitochondrial DNA and mitochondrial plasmids. *Curr. Genet.* 7:145-150.
29. Viji, G., Wu, B., Kang, S., Uddin, W., and Huff, D. R. 2001. *Pyricularia grisea* causing gray leaf spot of perennial ryegrass turf: Population structure and host specificity. *Plant Dis.* 85:817-826.
30. Weir, B. S., and Cockerham, C. C. 1984. Estimating F-statistics for the analysis of population structure. *Evolution* 38:1358-1370.
31. Weir, T. L., Huff, D. R., Christ, B. J., and Romaine, C. P. 1998. RAPD-PCR analysis of genetic variation among isolates of *Alternaria solani* and *Alternaria alternata* from potato and tomato. *Mycologia* 90:813-821.
32. White, J. F., Jr., Sullivan, R., Moy, M., Patel, R., and Duncan, R. 2000. An overview of problems in the classification of plant-parasitic *Clavicipitaceae*. *Studies in Mycology* No. 45:95-105.
33. Wright, S. 1978. *Evolution and the Genetics of Populations*. University of Chicago, Chicago.
34. Yeh, F. C., Yang, R.-C., Boyle, T. B. J., Ye, Z.-H., and Mao, J. X. 1997. POPGENE, the user-friendly shareware for population genetic analysis. *Molecular Biology and Biotechnology* Centre, University of Alberta, Canada.