Isothiocyanates Produced by Brassicaceae Species as Inhibitors of *Fusarium oxysporum*

U. Smolinska, Research Institute of Vegetable Crops, Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland; M. J. Morra and G. R. Knudsen, Soil Science Division, University of Idaho, Moscow 83844-2339; and R. L. James, USDA Forest Service, Northern Region, Forest Health Protection, 3815 Schreiber Way, Coeur d'Alene, ID 83814-8363

ABSTRACT

Smolinska, U., Morra, M. J., Knudsen, G. R., and James, R. L. 2003. Isothiocyanates produced by Brassicaceae species as inhibitors of *Fusarium oxysporum*. Plant Dis. 87:407-412.

Glucosinolates contained in members of the Brassicaceae release isothiocyanates potentially useful in controlling *Fusarium oxysporum* pathogens in conifer seedling nursery soils. Our objective was to determine the toxicity of individual isothiocyanates to different growth stages of the fungus. Bioassays with four *F. oxysporum* isolates were conducted using sealed containers in which 0.3 µl of 2-propenyl, ethyl, butyl, phenylethyl, benzyl, or phenyl isothiocyanate was allowed to volatilize. Propenyl and ethyl isothiocyanates were the most fungistatic of those compounds tested. The same concentrations of propenyl and ethyl isothiocyanates that inhibited mycelial growth completely suppressed conidial and chlamydospore germination of all isolates. Other isothiocyanates including ethyl, benzyl, and phenethyl were also fungitoxic to *F. oxysporum* conidia and chlamydospores. Reduction in pathogen populations resulting from a green-manure crop are likely achievable since chlamydospores are sensitive to isothiocyanate. Pathogenic *F. oxysporum* isolates infesting nursery soils would likely be most suppressed by species of plants such as *Brassica carinata*, *B. nigra*, and *B. juncea*, which contain glucosinolates that release high concentrations of propenyl isothiocyanate.

Additional keywords: forest nurseries, soil fumigation, soilborne fungi

Fusarium oxysporum Schlechtend. is a common soilborne fungus that causes damping-off, hypocotyl and root rot, and stunting or mortality of Douglas-fir nursery seedlings (3-5,22). These diseases, which occur most often in the first year of growth, are the primary causes of seedling mortality and can result in significant losses in forest nurseries (21). *F. oxysporum* is a soil inhabitant that infects plants through direct penetration or wounds in the roots. This fungus forms thick-walled resting structures called chlamydospores, thin-walled, sickle-shaped macroconidia, and smaller, one-celled microconidia, all of which act as potential sources of infection (6). The species also produces sclerotia that along with chlamydospores exist as resting structures in soil (35).

Fumigation of soil with compounds such as methyl bromide and chloropicrin reduces seedling losses (23). However, fumi-

gant use is being reduced due to environmental and human health concerns and prohibitive costs of reregistration. For example, methyl bromide will not be produced in or imported into the United States after the year 2005 as directed by the EPA, the 1990 Clean Air Act, and subsequent rulings. Production of alternate fumigants, Telone and Vorlex, containing the active ingredients dichloropropene/dichloropropane and methyl isothiocyanate, has been canceled in California (50). Alternative control measures are thus necessary to replace synthetic organic pesticides.

Plants of the order Capparales, especially agriculturally important *Brassica* spp. of the Brassicaceae, contain allelochemicals that reduce numbers of certain soilborne pathogens (11,34,36,37,44). These toxic effects are linked to the biologically active degradation products of glucosinolates. Glucosinolates themselves possess limited biological activity; however, enzymatic degradation by myrosinase (thioglucoside glucohydrolase, EC 3.2.3.1) results in the formation of a number of allelochemicals (9,12,47). More than 20 different aliphatic and aromatic isothiocyanates together with other potential allelochemicals have been identified among degradation products of glucosinolates originating from *Brassica napus*, *B. hirta*, *B. campestris*, *B. juncea*, and *B. nigra* (7,8,10,45). Each species has a unique

glucosinolate profile resulting in correspondingly different isothiocyanates.

Walker et al. (49) first described inhibition of fungi by pure isothiocyanates, and it has since been shown that isothiocyanates are general biocides that are highly toxic to some fungi (17,18). Inhibition of mycelial growth of *Gaeumannomyces graminis* (1), *Leptosphaeria maculans* (32), *Rhizoctonia solani*, *Fusarium graminearum*, *Bipolaris sorokiniana*, and *Pythium irregulare* (39) in the presence of isothiocyanates has been observed. Others have reported the effect of isothiocyanates on germination of *Botrytis cinerea* (16) and *Glomus mossae* (48) spores, and sporangia of *Peronospora parasitica* (20).

Toxicity or inhibition of fungal growth by Brassicaceae tissues has been demonstrated and linked to volatile glucosinolate hydrolysis products such as isothiocyanates (26,31,40,43,44). It has been determined that fungi respond differently to specific isothiocyanates (26,29,39,43), but regarding the differential sensitivity of *F. oxysporum* growth stages to individual compounds, little is known. This makes it difficult to select a suitable green-manure crop that might be used to reduce pathogen populations. The aims of this study were to (i) determine the sensitivity of four *F. oxysporum* strains to different isothiocyanates; and (ii) evaluate the stage in fungal development that is most sensitive to these compounds. Our overall goal was to use laboratory bioassays to predict the most promising green-manure crop to control diseases in forest nurseries caused by *F. oxysporum*.

MATERIALS AND METHODS

Fungal cultures. *F. oxysporum* isolates were obtained from forest tree nurseries in Idaho and Washington: (i) 9051C = container-grown white pine (*Pinus monticola*); (ii) 9243G = container-grown Douglas-fir (*Pseudotsuga menziesii* var. *glauca*); (iii) 9312F = bareroot nursery soil; and (iv) 9321A = bareroot nursery soil. All isolates were shown to be aggressive pathogens in laboratory tests on young Douglas-fir seedlings (24) and stored in sterile, silt-loam soil (13). Fungi were grown on potato dextrose agar (PDA) (Difco Laboratories, Detroit, MI).

Isothiocyanates. Pure chemical compounds were used in all experiments. Pro-

Corresponding author: M. J. Morra
E-mail: mmorra@uidaho.edu

Accepted for publication 17 November 2002.

Publication no. D-2003-0210-02R

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 2003.

penyl, ethyl, butyl, phenylethyl, and benzyl isothiocyanates were obtained from Aldrich Chemical (Milwaukee, WI). Phenyl isothiocyanate was obtained from Baker Chemical (Phillipsburg, NJ).

Mycelial growth. An agar disk (5 mm in diameter) was cut from a *F. oxysporum* colony growing on PDA and transferred to a new 100-mm-diameter polystyrene petri plate containing V8 juice agar (46). Each petri plate was placed in a large glass dish (330-cm³ volume) along with a glass filter (25 mm in diameter, Gelman AP20) to which 0.3 µl of the isothiocyanate was added. The glass dishes were quickly covered and sealed with plastic laboratory film (Parafilm "M", American National Can, Greenwich, CT). Dishes were incubated at 25°C and radial colony growth was measured daily for 8 days. In a second set of experiments, 1 µl of propenyl isothiocyanate was added to the glass filter and the container covered as noted above. However, the dish was opened after 24 h and the glass filter removed to evaluate if propenyl isothiocyanate was fungistatic or fungitoxic by allowing the fungus to grow for the next 7 days.

Conidial formation. Test isolates were grown on V8 juice agar for 8 days in a covered glass dish in which 0.3 µl of isothiocyanate was allowed to evaporate as

detailed in mycelial growth assays. An agar disk (12 mm in diameter) was cut from the colony approximately 1 cm from its edge and placed in 2 ml of sterile, distilled water. After stirring for 10 s to dislodge the conidia, conidial density was counted using a haemocytometer. No distinction was made between micro- and macroconidia.

Conidial germination. Conidia of *F. oxysporum* were produced according to Couteaudier and Alabouvette (14). Isolates were grown in a malt extract agar liquid medium (10 g/liter) on a rotary shaker at 22 to 25°C. Five 6-day-old cultures were filtered through four layers of cheesecloth to remove mycelial mats. Conidia were pelleted by centrifugation and washed three times in sterile, distilled water. The concentration of the final conidial suspension as determined with a haemocytometer was 10⁴ conidia/ml. Sterile microscope slides were coated with a layer of 2% agar in water to obtain a thin layer of medium on the slide surface, placed in glass petri dishes (330-cm³ volume) lined with moistened filter paper, and a 100-µl sample of spore suspension was spread on each slide. A glass filter with 0.3 µl of isothiocyanate was placed in each dish and then covered and sealed with plastic film. Slides were incubated overnight in the presence of individual isothiocyanates, followed by

incubation for 7 h at 25°C, and stored at 10°C to avoid overgrowth. One hundred conidia were assessed microscopically on each slide to determine germination percentage. Slides were placed in glass dishes, incubated overnight at 25°C without isothiocyanate, and 100 spores on each slide were examined for germination to determine if isothiocyanates were fungitoxic or fungistatic.

Chlamydospore germination. Macroconidia were produced at 25°C in a malt extract agar liquid medium on a rotary shaker (14). Conidia were pelleted by centrifugation and suspended in sterile, distilled water. A dry silt loam soil was sieved and autoclaved (1 h followed by 30 min of autoclaving 24 h later). Conidia were added to the moistened soil and then allowed to dry for 2 to 3 weeks. The soil was mixed, remoistened, and dried for another 2 to 3 weeks during which time conidia transformed into chlamydospores or lysed (41). Dry soil was crushed using a mortar and pestle, and a 5-g sample was added to 45 ml of sterile, distilled water followed by mixing for 10 min on a rotary shaker. Approximately 10⁵ chlamydospores per g of soil were formed and 100 µl of a 10⁻³ dilution of the original soil suspension was spread on petri plates with a selective agar medium for *F. oxysporum* (28). Each petri plate was placed in a glass dish (330-cm³ volume) along with a glass filter to which 0.3 µl of isothiocyanate was added. The dishes were covered, sealed with a plastic film, and *F. oxysporum* colonies were counted on each petri plate after 5 days of incubation at 25°C. Each colony was assumed to represent the germination of a single chlamydospore.

Mycelial growth, conidial formation, conidial germination, and chlamydospore germination experiments included three replicates per treatment. All experiments were performed twice, yielding similar data unless otherwise noted. Data for conidial germination were arcsine-transformed and subjected to analysis of variance using the General Linear Models program of SAS (SAS Institute, Inc. Cary, NC). Protected LSD values for the remaining data were determined with SAS using untransformed data.

RESULTS

Mycelial growth. Colony diameters were compared among treatments at day two and then again at day seven to determine whether initial growth response differences were maintained until experiment termination. Propenyl and ethyl isothiocyanate significantly reduced mycelial growth of each of the four *F. oxysporum* isolates at day two ($P < 0.05$) (Fig. 1). Butyl isothiocyanate reduced growth of three isolates (9051C, 9321A, and 9243G) at day two ($P < 0.05$), but growth inhibition was less than that induced by ethyl or propenyl isothiocyanate. In many cases,

Fig. 1. Mycelial growth of four *Fusarium oxysporum* isolates exposed to isothiocyanates (Prop = propenyl; Eth = ethyl; But = butyl; Phet = phenylethyl; Ben = benzyl; Ph = phenyl) present in the headspace of closed containers. The controls (ctrl) contained no isothiocyanate. Solid vertical bars at days two and six or seven represent LSD values for treatments at that respective day.

the inhibition noted at day two was not observed at day six or seven. None of the isothiocyanates caused a decrease in colony diameter for 9321A at day six and only ethyl isothiocyanate caused growth inhibition of 9312F at day seven. *F. oxysporum* isolate 9243G was the most sensitive toward all isothiocyanates. Benzyl and propenyl isothiocyanates showed the strongest, and butyl and phenyl isothiocyanates showed the weakest inhibition of this isolate at day seven.

Although propenyl and ethyl isothiocyanates had the most pronounced effect on all isolates of *F. oxysporum*, this effect was fungistatic, not fungitoxic, when grown in the presence of 0.3 μ l of each isothiocyanate. Propenyl isothiocyanate added to the enclosed container in an amount of 1.0 μ l completely suppressed growth of *F. oxysporum* isolate 9312F during the entire 8 days of monitoring (Fig. 2). All other isolates behaved similarly (data not shown). When exposed to 1.0 μ l of this compound for only 24 h, all isolates resumed growth a few days later. However, when this experiment was repeated, 24-h incubation of fungi in the presence of 1.0 μ l of propenyl isothiocyanate completely suppressed mycelial growth of all isolates except 9243G.

Conidia formation and viability. The numbers of conidia formed during 10 days of growth of *F. oxysporum* isolates on V8 juice agar in the presence of different isothiocyanates were compared (Table 1). In most cases, isothiocyanates did not reduce sporulation. Only benzyl isothiocyanate decreased the number of spores that isolate 9312F produced ($P < 0.05$), however this same compound increased sporulation of 9051C as compared to the control ($P < 0.05$). Viability of spores (only isolates 9321A and 9051C were tested) formed in the presence of isothiocyanates was no different from the control treatment (data not presented). Data from the repeated experiment were consistent with the presented results.

Fig. 2. Mycelial growth of *Fusarium oxysporum* 9312F exposed to propenyl isothiocyanate vapor for 24 h or 8 days in closed containers. The controls contained no isothiocyanate.

Conidial germination. Exposure of conidia of all *F. oxysporum* isolates to isothiocyanate for 24 h completely suppressed their germination (Fig. 3). After subsequent incubation of spores for another 24 h without isothiocyanates, conidia treated previously with butyl isothiocyanate and phenyl isothiocyanate germinated. Prolonged incubation of conidia treated during the first 24 h with propenyl, ethyl, phenethyl, or benzyl isothiocyanate

did not reverse the inhibition of germination. Data from the repeated experiment were consistent with the presented results.

Chlamydospore germination. All isothiocyanates used in this experiment in amounts of 0.3 μ l inhibited chlamydospore germination ($P < 0.05$) (Table 2). Propenyl and ethyl isothiocyanates completely suppressed the formation of fungal colonies of each tested isolate. The fungitoxic effect of phenethyl isothiocyanate was somewhat

Table 1. Sporulation of three isolates of *Fusarium oxysporum* in the presence of different types of isothiocyanates (ITC)^x

Treatment	Isolate		
	9051C	9321A	9312F
Control	47.7 b ^y	83.6 a ^z	13.5 ab
Propenyl ITC	42.8 b	82.5 a	16.5 a
Ethyl ITC	46.7 b	103.4 a	16.9 a
Butyl ITC	46.9 b	84.3 a	14.2 ab
Phenethyl ITC	50.4 ab	96.6 a	12.8 b
Benzyl ITC	58.0 a	100.4 a	9.3 c
Phenyl ITC	45.4 b	102.8 a	12.9 b

^x Sporulation = conidia $\times 10^5$ /cm² colony.

^y Each value is the mean of three replicates. Each replicate is the mean of two to three samples from one colony.

^z Numbers in the same column followed by the same letter are not statistically different ($P > 0.05$) according to LSD analysis.

Fig. 3. Conidial germination of four *Fusarium oxysporum* isolates exposed for 24 h to isothiocyanates (Prop = propenyl; Eth = ethyl; But = butyl; Phet = phenylethyl; Ben = benzyl; Ph = phenyl) present in the headspace of closed containers. Fungitoxicity was determined by monitoring germination in the absence of the respective isothiocyanates for an additional 24 h. The controls (ctrl) contained no isothiocyanate. Means labeled with the same letter and within the same 24-h assessment period are not different ($P < 0.05$).

variable. In the second experiment, although phenethyl isothiocyanate decreased the number of germinating chlamydospores compared to the control, this inhibition was never complete. Phenyl and butyl isothiocyanate reduced chlamydospore germination of *F. oxysporum* ($P < 0.05$) by 30 to 90% depending on the isolate tested.

DISCUSSION

Inhibition of mycelial growth as a result of exposure to isothiocyanates has been observed for a variety of fungi (1,30,39). Our work showed that isothiocyanates inhibited mycelial growth of *F. oxysporum*, but at exposure levels to which they were subjected in this study, the effects were predominantly fungistatic, not fungitoxic. Fungitoxicity could be induced only by increasing the concentration and respective vapor pressure of the most inhibitory isothiocyanate, and even then the results varied with the isolate. Sarwar et al. (39) similarly determined that some isothiocyanates are fungitoxic to *F. graminearum* after a 24-h exposure period, whereas others are only fungistatic.

Additional investigations demonstrate that volatile compounds released from glucosinolate-containing plants (26,31) also inhibit mycelial growth. Ramirez-Villapudua and Munnecke (38) indicate that growth inhibition occurs when subjecting *F. oxysporum* f. sp. *conglutinans* to gases evolved from decomposing cabbage residues. Radial growth was inhibited when the fungus was grown on petri plates suspended above cabbage-amended soil placed in closed containers. Mycelial growth resumed on transfer of the fungus to fresh PDA. In other studies, fungitoxicity to *Verticillium dahliae*, *V. albo-atrum*, *Pythium ultimum*, *Colletotrichum coccodes*, and *R. solani* grown on V8 juice agar was observed in the presence of gases evolved from *B. juncea* (31). Direct comparison of bioassays conducted using *Brassica* tissues to those conducted with specific isothiocyanates must be approached cautiously since tissues produce a variety of volatile inhibitory compounds (2,7,8) that may express additive or synergistic effects on the bioassay organism. This has led Bending and Lincoln (2) to suggest that

other volatile S compounds are likely to be as important as isothiocyanates in pest control. In addition, dose response comparisons are not possible without accurate measurement of gas phase concentrations of the responsible toxicants.

The extent of mycelial growth inhibition varied among isolates depending on the specific isothiocyanate to which the respective isolate was subjected. Differences in mycelial growth inhibition of different isolates of the same species in the presence of volatile compounds produced by cold-pressed *B. juncea* seed meal has been observed for *G. graminis* var. *tritici*, *R. solani*, *P. irregulare*, and *Bipolaris sorokiniana* (26). Variability in susceptibility to the isothiocyanates used here is especially important given the fact that all isolates were obtained from forest nurseries. This variable response must thus be considered when testing the efficacy of control strategies involving the use of isothiocyanate-releasing plant tissues.

Propenyl and ethyl isothiocyanates were the most inhibitory of all tested isothiocyanates to mycelial growth of the four *F. oxysporum* isolates. Correlations of reduced fungal growth with the release of propenyl isothiocyanate from *B. juncea* plant tissues have been observed (26,31). Inhibition of mycelial growth of *G. graminis* by propenyl isothiocyanate has also been reported (1). Other investigators have demonstrated the inhibitory nature of additional isothiocyanates including 2-phenylethyl (15) and those containing sulfur side chains such as 3-methyl-sulfinylpropyl, 3-methylsulfonylpropyl, and 4-methylthiobutyl isothiocyanates (29) to *Fusarium* species. However, incorporating isothiocyanates into the media as performed in the latter investigations makes comparisons with our bioassays conducted with only volatiles tenuous. Sarwar et al. (39) showed that aromatic isothiocyanates were more toxic than aliphatic isothiocyanates when incorporated in agar, but less toxic in volatile form as a result of lower vapor pressures. This work indicates that increased predictive capabilities concerning toxicities of individual isothiocyanates can only be achieved by considering their respective partitioning in a three-phase soil environment.

Mycelial growth was inhibited to a relatively small extent and thus represents a reasonably resistant stage in the life cycle of *F. oxysporum* isolates used in this work. Likewise, conidial formation was little affected by those isothiocyanates tested. In contrast, conidial and chlamydospore germination were highly susceptible to inactivation by isothiocyanates. Thus these two stages in the life cycle of *F. oxysporum* are the most susceptible to inhibition by a green-manure treatment. Propenyl, ethyl, benzyl, and in most cases, phenethyl isothiocyanates completely suppressed conidial and chlamydospore germination. In contrast, the toxic effects of phenyl and butyl isothiocyanates, although significant compared to the control, were almost always less than other isothiocyanates. These results indicate that fungicidal activity of isothiocyanates is strongly determined by the nature of the molecular structure, but that activity is not correlated with aliphatic or aromatic character.

These results are consistent with those of Smolinska and Horbowicz (43) in which the investigators measured the release of propenyl isothiocyanate from *B. juncea* tissues and determined that a correlation exists between the presence of this compound in volatile form and the inhibition of germination of chlamydospores of *F. oxysporum* var. *radicis* f. sp. *lycopersici*. As in our work, the effect was nonreversible, and thus propenyl isothiocyanate appeared to be fungitoxic. In addition, *B. juncea* tissues have been shown to reduce *F. oxysporum* f. sp. *lycopersici* chlamydospore viability and lessen wilt of tomato disease symptoms (42). These observations are inconsistent with speculation of Greenhalgh and Mitchell (20) that activity of propenyl isothiocyanate toward *Phytophthora parasitica* mycelium should be more deleterious than to sporangia, because the first is an active and the second is an inactive stage in fungal development.

From these results, we conclude that decreased pathogen populations resulting from the direct inhibition of *F. oxysporum* mycelial growth seem unlikely. Changes in populations of these organisms resulting from a green-manure crop will only be achievable if conidia and chlamydospores are targeted. This is a feasible approach since *F. oxysporum* may exist almost exclusively as chlamydospores until stimulated to germinate by the root tip of a susceptible host. Incorporation of a green-manure crop prior to sowing conifers may reduce the number of germinating chlamydospores, thereby decreasing the likelihood of seedling infection. Attempts at controlling growth stages of the fungus following chlamydospore germination will be less effective.

Pathogenic *F. oxysporum* isolates typically infesting nursery soils in western North America will likely be most suppressed by species of plants containing

Table 2. Germination of *Fusarium oxysporum* chlamydospores in the presence of isothiocyanates (ITC)

Treatment	Isolate			
	9051C	9321A	9312F	9243G
Control	169.7 a ^y	54.0 a ^z	63.0 a	123.7 a
Propenyl ITC	0.0 c	0.0 c	0.0 c	0.0 d
Ethyl ITC	0.0 c	0.0 c	0.0 c	0.0 d
Butyl ITC	16.3 c	21.3 b	38.3 b	31.7 b
Phenethyl ITC	0.0 c	0.0 c	32.0 b	0.0 d
Benzyl ITC	3.0 c	0.0 c	0.7 c	2.7 d
Phenyl ITC	70.0 b	19.3 b	44.0 b	20.3 c

^y All values represent CFU. Each value is the mean of three replicates.

^z Numbers in the same column followed by the same letter are not statistically different ($P > 0.05$) according to LSD analysis.

glucosinolates that release the highest concentrations of propenyl isothiocyanate. Other glucosinolates releasing ethyl, benzyl, and phenethyl isothiocyanates should also contribute to potential suppression of *F. oxysporum*. Propenyl glucosinolate is found in high concentrations in shoots of three mustard species (*B. carinata*, *B. nigra*, and *B. juncea*) and phenylethyl glucosinolate is most commonly found at high concentrations in the roots of rapeseed or *B. napus* cultivars (25). Several weed species contain high concentrations of benzyl glucosinolate (25), whereas *Lepidium menziesii* is the only species in which ethyl glucosinolate has been identified (27).

It is important to note that only a fraction of the isothiocyanate potentially available from the measured molar quantity of glucosinolate within the tissues is actually released and available for pest inhibition. In laboratory studies, leaf and stem tissues of *B. napus* have been shown to release only 8% of the butenyl isothiocyanate theoretically possible as calculated from the tissue glucosinolate concentration (8). Field investigations conducted by measuring allelochemicals released in soil after a *B. napus* green-manure crop was incorporated showed that compounds derived from root tissues dominated the profile of measured compounds and only traces of shoot glucosinolate products were detected (19). More recent investigations indicate that physical methods are required to maximize the rupture of plant cell membranes and release isothiocyanate (33). Ensuring efficient isothiocyanate release from a green-manure crop is critical since organic carbon inputs without effective *F. oxysporum* control may serve to increase pathogen populations.

The first step in the use of green-manure crops to control pathogenic *F. oxysporum* populations in forest nursery soils is thus to choose plants with high concentrations of the appropriate glucosinolates. Release of the most inhibitory isothiocyanates at a time appropriate to target a susceptible growth stage of the fungus will increase the likelihood of pest suppression. Our work indicates that chlamydospores may be inhibited prior to the introduction of conifer seedlings if the release of isothiocyanate from the selected green-manure crop is maximized.

ACKNOWLEDGMENTS

Research supported by USDA CSREES grants 93-34103-8693 and 93-COOP-1-8648, and state and federal funds provided to the Idaho Agricultural Experiment Station.

LITERATURE CITED

1. Angus, J. F., Gardner, P. A., Kirkegaard, J. A., and Desmarchelier, J. M. 1994. Biofumigation: Isothiocyanates released from *Brassica* roots inhibit growth of the take-all fungus. *Plant Soil* 162:107-112.
2. Bending, G. D., and Lincoln, S. D. 1999. Characterisation of volatile sulphur-containing compounds produced during de-

- composition of *Brassica juncea* tissues in soil. *Soil Biol. Biochem.* 31:695-703.
3. Bloomberg, W. J. 1971. Diseases of Douglas-fir seedlings caused by *Fusarium oxysporum*. *Phytopathology* 61:467-470.
4. Bloomberg, W. J. 1976. Distribution and pathogenicity of *Fusarium oxysporum* in a forest nursery soil. *Phytopathology* 66:1090-1092.
5. Bloomberg, W. J., and Lock, W. 1972. Strain differences in *Fusarium oxysporum* causing diseases of Douglas-fir seedlings. *Phytopathology* 62:481-485.
6. Booth, C. 1971. *The Genus Fusarium*. Commonwealth Mycological Institute, Kew, Surrey, England.
7. Brown, P. D., and Morra, M. J. 1995. Glucosinolate-containing plant tissues as bioherbicides. *J. Agric. Food Chem.* 43:3070-3074.
8. Brown, P. D., and Morra, M. J. 1996. Hydrolysis products of glucosinolates in *Brassica napus* tissues as inhibitors of seed germination. *Plant Soil* 181:307-316.
9. Brown, P. D., and Morra, M. J. 1997. Control of soil-borne plant pests using glucosinolate-containing plants. *Adv. Agron.* 61:167-231.
10. Brown, P. D., Morra, M. J., McCaffrey, J. P., and Williams, L. III. 1991. Allelochemicals produced during glucosinolate degradation in soil. *J. Chem. Ecol.* 17:2021-2034.
11. Chan, M. K. Y., and Close, R. C. 1987. *Aphanomyces* root rot of peas 3. Control by use of cruciferous amendments. *N.Z. J. Agric. Res.* 30:225-233.
12. Chew, F. S. 1988. Biological effects of glucosinolates. Pages 155-181 in: *Biologically Active Natural Products: Potential Use in Agriculture*. H. G. Cutler, ed. American Chemical Society, Washington, DC.
13. Cormack, M. W. 1951. Variation in the cultural characteristics and pathogenicity of *Fusarium avenaceum* and *F. arthrosporioides*. *Can. J. Bot.* 29:32-45.
14. Couteaudier, Y., and Alabouvette, C. 1990. Survival and inoculum potential of conidia and chlamydospores of *Fusarium oxysporum* f. sp. *lini* in soil. *Can. J. Microbiol.* 36:551-556.
15. Davis, D. 1964. Host fungitoxins in selective pathogenicity of *Fusarium oxysporum*. *Phytopathology* 54:290-293.
16. Dawson, G. W., Doughty, K. J., Hick, A. J., Pickett, J. A., Pye, B. J., Smart, L. E., and Wadhams, L. J. 1993. Chemical precursors for studying the effects of glucosinolate catabolites on diseases and pests of oilseed rape (*Brassica napus*) or related plants. *Pestic. Sci.* 39:271-278.
17. Drobnicova, L., Zemanova, M., Nemec, P., Antos, K., Kristian, P., Stullerova, A., Knopova V., and Nemec, J. R. P. 1967. Antifungal activity of isothiocyanates and related compounds. I. Naturally occurring isothiocyanates and their analogues. *Appl. Microbiol.* 15:701-709.
18. Drobnicova, L., Zemanova, M., Nemec, P., Kristian, P., Antos, K., and Hulka, A. 1967. Antifungal activity of isothiocyanates and related compounds. II. Mononuclear aromatic isothiocyanates. *Appl. Microbiol.* 15:710-717.
19. Gardiner, J. B., Morra, M. J., Eberlein, C. V., Brown, P. D., and Borek, V. 1999. Allelochemicals released in soil following incorporation of rapeseed (*Brassica napus*) green-manures. *J. Agric. Food Chem.* 47:3837-3842.
20. Greenhalgh, J. R., and Mitchell, N. D. 1976. The involvement of flavour volatiles in the resistance to downy mildew of wild and cultivated forms of *Brassica oleracea*. *New Phytol.* 77:391-398.
21. Hamm, P. B., Campbell, S. J., and Hansen, E. M. 1990. Growing healthy seedlings. Identifi-

- cation and management of pests in Northwest forest nurseries. Forest Pest Management, USDA Forest Service, Pacific Northwest Region, and Forest Res. Lab., College of Forestry, Oregon State University, Corvallis.
22. Hansen, E. M., and Hamm, P. B. 1988. Canker diseases of Douglas-fir seedlings in Oregon and Washington bareroot nurseries. *Can. J. For. Res.* 18:1053-1058.
23. Hansen, E. M., Myrold, D. D., and Hamm, P. B. 1990. Effects of soil fumigation and cover crops on potential pathogens, microbial activity, nitrogen availability, and seedling quality in conifer nurseries. *Phytopathology* 80:698-704.
24. James, R. L., Perez, R., Dumroese, R. K., and Wenny, D. L. 2000. Virulence of *Fusarium oxysporum* on Douglas-fir germinants: Comparison of isolates from nursery soil and roots of healthy and diseased seedlings. Pages 49-64 in: *Proceeding of the 4th meeting of IUFRO Working Party 7.03.04-Diseases and insects in forest nurseries*. A. Lilja and J.R. Sutherland, eds. Finnish Forest Research Institute, Research Papers 781.
25. Kirkegaard, J. A., and Sarwar, M. 1998. Biofumigation potential of brassicas. I. Variation in glucosinolate profiles of diverse field-grown brassicas. *Plant Soil* 201:71-89.
26. Kirkegaard, J. A., Wong, P. T. W., and Desmarchelier, J. M. 1996. *In vitro* suppression of fungal root pathogens of cereals by *Brassica* tissues. *Plant Pathol.* 45:593-603.
27. Kjaer, A., and Larsen, I. 1954. Isothiocyanate IX. The occurrence of ethyl isothiocyanate in nature. *Acta Chem. Scand.* 8:699-701.
28. Komada, H. 1975. Development of a selective medium for quantitative isolation of *Fusarium oxysporum* from natural soil. *Rev. Plant Prot. Res. (Japan)* 8:114-125.
29. Manici, L. M., Lazzeri, L., and Palmieri, S. 1997. *In vitro* fungitoxic activity of some glucosinolates and their enzyme-derived products toward plant pathogenic fungi. *J. Agric. Food Chem.* 45:2768-2773.
30. Mari, M., Iori, R., Leont, O., and Marchi, A. 1993. *In vitro* activity of glucosinolate-derived isothiocyanates against postharvest fruit pathogens. *Ann. Appl. Biol.* 123:155-164.
31. Mayton, H. S., Olivier, C., Vaughn, S. F., and Loria, R. 1996. Correlation of fungicidal activity of *Brassica* species with allyl isothiocyanate production in macerated leaf tissue. *Phytopathology* 86:267-271.
32. Mithen, R. F., and Lewis, B. G. 1986. *In vitro* activity of glucosinolates and their products against *Leptosphaeria maculans*. *Trans. Br. Mycol. Soc.* 87:433-440.
33. Morra, M. J., and Kirkegaard, J. A. 2002. Isothiocyanate release from soil-incorporated *Brassica* tissues. *Soil Biol. Biochem.* 34:1683-1690.
34. Muehlchen, A. M., Rand, R. E., and Parke, J. L. 1990. Evaluation of crucifer green-manures for controlling *Aphanomyces* root rot of peas. *Plant Dis.* 74:651-654.
35. Nelson, P. E., Toussoun, T. A., and Marasas, W. F. O. 1983. *Fusarium* Species: An Illustrated Manual for Identification. The Pennsylvania State University Press, University Park.
36. Papavizas, G. C. 1966. Suppression of *Aphanomyces* root rot of peas by cruciferous soil amendments. *Phytopathology* 56:1071-1075.
37. Ramirez-Villapudua, J., and Munnecke, D. E. 1987. Control of cabbage yellow (*Fusarium oxysporum* f. sp. *conglutinans*) by solar heating of field soils amended with dry cabbage residues. *Plant Dis.* 71:217-221.
38. Ramirez-Villapudua, J., and Munnecke, D. E. 1988. Effect of solar heating and soil amend-

- ments of cruciferous residues on *Fusarium oxysporum* f. sp. *conglutinans* and other organisms. *Phytopathology* 78:289-295.
39. Sarwar, M., Kirkegaard, J. A., Wong, P. T. W., and Desmarchelier, J. M. 1998. Biofumigation potential of brassicas. III. *In vitro* toxicity of isothiocyanates to soil-borne fungal pathogens. *Plant Soil* 201:103-112.
 40. Singh, R. S., and Singh, N. 1970. Effect of oil-cake amendment of soil on populations of some wilt causing species of *Fusarium*. *Phytopathology* 69:160-167.
 41. Smith, S. N., and Snyder, W. C. 1971. Germination of *Fusarium oxysporum* chlamydospores in soil favorable and unfavorable to wilt establishment. *Phytopathology* 62:273-277.
 42. Smolinska, U. 2000. Survival of *Sclerotium cepivorum* sclerotia and *Fusarium oxysporum* chlamydospores in soil amended with cruciferous residues. *J. Phytopathol.* 148:343-349.
 43. Smolinska, U., and Horbowicz, M. 1999. Fungicidal activity of volatiles from selected cruciferous plants against resting propagules of soil-borne fungal pathogens. *J. Phytopathol.* 147:119-124.
 44. Smolinska, U., Knudsen, G. R., Morra, M. J. and Borek, V. 1997. Inhibition of *Aphanomyces euteiches* f. sp. *pisi* by volatiles produced by hydrolysis of *Brassica napus* seed meal. *Plant Dis.* 81:288-292.
 45. Spencer, G. F., and Daxenbichler, M. E. 1980. Gas chromatography-mass spectrometry of nitriles, isothiocyanates and oxazolidinethiones derived from cruciferous glucosinolates. *J. Sci. Food Agric.* 31:359-367.
 46. Toussoun, T. A., and Nelson, P. E. 1976. A Pictorial Guide to the Identification of *Fusarium* Species. The Pennsylvania State University Press, University Park.
 47. Underhill, E. W. 1980. Glucosinolates. Pages 493-511 in: *Encyclopedia of Plant Physiology*. Vol.8. Secondary Plant Products. E. A. Bell and B. V. Charwood, eds. Springer-Verlag, New York.
 48. Vierheilig, H., and Ocampo, J. A. 1990. Effect of isothiocyanates on germination *G. mosseae*. *Soil Biol. Biochem.* 22:1161-1162.
 49. Walker, J. C., Morell, S., and Foster, H. H. 1937. Toxicity of mustard oils and related sulfur compounds to certain fungi. *Am. J. Bot.* 24:536-541.
 50. White, J. 1994. After methyl bromide: No easy answers. *Calif. Agric.* 48:7-9.