Gray Leaf Spot (Blast) of Perennial Ryegrass Turf An Emerging Problem for the Turfgrass Industry

Perennial ryegrass (*Lolium perenne* L.) is a cool-season grass originated from open areas and forest fringes of southern Europe and western Asia. It was first used in England in 1677 as a forage grass. At present, it is widely used in the turf industry, especially on golf course fairways, due to its superior agronomic attributes such as turf color, upright and bunch growth habit, rapid germination and coverage, tolerance to close-mowing and soil compaction, and absence of thatch (25). Additionally, tolerance to cold weather has led to the use of perennial ryegrass for over-seeding dormant bermudagrass golf tees and fairways during the fall in the southern United States.

Gray leaf spot or blast is caused by *Magaporthe grisea* (T.T. Hebert) Yaegashi & Udagawa, anamorph *Pyricularia grisea* (Cooke) Sacc. Recently, Couch and Kohn (8) proposed dividing *M. grisea* into two species, *M. grisea* and *M. oryzae*, for isolates causing disease on several gramineous hosts including perennial ryegrass. However, *M. grisea* is likely to remain as the standard name for the teleomorph of *P. grisea* until a formal name is adopted by the International Committee on Fungal Nomenclature.

In recent years, severe outbreaks of gray leaf spot have resulted in extensive damage to perennial ryegrass golf course fairways and athletic fields, particularly in the mid-western and northeastern United States. Under favorable conditions, the disease develops rapidly, and entire ryegrass swards can be killed within a few days, leaving only annual bluegrass and other grassy weeds that are not affected by the

disease. Since gray leaf spot was first detected on perennial ryegrass in 1991, the disease commonly has resulted in increased fungicide costs of \$20,000 to \$25,000 per year on courses with perennial ryegrass fairways, and certain resorts have reported losses of up to \$500,000 in revenue due to severe outbreaks of the disease (P. Vincelli, *unpublished*). Turfgrass managers are now considering replacing perennial ryegrass with other turfgrass species such as creeping bentgrass and Kentucky bluegrass due to the extensive and unpredictable damage caused by gray leaf spot. However, replacement of perennial ryegrass with these turfgrass species does not provide the best solution because of the superior agronomic characteristics of perennial ryegrass and the rapid thatch buildup and patch disease problems in other turfgrass species.

Hosts

M. grisea is pathogenic to more than 50 gramineous hosts including small grains, forage, and turfgrasses (1,2,16,28,31,35, 42,75). The fungus is probably best known for the devastating losses it can cause on rice (*Oryza sativa* L.). It may infect at any growth stage of the rice plant causing rapid leaf, nodal, neck, and panicle blighting that is referred to as blast. The term blast has also been used to describe the disease in other cereal grains such as wheat (*Triticum aestivum* L.) and finger millet (*Eleusine corocana*) (1). The disease in broadleaf hosts such as species of *Ctenanthe*, *Marantha*, and *Stromanthe* has been referred to as *Pyricularia* leaf spot (29).

In 1957, Malca and Owen (24) reported a leaf spot disease of St. Augustinegrass (*Stenotaphrum secundatum* Walt.) in Florida caused by *M. grisea*. They named the disease gray leaf spot rather than blast because it did not result in rapid leaf blighting or plant death. Gray leaf spot is a common foliar disease of St. Augustine-

grass in the southeastern United States (33,57); however, it generally does not cause damage to this grass species to the extent reported in perennial ryegrass (49). Severe outbreaks of a disease caused by *M. grisea* were observed on annual ryegrass (*Lolium multiflorum* Lam.) used for forage in the southern United States during the early 1970s (2,6). The disease was referred to as blast because of the resemblance of foliar symptoms (leaf spot and blighting) to leaf blast symptoms on rice (2,40).

In 1991, Landschoot and Hoyland (22) reported gray leaf spot on perennial ryegrass turf in golf course fairways in Pennsylvania. The epidemic was confined to the southeastern region of Pennsylvania where extensive damage was reported. Since this first report, outbreaks of gray leaf spot have occurred sporadically, resulting in serious loss of turf in 1995, 1998, and 2000 in the mid-Atlantic region (10,43). The disease has recently been reported from the Midwest (19,30,64), New England (32), and the western United States (54).

Pathogen Diversity and Biology

Significant progress has been made in research on population structure and biology of the gray leaf spot pathogen in recent years. Perennial ryegrass isolates collected from various regions of the United States using different transposons as DNA probes in restriction fragment length polymorphism (RFLP) analysis indicated the presence of three lineages (Fig. 1) (61). Comparison of DNA fingerprints obtained with the probe Pot2 showed a close relationship between isolates of *M. grisea* from perennial ryegrass and those from wheat and triticale. Furthermore, the perennial ryegrass isolates had the same sequence in the rDNA internal transcribed spacer (ITS) region as those of the wheat and triticale isolates (61). A close relationship between an isolate from perennial ryegrass and an isolate from wheat has also been reported

Corresponding author: Wakar Uddin
E-mail: wxu2@psu.edu

by Farman (15). However, the study indicated that the vast majority of perennial ryegrass isolates tested were distinct from wheat blast isolates based on the fingerprints obtained with Pot2 and MGR583. Additionally, the perennial ryegrass isolates were closely related to isolates from tall fescue (*Festuca arundinacea* L.) based on fingerprints obtained with these probes (15). *M. grisea* isolates from perennial ryegrass were genetically distinct from the isolates from rice, crabgrass (*Digitaria* spp.), and *Setaria* spp. This was evidenced by the uneven distribution of the MAGGY retrotransposon in the perennial ryegrass isolates, where some isolates contained 6 to 8 or 10 to 30 copies of the element and other isolates completely lacked the element (15).

Isolates of *M. grisea* from perennial ryegrass and wheat have been shown to cross-infect each other's hosts. Pathogenicity tests have indicated that isolates of *M. grisea* from perennial ryegrass are virulent on wheat and tall fescue (Fig. 2). The isolates also caused gray leaf spot symptoms on triticale but not on rice (61). Isolates from wheat and triticale were also virulent on perennial ryegrass, indicating that these cereal grain crops have the potential to serve as an additional source of inoculum for the spread of gray leaf spot.

The perfect stage of *P. grisea* was first described in 1971 (20) in crosses between isolates from cereals and wild grasses. Since then, efforts have been made to produce perithecia under controlled conditions using hermaphroditic tester strains to study the genetic relationships among isolates. In a recent study, 73 isolates of perennial ryegrass were paired with fertile hermaphroditic tester strains from finger millet, rice, and wheat in order to determine the mating type distribution, fertility, and degree of sexual compatibility (59). All isolates from perennial ryegrass produced perithecia only when crossed with *MAT1-1* tester strains and belonged to one mating type (*MAT1-2*), indicating that sexual recombination does not occur among these isolates. The perennial ryegrass isolates that formed perithecia were male-fertile (female sterile), suggesting that the capacity for expressing the female characteristic may have been lost. Additionally, none of the perennial ryegrass isolates were capable of forming asci or ascospores, regardless of their developmental stage. The low levels of fertility in the *M. grisea* isolates of perennial ryegrass along with the relatively simple population structure suggests the clonality of the pathogen, and also indicates an exclusive asexual reproduction.

Symptoms

Gray leaf spot develops on the leaf blades of perennial ryegrass as small, water-soaked lesions that subsequently turn into dark-colored, 1- to 3-mm-diameter necrotic spots. The spots expand rapidly and become gray, grayish-brown, or light brown, circular to oblong lesions with purple to dark brown borders that often are surrounded by a yellow halo (Fig. 3A). The necrotic lesions coalesce, become irregular in shape, and cause partial blighting (tip blighting) or complete blighting of the leaves. Blighted leaf blades may also exhibit twisting or flagging (Fig. 3B). Complete necrosis of the leaves results in death of the entire plant. There is no evidence of infection of crown tissue by the pathogen. Blighted leaf blades may appear grayish-white to tan and may have a dusty or velvety texture when conidia are produced profusely. Aerial mycelium is usually not evident on necrotic leaves under humid conditions.

Diseased turfgrass stands develop an off-color, diffuse blighted or wilted appearance. This is followed by the development of sunken or pocketed areas or irregularly shaped large patches. When severe, the entire ryegrass stand may be killed, leaving the annual bluegrass and other grassy weeds in the fairways (Fig. 4). The disease may be distributed along low-lying or

Fig. 1. Comparison of genetic similarity among *Magnaporthe grisea* isolates from perennial ryegrass using Pot2 fingerprinting probe. A, Fingerprints of representative isolates from Kansas (KS), Maryland (MD), New Jersey (NJ), Virginia (VA), West Virginia (WV), New York (NY), and Pennsylvania (PA). B, Phenogram derived by unweighted pair group method, arithmetic mean (UPGMA) based on Pot2 restriction fragment length polymorphism data.

drainage areas where high relative humidity and prolonged leaf wetness periods occur in the turf canopy (Fig. 5).

Gray leaf spot is usually confirmed by observing the hyaline, pyriform, septate (one to three septa) conidia on diseased leaf tissue (14) (Fig. 6). One of the major challenges in using such a conventional method is that the conidia of *M. grisea* are not always present on the infected leaf tissue during microscopic examination. Symptomatic leaves may require incubation in a moist chamber for up to 48 h to induce sporulation. Recently, a rapid immuno-recognition assay using a monoclonal antibody was developed for diagnosis of gray leaf spot (53). This method of detection has a great potential for practical use as a diagnostic kit in processing turf

samples. In this assay, the monoclonal antibody did not react with antigens from *Rhizoctonia solani* (brown patch), *Bipolaris sorokiniana* (Bipolaris leaf spot), or *Pythium aphanidermatum* (Pythium blight). These pathogens are prevalent in perennial ryegrass during the periods of gray leaf spot epidemic development, and the symptoms they cause often resemble those of gray leaf spot (Fig. 7). The monoclonal antibody was effective in detecting *M. grisea* in symptomatic perennial ryegrass leaves, but not during the incubation period. Although significant advances have been made in polymerase chain reaction (PCR)-based procedures for diagnosis of a number of turfgrass pathogens in recent years, this procedure is not yet available for diagnosing gray leaf spot.

Disease Development

M. grisea overwinters as dormant mycelium in dead plant debris. Harmon and Latin (18) found that survival of *M. grisea* was greatly reduced during the winter, but they successfully induced sporulation of the fungus from infested plant debris in the spring. Conidia produced from the leaf debris apparently serve as the primary inoculum for leaf infections early in the growing season, although details of this early infection process need to be determined. It is also possible that at least some infection foci are established via long-distance dispersal of conidia. Based on field observations, we hypothesize that gray leaf spot develops at visually undetectable levels in early to mid-summer. Conidia produced on infected leaves during this period eventually trigger a series of secondary infections that contribute to the build-up of inoculum during the late summer periods (Fig. 8).

Gray leaf spot is often observed first in turf in golf course roughs maintained at higher mowing heights (Fig. 9). These areas may be partially shaded and have extended leaf wetness periods and high humidity that are more conducive for infection. The disease may be detected in roughs several days before extensive damage of turf in fairways becomes evident. Dispersal of inoculum is by wind, wind-blown rain, water-splash from sprinkler irrigation, movement by ground maintenance equipment, and other golfing activities. Dispersal of conidia by mowers, spray rigs, spreaders, core aerifiers, and golf carts is important in the spread of the disease in golf courses. Although transmission of *M. grisea* by seeds is well known in rice blast disease, this has not been documented in perennial ryegrass.

Gray leaf spot normally develops from early August to mid-October. Environmental conditions prevailing during this late summer period and availability of inoculum are major determinants in the development of gray leaf spot epidemics. Efforts to quantitatively describe the relationships between environmental factors and gray leaf spot development were first studied on Italian ryegrass in 1987 (27) and continued on tall fescue (41) and perennial ryegrass (22). These studies showed that temperatures between 26 and 29°C were optimal for disease development. A recent study on effects of environmental factors on development of gray leaf spot on perennial ryegrass showed a similar temperature effect on disease development (44). The study showed that the effect of temperature on gray leaf spot incidence and severity is cubic, indicating that disease incidence and severity increase with an increase in temperature from 20 to 28°C and decrease with an increase in temperature above 28°C. Leaf wetness duration is also important in the development of gray leaf spot (41,44). Uddin et al. (44) reported that

Fig. 2. Incidence and severity of gray leaf spot or blast on tall fescue, hard red winter wheat, and perennial ryegrass inoculated with isolates of *Magnaporthe grisea* from perennial ryegrass.

disease incidence and severity increased with increased leaf wetness duration at all temperatures (Fig. 10). A shorter leaf wetness duration was required for disease development under warmer temperatures. They proposed a polynomial regression model, $Y_{is} = b_0 + b_1T + b_2T^2 + b_3T^3 + b_4W + b_5W^2 + b_6TW$ (Y_{is} = disease incidence or disease severity, T = temperature, and W = leaf wetness duration in hours), as part of a forecasting system for predicting the disease.

In addition to leaf wetness duration, relative humidity is also an important environmental component influencing gray leaf spot development. Although expansion of necrotic lesions is rapid under prolonged leaf wetness periods, conidia are not produced when excessive free moisture is present on the leaf tissue. Removal of free moisture from the infected leaf blades under high humidity is necessary for production of conidia. Therefore, warm day and night temperatures, subsequent wetting and drying of leaf blades, and high humidity regimes are major contributing factors to the development of gray leaf spot epidemics and perpetuation of the disease. Further studies on epidemiology, particularly the role of relative humidity in gray leaf spot epidemic development and testing of the model under various turfgrass cultural management practices, are warranted.

In turfgrass management, the assessment of damage caused by gray leaf spot is exclusively based on disease severity measured by the percent area of dead turf in fairways. Such an assessment may be impractical when the disease appears as off-color, diffused blight at its early stage of development. At this stage when the disease is initially detected, incidence data (percent leaf blades symptomatic) can be obtained and the severity estimated; thus a disease threshold for a fungicide spray program can be established. A recent study has shown that gray leaf spot severity can be estimated by incidence ($Y = 0.25 - 0.002x + 0.0009x^2$; $r^2 = 0.90$, where Y = disease severity, and x = disease incidence) (44).

Disease Management Strategies

Cultural management practices often do not provide adequate control of gray leaf spot due to rapid development of the disease and high susceptibility of currently available cultivars. An integrated approach that entails various cultural management practices and a sound fungicide program provide effective control of gray leaf spot.

Fungicidal control. Among currently labeled fungicides, the most effective materials (Fig. 11) for gray leaf spot control are azoxystrobin, trifloxystrobin, and thiophanate methyl. Azoxystrobin is labeled for gray leaf spot as Heritage 50WG at the rates of 0.61 to 1.22 kg of formulated product per hectare (0.2 to 0.4 oz/1,000 ft²) at 14- to 28-day intervals. While the 0.61

kg/ha rate of Heritage 50WG has proven very effective in some tests (11,36–38,45,50,72), biweekly applications of the 0.61 kg/ha rate sometimes have resulted in a small but significant amount of foliar blighting (12,39,56,69). In contrast, application of Heritage 50WG at the 1.22 kg/ha rate typically has provided excellent disease control for at least 3 weeks under high disease pressure (45,67,69–71). The performance of trifloxystrobin (formulated as Compass 50WG) in field tests typically has been very satisfactory. In some studies, there was no significant ($P \geq 0.05$) difference between Compass used at labeled rates at 2-week intervals and the top-performing treatment in the test (12,38,45, 63,70). However, in several tests, use of the compound according to label directions

provided slightly lower disease control than the top treatment in the test (56,65). The product has not always provided acceptable disease control when used at labeled intervals exceeding 2 weeks, even at the highest labeled rate (65). Thiophanate methyl (Cleary's 3336 50WP) and similar products typically have provided excellent control under high disease pressure when used at a minimum of 9.15 kg a.i./ha at 14-day intervals (37,50,56,70). Rates as low as 6.1 kg a.i./ha have been effective under low to moderate disease pressure (38,39,63). In one test, thiophanate methyl applied biweekly at 9.15 kg a.i./ha provided excellent control for most of the season, but control diminished somewhat at the end of the epidemic (11), suggesting that under the highest disease pressure, the

Fig. 3. Gray leaf spot symptoms on perennial ryegrass. A, Characteristic necrotic lesion. B, Flagging of leaf blade.

Fig. 4. Extensive damage of perennial ryegrass on a golf course fairway caused by gray leaf spot.

Fig. 5. Gray leaf spot development in drainage areas where high relative humidity and prolonged leaf wetness periods occur.

Fig. 6. Cryo-scanning electron micrograph of A, conidia, B, germinating conidium with appressorium, and C, fluorescent micrograph of conidia of *Pyricularia grisea* causing gray leaf spot of perennial ryegrass turf.

12.2 kg a.i./ha rate may be necessary. Formulations of two demethylation inhibitor (DMI) fungicides, propiconazole and triadimefon, are labeled for gray leaf spot. These DMI fungicides, when used alone following label directions, have usually provided poor control under high disease pressure (36–38,45,56,68–70,73). Chlorothalonil and mancozeb, both contact fungicides that act as nonspecific enzyme inhibitors, are also labeled for this disease. Although some studies have shown good results with these materials (12,38,39,48, 50,56,72), chlorothalonil (11,36,37,65, 69,70) and mancozeb (56,68,73) have not consistently provided acceptable control

under high disease pressure. Tank mixes of a DMI fungicide and chlorothalonil at labeled rates can often provide excellent control (11,12,38,48,67,70), but control is sometimes not complete under high disease pressure (65,68).

Timely use of fungicides can help prevent the disease epidemic from reaching its logarithmic phase. Although infections of *M. grisea* can be found as much as 4 to 6 weeks in advance of the logarithmic phase, rapid increase in disease development must be prevented (66,71). In the absence of site-specific information on disease development, some turf managers will initiate a spray program a week or two before the time of year when epidemics historically have begun in the area; others will begin spraying when the disease is first reported in the region. Both approaches carry risks. The frequency and longevity of a spray program often depend on a combination of the past history of the disease at the site and in the region, and the weather conditions favorable to disease development. In the Midwest and Northeast, the window where fungicide protection is needed is usually from early August to early September and often beyond. In seasons with low disease pressure, no fungicide protection may be needed beyond early September (66), whereas under high disease pressure, fungicides are needed into October (71).

Fungicide resistance. The three most effective active ingredients against gray leaf spot belong to two fungicide groups with a high risk for selection of fungicide-resistant subpopulations. Azoxystrobin and trifloxystrobin are strobilurins in a relatively new fungicide class, the QI fungicides, to which several pathogens have developed resistance (3); and thiophanate methyl is a benzimidazole fungicide, a group in which highly resistant pathogen subpopulations have been selected rela-

Fig. 7. Gray leaf spot symptoms resemble those of A, brown patch, and B, Pythium blight.

tively frequently (34). To date, *M. grisea* isolates from perennial ryegrass with resistance to Q₀I fungicides have been found in a total of nine locations in five states (Ken-

tucky, Illinois, Indiana, Maryland, and Virginia) since azoxystrobin was first commercially available for gray leaf spot control in 1998. In cases where we have access

to records, the disease outbreaks occurred after Q₀I fungicides had been applied according to label directions. Resistant isolates have exhibited significantly reduced

Fig. 8. Life cycle of *Pyricularia grisea*.

sensitivity to azoxystrobin in vitro compared with baseline isolates (64; P. Vincelli and E. Dixon, *unpublished*), and they have one or the other of the target site mutations in the mitochondrial cytochrome *b* gene associated with resistance to Q_oI fungicides (3,21).

Although there appears to be a significant resistance risk to Q_oI fungicides in *M. grisea* isolates from perennial ryegrass, these materials should continue to have a place in control programs for gray leaf spot. Rapid development of a resistant subpopulation in the field does not necessarily preclude the effective use of the fungicide in other locations for a number of years. For example, resistance in strains

of *Podosphaera xanthii* to benomyl and to DMI fungicides occurred 1 and 2 years after commercial introduction, respectively, and yet these fungicides remained important disease control tools in other areas for years until resistant strains became widespread (26). The Q_oI fungicides have been an important component of a gray leaf spot management program since 1998, when azoxystrobin received a federal registration for the disease. Avoiding the use of Q_oI fungicides in the many locations with no evidence of control failure would exert additional selection pressure on *M. grisea* toward resistance to the few remaining effective fungicides, e.g., thiophanate-methyl and DMI fungicides (62). The po-

tential for rapid selection for resistance does, however, suggest that users of Q_oI fungicides for gray leaf spot control should aggressively use strategies for reducing the risk of resistance.

Biological control. Significant efforts have been made in development of biological control that can be incorporated into a broader integrated management strategy for gray leaf spot (52). A recent study has shown that *Pseudomonas aeruginosa* isolated from spent mushroom substrate is highly antagonistic to the gray leaf spot pathogen (60). In controlled environmental chamber studies, bacterial isolates provided a level of control that was comparable to that of propiconazole but significantly lower than that of azoxystrobin. In field experiments, *P. aeruginosa* provided significant suppression of gray leaf spot in perennial ryegrass turf with efficacy up to 14 days when applied as a preventive (60). Another bacterium, *Bacillus lentimorbus*, isolated from the rhizosphere of perennial ryegrass has also been shown to be suppressive to *M. grisea* (58). In dual plate assays, the antagonist significantly inhibited mycelial growth of *M. grisea* isolates from various hosts including perennial ryegrass, tall fescue, and rice. In controlled environment chamber experiments, the bacterium significantly reduced disease severity and incidence with efficacy up to 20 days when applied as a preventive. In addition to bacteria, various pathogenic and nonpathogenic isolates of *R. solani* have been shown to be suppressive to gray leaf spot (46). Further studies addressing the safety and practical use of the biocontrol agents are warranted.

Turfgrass height and grass clipping management. Results from experiments on the effects of mowing height on gray leaf

Fig. 9. Development of gray leaf spot in rough (far right) and subsequent spread of the disease to fairways.

Fig. 10. A surface response for disease incidence and severity derived from the polynomial model, $Y_{is} = b_0 + b_1T + b_2T^2 + b_3T^3 + b_4W + b_5W^2 + b_6TW$, where Y_{is} = disease incidence or disease severity, T = temperature, and W = leaf wetness duration in hours.

spot severity have been contradictory. A study conducted in the northeastern region of the United States indicated that gray leaf spot was more severe at a mowing height of 8.9 cm than of 1.27 cm (55). In Kentucky, Williams et al. (74) found no differences in disease severity on perennial ryegrass mowed at 1.9 cm and 6.4 cm. This discrepancy may have been due to various cultural and environmental factors influencing disease development. Turf managers in the Northeast and Mid-Atlantic States are generally advised to avoid raising mowing heights, particularly during gray leaf spot epidemics. Grass clipping management is also important, as the removal of clippings can reduce disease incidence substantially under low disease intensity; however, under high disease intensity the effect of clipping removal is not significant (7,55). Although clipping removal effectively reduces disease intensity under low to moderate disease intensity, collecting the clippings from large fairways and disposing them is impractical in the operation of most golf courses.

Fertility. Increasing the amount nitrogen increases gray leaf spot severity (51,55,74). Source of nitrogen also influences gray leaf spot development. A recent study has shown that gray leaf spot severity was lower when controlled-release forms of nitrogen such as isobutylidene diurea (IBDU) and sewage sludge-based Milorganite were applied compared to quick-release forms such as ammonium nitrate and urea (51).

Herbicide. The herbicide ethofumesate is widely used to control annual bluegrass in perennial ryegrass fairways. Its use in the spring has been associated with in-

creased severity of gray leaf spot (47). These effects were not evident when the herbicide was applied during fall. The mechanism for the increase in severity of gray leaf spot by ethofumesate is unclear. The herbicide interferes with fatty acid biosynthesis in plants (13) and causes aggregation of the epicuticular wax on leaves (23). More efficient penetration of host plant surface at the thinning areas of leaf tissue by *M. grisea* may have resulted in higher gray leaf spot severity (Fig. 12; W. Uddin and M. D. Soika, *unpublished*). A more rigorous fungicide spray program for gray leaf spot may be required in areas where ethofumesate is applied in spring for annual bluegrass seedhead suppression (47).

Breeding for resistance. Cultivars of perennial ryegrass resistant to the gray leaf spot pathogen are not currently available. Polygenic resistance appears to be present in perennial ryegrass (9). A number of field studies to identify novel sources of resistance in a worldwide collection of perennial ryegrass cultivars and lines are currently underway. Thus far, some progress in identifying resistant germplasm has been made, and several improved lines and cultivars have been identified (5,17). Research to identify patterns of inheritance and DNA markers associated with resistance in progenies from cultivars, experimental selections, and single-plot progenies of perennial ryegrass are in progress. Recently, microsatellite DNA markers in perennial ryegrass progenies associated with moderate to good resistance to *M. grisea* have been identified (4). Breeding for gray leaf spot resistance in perennial

ryegrass still remains a long-term goal in the turfgrass industry.

Current Status and Future Outlook

The impact of gray leaf spot on the management of perennial ryegrass in golf courses has been profound. Additionally, the disease is an emerging problem on ryegrass turf in residential lawns and sports and recreational turf. Although the economic loss due to damage in residential and sports turf has not been well documented, the impact of gray leaf spot in these turf types appears to be significant. As gray leaf spot emerges as a serious disease of perennial ryegrass turf, significant progress in research on various aspects of the disease, such as biology of the pathogen and epidemiology and management of the disease, has been made in recent years. The genetic relationship among the isolates of *M. grisea* from perennial ryegrass and other gramineous hosts has provided basic understanding of the biology of the pathogen (15,61). Development of a weather-based model for predicting gray leaf spot epidemic has been a focus of our research, and currently a temperature and leaf wetness duration based model is available (44). Efforts to quantitatively describe the relationship between the relative humidity and gray leaf spot development are currently underway. A major breakthrough in research on gray leaf spot management is the development of effective fungicide programs for the disease. While intensive use of fungicides is not a desirable long-term disease control strategy, spray programs can be used to prevent epidemics until more sustainable management options are available.

Fig. 11. Efficacy of various fungicides for control of gray leaf spot. Azoxystrobin, trifloxystrobin, and thiophanate methyl provided excellent control (plots in background), and chlorothalonil provided moderate control (plot with small patches in left). Flutolanil and propiconazole provided poor control (plots in right) which was not different from nontreated control plot (foreground).

Fig. 12. Effect of ethofumesate on epicuticular wax deposition on perennial ryegrass leaf blade. A, Cryo-scanning electron micrograph of perennial ryegrass leaf blade 4 weeks after treatment with ethofumesate. B, Control (distilled water).

While significant advances have been made in understanding gray leaf spot of perennial ryegrass turf, some major challenges still exist. One of the most serious challenges is the identification of resistant germplasms and development of resistant cultivars. Additionally, very little is known about the life cycle of the gray leaf spot pathogen. Further understanding of the life cycle and its significance to the epidemic development in ryegrass fairways in late summer periods will require major efforts in gray leaf spot research. Although there may be conclusive evidence that the teleomorph of the rice blast fungus is *M. oryzae* as suggested by a recent study (8), genetic relationships between the pathogens of rice blast and gray leaf spot disease of perennial ryegrass need to be examined more closely. Phylogenetic analysis of a large pool of isolates of *P. grisea* from perennial ryegrass from a diverse population may be required to reach a reasonable conclusion that the teleomorph of the gray leaf spot pathogen is indeed *M. oryzae*. Additionally, determination of the population structure of the gray leaf spot pathogen that encompasses wider geographic regions of the United States will require development of a database on *M. grisea* isolates to support such an undertaking.

Acknowledgments

We thank Michael Soika, Frances Moorman, Ed Dixon, Jack Doney, Paul Burrus, David Williams, Yun-sik Kim, Jesse Shelton, Nasreen Kabir, Erin Soika, Lesli Lucas, and Billie Lucas for providing technical assistance in studies that have provided information for this article. Financial support for these studies was provided by the Golf Course Superintendents' Association of America, United States Golf Association, The Pennsylvania Turfgrass Council, and the fungicide industry.

Literature Cited

- Asuyama, H. 1963. Morphology, taxonomy, host range, and life cycle of *Piricularia oryzae*. Pages 9-22 in: The Rice Blast Disease. S. H. Ou, ed. John Hopkins Press, Baltimore, MD.
- Bain, D. C., Patel, B. M., and Patel, M. V. 1972. Blast of ryegrass in Mississippi. Plant Dis. Rep. 56:210.
- Bartlett, D. W., Clough, J. M., Godwin, J. R., Hall, A. A., Hamer, M., and Parr-Dobrzanski, B. 2002. The strobilurin fungicides. Pest Manage. Sci. 58:649-662.
- Bonos, S. A., Kubik, C., Clarke, B. B., and Myer, W. A. Breeding perennial ryegrass for resistance to gray leaf spot. Crop Sci. In press.
- Bonos, S. A., Kubik, C., Han, Y., and Meyer, W. A. 2001. Evaluation of perennial ryegrass germplasm for gray leaf spot. Agron. Abstr. American Society of Agronomy, Madison, WI.
- Carver, R. B., Rush, M., and Lindberg, G. D. 1972. An epiphytotic of ryegrass blast in Louisiana. Plant Dis. Rep. 56:157-159.
- Clarke, B. B., and Vaiciunas, S. S. 2001. Best management practices for control of gray leaf spot. (Abstr.) Phytopathology 91:S148.
- Couch, B. C., and Kohn, L. M. 2002. A multi-locus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. Mycologia 94:683-693.
- Curley, J., Jung, G., Warnkes, S., Barker, R., and Leong, S. 2001. QTL mapping for resistance to gray leaf spot in perennial ryegrass. (Abstr.) Phytopathology 91:S20.
- Dernoe, P. H. 1996. Perennial ryegrass and gray leaf spot. Golf Course Manage. 64:49-52.
- Dernoe, P. H. 2001. Gray leaf spot control with fungicides, 1998. Fungic. Nematicide Tests 56:T32.
- Dernoe, P. H., Krouse, J. M., and Kaminski, J. E. 2001. Gray leaf spot control with fungicides, 2000. Fungic. Nematicide Tests 56:T31.
- Devine, M. D., Duke, S. O., and Fedtke, C. 1993. Physiology of Herbicide Action. Prentice Hall, Englewood Cliffs, NJ.
- Ellis, M. B. 1993. Dematiaceous Hyphomycetes. CAB International, Wallingford, UK.
- Farman, M. L. 2002. *Piricularia grisea* isolates causing gray leaf spot on perennial ryegrass (*Lolium perenne*) in the United States: Relationship to *P. grisea* isolates from other host plants. Phytopathology 92:245-254.
- Farr, D. F., Bills, G. F., Chamuris, G. P., and Rossman, A. Y. 1989. Fungi on Plants and Plant Products in the United States. American Phytopathological Society, St. Paul, MN.
- Hamblin, A. M., and Hofmann, N. E. 2001. Breeding for gray leaf spot resistance. (Abstr.) Phytopathology 91:S148.
- Harmon, P. F., and Latin, R. 2001. Perennation of *Magnaporthe grisea* in the Midwest: Disease management implications. (Abstr.) Phytopathology 91:S148.
- Harmon, P., Rane, K., Ruhl, G., and Latin, R. 2000. First report of gray leaf spot on perennial ryegrass in Indiana. Plant Dis. 84:492.
- Hebert, T. T. 1971. The perfect stage of *Piricularia grisea*. Phytopathology 61:83-87.
- Kim, Y.-S., Dixon, E. W., Vincelli, P., and Farman, M. L. 2003. Field resistance to strobilurin (Q_0) fungicides in *Piricularia grisea* caused by mutations in the mitochondrial cytochrome b gene. Phytopathology 93:891-900.
- Landschoot, P. J., and Hoyland, B. F. 1992. Gray leaf spot of perennial ryegrass turf in Pennsylvania. Plant Dis. 76:1280-1282.
- Levitt, J. R. C., and Penner, D. 1978. A comparison of the effects of EPTC and R-25788 (2,2-dichloro-N,N-diallyl acetamide) on epicuticular wax deposition on corn and cabbage. (Abstr.) Weed Sci. Soc. Am. 79.
- Malca, I., and Owen, J. H. 1957. The gray leaf spot disease of St. Augustinegrass. Plant Dis. Rep. 41:871-875.
- McCarty, B. 2000. Perennial ryegrass on golf courses: Friend or foe? Grounds Maint. Jan. 67-74.
- McGrath, M. T. 2001. Fungicide resistance in cucurbit powdery mildew: Experiences and challenges. Plant Dis. 85:236-245.
- Moss, M. A., and Trevathan, L. E. 1987. Environmental conditions conducive to infection of ryegrass by *Piricularia grisea*. Phytopathology 77:863-866.
- Ou, S. H. 1985. Rice Diseases. Commonwealth Mycological Institute, Kew, Surrey, UK.
- Pappas, A. C., and Paplomatas, E. J. 1998. *Piricularia* leaf spot: A new disease of ornamental plants of the family Marantaceae. Plant Dis. 82:465-469.
- Pedersen, D. K., Kane, R. T., and Wilkinson, H. T. 2000. First report of gray leaf spot caused by *Piricularia grisea* on *Lolium perenne* in Illinois. Plant Dis. 84:1151.
- Rossman, A. Y., Howard, R. J., and Valent, B. 1990. *Piricularia grisea*, the correct name for the rice blast fungus. Mycologia 82:509-512.
- Schumann, G. L., and Jackson, N. 1999. First report of gray leaf spot (*Piricularia grisea*) on perennial ryegrass (*Lolium perenne*) in New England. Plant Dis. 83:1073.
- Smiley, R. W., Dernoe, P. H., and Clarke, B. B. 1996. Compendium of Turfgrass Diseases. American Phytopathological Society, St. Paul, MN.
- Smith, C. M. 1988. History of benzimidazole use and resistance. Pages 23-24 in: Fungicide Resistance in North America, C. J. Delp, ed. American Phytopathological Society, St. Paul, MN.
- Sundaram, N. V., Palmer, L. T., Nagarajan, K., and Prescott, J. M. 1972. Disease survey of sorghum and millet in India. Plant Dis. Rep. 56:740-743.
- Tisserat, N., and Fry, J. 1999. Preventive fungicide applications for control of gray leaf spot on perennial ryegrass, 1998. Fungic. Nematicide Tests 54:518.
- Towers, G. W., Majumdar, P. R., Weibel, E. N., DiMarco, J. N., Peacos, M., and Clarke, B. B. 2000. Evaluation of fungicides for the control of gray leaf spot, 1999. Fungic. Nematicide Tests 55:519.
- Towers, G. W., Majumdar, P. R., Weibel, E. N., DiMarco, J. N., Peacos, M., and Clarke, B. B. 2000. Efficacy of fungicides for control of gray leaf spot on perennial ryegrass, 1999. Fungic. Nematicide Tests 55:520-521.
- Towers, G. W., Majumdar, P. R., Weibel, E. N., DiMarco, J. N., Peacos, M., and Clarke, B. B. 2000. Evaluation of selected fungicides for the control of gray leaf spot, 1999. Fungic. Nematicide Tests 55:522.
- Trevathan, L. E., Moss, M. A., and Blasingame, D. 1994. Ryegrass Blast. Plant Dis. 78:113-117.
- Uddin, W., Burpee, L. L., and Stevenson, K. L. 1998. Influence of temperature, and leaf wetness duration on development of gray leaf spot (blast) of tall fescue. Pages 136-141 in: Int. Conf. Plant Prot. Tropics, 5th. Kuala Lumpur, Malaysia.
- Uddin, W., Moorman, F. E., Soika, M. D., and Viji, G. 1999. Influence of turfgrass age on development of gray leaf spot (blast) of tall fescue. (Abstr.) Phytopathology 89:S78.
- Uddin, W., Schumann, G. L., Viji, G., and Boyd, S. H. 2001. Monitoring of gray leaf spot epidemics in golf courses in New England, New Jersey, and Pennsylvania in 2000. (Abstr.) Phytopathology 91:S198.
- Uddin, W., Serlemitsos, K., and Viji, G. 2003. A temperature and leaf wetness duration-based model for prediction of gray leaf spot of perennial ryegrass turf. Phytopathology 93:336-343.
- Uddin, W., and Shelton, J. 2002. Evaluation of fungicides for control of gray leaf spot on perennial ryegrass, 2001. Fungic. Nematicide Tests 57:T34.
- Uddin, W., Shelton, J., and Viji, G. 2002. Interactions between *Piricularia grisea* and *Rhizoctonia solani* in gray leaf spot pathosystem in perennial ryegrass turf. (Abstr.) Phytopathology 92:S82.
- Uddin, W., and Soika, M. D. 2000. Effects of plant growth regulators, herbicides, and fungicides on development of blast disease (gray leaf spot) of perennial ryegrass turf. (Abstr.) Phytopathology 90:S78.
- Uddin, W., and Soika, M. D. 2000. Evaluation of fungicides for control of gray leaf spot (blast) on perennial ryegrass fairway, 1999. Fungic. Nematicide Tests 55:526.
- Uddin, W., Soika, M. D., Moorman, F. E., and Viji, G. 1999. A serious outbreak of blast disease (gray leaf spot) of perennial ryegrass in golf course fairways in Pennsylvania. Plant Dis. 83:783.
- Uddin, W., Soika, M. D., Shelton, J., and Soika, E. L. 2001. Evaluation of fungicides for control of gray leaf spot (blast) on a perennial ryegrass fairway, 2000. Fungic. Nematicide Tests 56:T40.
- Uddin, W., Soika, M. D., Shelton, J., and Soika, E. L. 2001. Severity of gray leaf spot (blast) of perennial ryegrass turf as influenced by nitrogen source. (Abstr.) Phytopathology 91:S90.
- Uddin, W., and Viji, G. 2002. Biological

- control of turfgrass diseases. Pages 313-337 in: Biological Control of Crop Diseases. S. S. Gnanamanickam, ed. Marcel Dekker, Inc., New York.
53. Uddin, W., Viji, G., Schumann, G. L., and Boyd, S. H. 2003. Detection of *Pyricularia grisea* causing gray leaf spot of perennial ryegrass turf by a rapid immuno-recognition assay. Plant Dis. 87:772-778.
 54. Uddin, W., Viji, G., and Stowell, L. 2002. First report of gray leaf spot on perennial ryegrass turf in California. Plant Dis. 86:75.
 55. Vaiciunas, S., and Clarke, B. B. 2000. Impact of cultural practices and genotype on the development of gray leaf spot in cool season turfgrasses. Proc. Rutgers Turfgrass Sympos., New Brunswick, NJ.
 56. Vaiciunas, S. S., Majumdar, P. R., DiMarco, J. N., Tredway, L. P., Peacos, M., and Clarke, B. B. 1999. Curative control of gray leaf spot with selected fungicides, 1998. Fungic. Nematicide Tests 54:522-523.
 57. Vargas, J. M., Jr. 1994. Management of Turfgrass Diseases. Lewis Publishers, Ann Arbor, MI.
 58. Viji, G., and Uddin, W. 2001. Biocontrol of gray leaf spot (blast) of perennial ryegrass by *Bacillus lentimorbus*. (Abstr.) Phytopathology 91:S92.
 59. Viji, G., and Uddin, W. 2002. Distribution of mating type alleles and fertility status of *Magnaporthe grisea* causing gray leaf spot of perennial ryegrass and St. Augustinegrass turf. Plant Dis. 86:827-832.
 60. Viji, G., Uddin, W., and Romaine, C. P. 2003. Suppression of gray leaf spot of perennial ryegrass turf by *Pseudomonas aeruginosa* from spent mushroom substrate. Biol. Control 26:233-243.
 61. Viji, G., Wu, B., Kang, S., Uddin, W., and Huff, D. R. 2001. *Pyricularia grisea* causing gray leaf spot of perennial ryegrass turf: Population structure and host specificity. Plant Dis. 85:817-826.
 62. Vincelli, P. 2000. Fungicidal control of gray leaf spot. Golf Course Manage. 68:68-74.
 63. Vincelli, P., and Dixon, E. 2001. Efficacy and timing of fungicides for control of gray leaf spot of perennial ryegrass, 2000. Fungic. Nematicide Tests 56:T37.
 64. Vincelli, P., and Dixon, E. 2002. Resistance to Q_oI (Strobilurin-like) fungicides in isolates of *Pyricularia grisea* from perennial ryegrass. Plant Dis. 86:235-240.
 65. Vincelli, P., Dixon, E., and Williams, D. 2000. Efficacy of fungicides for control of gray leaf spot of perennial ryegrass, 1999. Fungic. Nematicide Tests 55:529.
 66. Vincelli, P., Dixon, E., and Williams, D. 2000. Timing of fungicides for control of gray leaf spot of perennial ryegrass, 1999. Fungic. Nematicide Tests 55:531.
 67. Vincelli, P., Dixon, E., and Williams, D. 2001. Efficacy of fungicides for control of gray leaf spot of perennial ryegrass, 2000. Fungic. Nematicide Tests 56:T38.
 68. Vincelli, P., Dixon, E., Williams, D., and Burrus, P. 2002. Efficacy of fungicides for control of gray leaf spot of perennial ryegrass, 2001. Fungic. Nematicide Tests 57:T33.
 69. Vincelli, P., and Doney, J. C. 1997. Evaluation of fungicides for control of gray leaf spot of perennial ryegrass, 1996. Fungic. Nematicide Tests 52:398.
 70. Vincelli, P., and Williams, D. W. 1999. Efficacy of fungicides for control of gray leaf spot of perennial ryegrass, 1998. Fungic. Nematicide Tests 54:524-525.
 71. Vincelli, P., and Williams, D. W. 1999. Timing of Heritage 50WG fungicide for control of gray leaf spot of perennial ryegrass, 1998. Fungic. Nematicide Tests 54:526.
 72. Watkins, J. E. 2000. Evaluation of fungicides for control of gray leaf spot of perennial ryegrass, 1999. Fungic. Nematicide Tests 55:534.
 73. Watkins, J. E. 2001. Fungicides evaluated for control of gray leaf spot, 2000. Fungic. Nematicide Tests 56:T35.
 74. Williams, D. W., Burrus, P. B., and Vincelli, P. 2001. Severity of gray leaf spot in perennial ryegrass as influenced by mowing height and nitrogen level. Crop Sci. 41:1207-1211.
 75. Yaegashi, H., and Udagawa, S. 1978. The taxonomical identity of the perfect state of *Pyricularia grisea* and its allies. Can. J. Bot. 56:180-183.

Wakar Uddin

Dr. Uddin is an assistant professor in the Department of Plant Pathology at the Pennsylvania State University. He received a B.S. in plant science and an M.S. in IPM from University of Nevada-Reno in 1985 and 1987, respectively, and a Ph.D. in plant pathology from the University of Georgia in 1996. After receiving the M.S., he worked as a research assistant at the Columbia Basin Agricultural Research Center, Oregon State University, and as a diagnostician at the Plant Disease Clinic at the University of Georgia. He spent a year as a postdoctoral research associate at the University of Georgia in 1997 prior to joining the Pennsylvania State University in 1998. His research involves epidemiology and management of gray leaf spot of perennial ryegrass turf, and population biology of *Magnaporthe grisea*. He teaches two resident courses and an online course (Penn State World Campus) in turfgrass disease diagnosis and management.

Gnana Viji

Dr. Viji is a postdoctoral research associate in the Department of Plant Pathology at the Pennsylvania State University. She received her B.Sc. in 1987, M.Sc. in 1989, and Ph.D. in 1997 in botany from the University of Madras, India. Her Ph.D. research involved genetic diversity and fertility status of *Magnaporthe grisea* causing blast disease of finger millet. She was a senior research fellow of the Rockefeller Foundation Rice Biotechnology International Fellowship from 1993 to 1997 and a visiting scholar at the International Rice Research Institute, Philippines, in 1995. Dr. Viji was also a lecturer at the Loyola College at Madras in India. Her current research focuses on the biology of fungal pathogens, particularly *M. grisea*, associated with gray leaf spot of perennial ryegrass turf with emphasis on population biology.

Paul Vincelli

Dr. Vincelli received his bachelor's degree in botany from Rutgers University. He served in the Peace Corps for 3 years in Nicaragua and Colombia, and then pursued graduate studies in plant pathology at Rutgers University (M.S.) and Cornell University (Ph.D.). He worked as the extension plant pathologist at the University of Wyoming during 1988-1989. He joined the faculty of the Department of Plant Pathology at the University of Kentucky in 1990 and was promoted to extension professor in 2001. His current responsibilities include extension and applied research for pathology of turfgrasses, forage crops, and corn; he also teaches the introductory course in plant pathology and a graduate course in epidemiology and management of plant diseases.