The Role of Hybridization in the Evolution and Emergence of New Fungal Plant Pathogens

Eva H. Stukenbrock

Environmental Genomics, Christian-Albrechts University of Kiel, Am Botanischen Garten 9-11, 24118 Kiel, Germany and Max Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, 24306 Plön, Germany.

Accepted for publication 13 November 2015.

ABSTRACT

Stukenbrock, E. H. 2016. The role of hybridization in the evolution and emergence of new fungal plant pathogens. *Phytopathology* 106:104-112.

Hybridization in fungi has recently been recognized as a major force in the generation of new fungal plant pathogens. These include the grass pathogen *Zymoseptoria pseudotritici* and the powdery mildew pathogen *Blumeria graminis triticales* of triticale. Hybridization also plays an important role in the transfer of genetic material between species. This process is termed introgressive hybridization and involves extensive backcrossing between hybrid and the parental species. Introgressive hybridization has contributed substantially to the successful spread of plant pathogens such as *Ophiostoma ulmi* and *O. novo-ulmi*, the causal agents of Dutch elm disease, and other tree pathogens such as the rust pathogen *Melampsora*. Hybridization occurs more readily between species that have previously not coexisted, so-called allopatric species. Reproductive barriers between allopatric species are likely to be more permissive allowing interspecific mating to occur. The bringing together of allopatric species of plant pathogens by global agricultural trade consequently increases the potential for hybridization between pathogen species. In light of global environmental changes, agricultural development, and the facilitated long-distance spread of fungal plant pathogens, hybridization should be considered an important mechanism whereby new pathogens may emerge. Recent studies have gained insight into the genetics and biology of fungal hybrids. Here I summarize current knowledge about hybrid speciation and introgressive hybridization. I propose that future studies will benefit greatly from the availability of large genome data sets and that genome data provide a powerful resource in combination with experimental approaches for analyses of hybrid species.

Additional keywords: genome evolution, genomic conflicts, host specificities.

Hybrids emerge from successful genetic crosses of non-conspecific individuals. In fungi, hybrids can be generated both by sexual mating and asexual fusion of cells or hyphae (Kohn 2005). Phylogenetic analyses can identify hybrids as intermediate clades with incongruent phylogenetic topologies that reflect the different evolutionary histories of different loci in the genomes of hybrids (Schardl and Craven 2003). Phenotypically hybrids are often characterized by intermediate phenotypes of the parental species (Greig et al. 2002).

Pure hybrids are found relatively rare in nature. One reason is that hybrids exhibiting intermediate phenotypes often have a reduced fitness compared with their parents. Due to their non-optimized phenotypes, they are outcompeted by their parents if co-existing in the same environment (Barton 2001). Additionally hybrids may suffer negative fitness effects due to genetic incompatibilities of

parental alleles. Many allelic combinations in the hybrid will consist of alleles that have not co-evolved in the same genomic background and therefore have not been optimized in parallel by natural selection. Such genomic incompatibilities in hybrids are also referred to as negative epistatic interactions or Dobzhansky-Muller interactions (Kondrashov et al. 2002; see also Kohn 2005 for discussion about the importance of Dobzhansky-Muller interactions in speciation of fungi). The finding of pure hybrid species in nature however suggests that genomic incompatibilities and fitness effects under some conditions can be overcome (Fig. 1A) (Greig et al. 2002; see also review by Yakimowski and Rieseberg 2014).

Another more common result of mating between non-conspecific individuals is introgression by which “transient” hybrids backcross with individuals of the parental species (Arnold 2004; Mallet 2005) (Fig. 1B). Repeated backcrossing can “dilute” the hybrid genome to only maintain fragments of the genome of the other parental species (Baack and Rieseberg 2007). Examples from plants, animals, and oomycetes suggest that introgressive hybridization can act to transfer adaptive traits between species and thereby be a mechanism that speeds up adaptive evolution (see Arnold 2004 for a review).

Corresponding author: E. H. Stukenbrock;
E-mail address: estukenbrock@bot.uni-kiel.de

<http://dx.doi.org/10.1094/PHYTO-08-15-0184-RVW>
© 2016 The American Phytopathological Society

Genomics have increased our understanding of hybrid genetics and have provided new insights into the importance of introgressive hybridization for species evolution (Mallet 2005). In a recent paper using comparative population genomics analyses, Lamichhaney and colleagues explored the adaptive radiation in one of the most famous models in evolutionary biology: Darwin's finches at the Galapagos Islands (Lamichhaney et al. 2015). The authors showed

evidence for extensive gene flow between species of finches. Clearly introgressive hybridization has played a role in adaptive radiation of finches together with natural selection on standing and introgressed variation. Adaptive introgression has also been identified in other species using genome analyses. Recent examples include *Heliconius* butterflies (Pardo-Diaz et al. 2012), monkey-flowers (Stankowski and Streisfeld 2015), and fungal plant

FIGURE 1

A, Hybridization can occur by asexual fusion or sexual mating between diverged species. **B**, Recurrent crossing between species and backcrossing of the hybrid with parental species is referred to as introgressive hybridization. In the extreme case of introgressive hybridization, the hybrid is a transient form, while the important outcome of the interspecific mating is the transfer of genetic material.

pathogens causing mildew (Menardo et al. in press). Together these studies demonstrate that hybridization in diverse organismal groups has been instrumental in the formation of adaptive diversity.

Among plant-associated fungi (pathogens, endophytes, or symbionts) there are several well-characterized hybrid systems including the grass endophyte *Epichloe* (Schardl 2001), the grass pathogen *Zymoseptoria pseudotritici* (Stukenbrock et al. 2012a), and the triticale infecting mildew pathogen *Blumeria graminis* f. sp. *triticales* (Menardo et al. in press). Moreover, introgressive hybridization has been documented in several systems such as the Dutch Elm disease pathogens *Ophiostoma ulmi* and *O. novo-ulmi* (Brasier et al. 1998; Brasier and Kirk 2010) and the anther smut pathogens *Microbotryum silene-dioicae* and *M. lychnidis-dioicae* (Gladieux et al. 2011).

Hybridization has been proposed a major force in the evolution of plant pathogens and in particular the emergence and adaptation of crop infecting pathogens (Brasier 2000, 2001). Global trade moves pathogens and hosts around the world and thereby brings together species, which previously did not co-exist (Brown and Hovmøller 2002). Introduced pathogens provide a threat to naïve hosts (cultivated as well as wild plants). Another concern is the potential of hybridization between introduced and native pathogen species. As will be discussed in more details below, reproductive barriers between species that have previously not co-existed can be more permissive and thereby allow hybrids to be formed more readily (Anderson et al. 1980; Dettman et al. 2003; Stenlid and Karlsson 1991). The result can be either the emergence of new hybrid pathogen species or the transfer of genetic material between pathogen species.

In this review, I will present examples of hybrid fungal plant pathogens and discuss the genetic basis of hybrid formation in the context of adaptive evolution. I propose that more studies integrating genomic and experimental data are essential to understand the biology and evolution of hybrids fungal plant pathogens in natural environments as well as agroecosystems.

HOW HYBRIDS ARISE

A hybrid is the offspring of two non-conspecific individuals (Mallet 2007). Normally, reproductive barriers act to maintain the genetic integrity of species and prevent the introduction of “foreign” DNA by mating with non-conspecific individuals (Orr 1995). It may however occur that hybrids can form if reproductive barriers are permissive, allowing mating between individuals of different species. To understand how reproductive barriers can be permissive, we first need to identify the underlying genetics of reproductive barriers. Detailed studies in the two ascomycete model species *Saccharomyces cerevisiae* and *Neurospora crassa* have shed light on speciation genetics and reproductive isolation in fungi (Dettman et al. 2007, 2008; Turner et al. 2011). These studies have documented how both prezygotic and postzygotic reproductive barriers either impair the formation of hybrid zygotes (prezygotic reproductive barriers) or severely decrease fitness of hybrid offspring (postzygotic reproductive barriers). Dettman and colleagues used an experimental evolution approach to generate two diverged yeast populations adapted to distinct environments, high salinity (S) and low glucose minimal (M) media (Dettman et al. 2007). As expected, fitness of SM hybrids was severely reduced compared with fitness of the parental populations. However, the hybrids were also impaired in meiotic efficiency suggesting additional genomic “defects.” The meiotic impairment could be linked to antagonistic interactions between diverged alleles brought together in the SM hybrids. Dettman and colleagues demonstrated how rapidly (500 cell transfers) negative interactions can arise between diverging populations. Moreover, the study shows that one or few genetic incompatibilities may be sufficient to prevent mating between diverged populations and thereby be the drivers of incipient speciation.

Turner and colleagues took a different approach to assess the underlying genetic traits of incompatibility between sympatric strains of *N. crassa* and *N. intermedia* (Turner et al. 2011). The authors generated hybrids of allopatric and sympatric strains of the two *Neurospora* species and used a quantitative trait locus (QTL) approach to identify the genetic traits associated with prezygotic reproductive barriers. By crossing sympatric and allopatric conspecific and non-conspecific strains, it was shown that hybrid fruiting bodies of sympatric crosses are aborted at a significantly higher frequency than hybrid fruiting bodies of allopatric species. This finding is in agreement with the presence of reinforced reproductive isolation between coexisting *N. crassa* and *N. intermedia* strains. Interestingly, maternal *N. crassa* colonies that aborted fruiting bodies in crosses with sympatric *N. intermedia* strains were subsequently able to mate normally with a conspecific partner. In comparison, a maternal *N. crassa* strain already fertilized by a conspecific partner (sympatric or allopatric) could not produce new fruiting bodies in a second round of fertilization with a conspecific partner. A key finding in the study by Turner and colleagues is the identification of genetic traits underlying fruiting body abortion and thereby the reinforced reproductive barriers (Turner et al. 2011). These major QTLs located on the mating type determining chromosome of *Neurospora* and associated with conspecific reproduction represent interesting candidates in future studies of speciation genomics of *Neurospora*.

Reinforced reproductive barriers between sympatric species have also been elegantly demonstrated by in vitro crossing experiments in other ascomycetes including wild populations of *S. cerevisiae* (Kuehne et al. 2007) and in Basidiomycetes such as *Armilaria mella* (Anderson et al. 1980) and *Heterobasidion annosum* (Stenlid and Karlsson 1991).

We know very little about reproductive barriers in plant-pathogenic fungi, although important insight has been gleaned from studies of the anther smut fungus *Microbotryum* (Gladieux et al. 2011; Le Gac et al. 2007b; Yockteng et al. 2008). Studies of *Microbotryum* have shown an effect of prezygotic reproductive barriers on hybridization between the two species *M. lychnidis-dioicae* and *M. silene-dioicae* and show a central role of host-dependent effects on hybrid fitness and establishment (Büker et al. 2013). The nature of these effects is not known in the *Microbotryum* system. However, they are likely to involve an optimal repertoire of effector genes and associated regulatory machinery essential for successful host infection. An optimal intact combination of effector genes is likely not preserved in hybrid plant pathogen genomes (Le Gac et al. 2007b). Studies from *Microbotryum* and another plant-associated fungus, *Epichloe*, document that specialization to distinct hosts is a primary barrier for interspecific mating between pathogens (Le Gac et al. 2007a; Schirrmann and Leuchtman 2015). Pathogens belonging to distinct species may never encounter each other if they are not compatible on the same host. Hybridization will in this case only be possible if the sexual cycle is detached from host infection or if the life cycle of both parental species includes a saprotrophic stage where hyphal fusions can occur. For most plant pathogens, the biology of growth outside the host is still poorly known.

THE HYBRID GENOME: CONFLICTS AND POTENTIALS

The formation of a fungal hybrid can occur by sexual mating between parental species or fusion of hyphae or vegetative cells. Fusion of vegetative cells can be followed by parasexual mating where mitotic crossover generates recombined hybrid cells (Schardl and Craven 2003). The outcome of hybridization can be an off-spring with the same chromosome number as the parent, known as a homoploid hybrid, or it can be a hybrid with an increased chromosome number known as an allopolyploid or heteroploid hybrid (Mallet 2007).

The genome structure can have a determining effect on fitness and evolution of the hybrid (Yakimowski and Rieseberg 2014). First of all, the hybrid can be affected by genomic incompatibilities between interacting alleles. Secondly, the consequence of chromosome number changes can be reproductive isolation. This is the case when backcrossing with the parental species is present due to a novel chromosome structure in the heteroploid hybrid.

We first consider genomic incompatibilities. Examples of hybrid sterility and inferior fitness observed in plants, animals, and fungi support the hypothesis that incompatibilities between individual genes with major effects, or larger numbers of genes with a cumulative effect, are responsible for deleterious effects in hybrids (Orr 1995). As found by Dettman and colleagues, divergently adaptive yeast strains rapidly evolve genetic incompatibilities, negative epistasis that lower mitotic fitness in yeast hybrid cells (Dettman et al. 2007). When the authors dissected the underlying genetic nature of the evolved incompatibilities in *S. cerevisiae* they found that two genes involved in cell wall synthesis were underexpressed in the hybrid while two other genes involved in sporulation were overexpressed (Dettman et al. 2007; Parreiras et al. 2011). Regulatory defects in hybrids were also shown in crosses of *S. cerevisiae* and *S. bayanus* (Lee et al. 2008) suggesting that negative epistatic interactions in hybrids often are determined by nonoptimal expression of genes.

Several examples of successful hybrids suggest that genomic incompatibilities in some cases, under some circumstances can be overcome (Mallet 2007). In heteroploid hybrids, the genomic contents of both parents may be entirely or partly retained, and genomic incompatibilities in the form of negative epistasis can be reduced, when both parental alleles are expressed. Instead, it is possible that incompatibilities relating to gene dosage may become relevant. Moreover, negative fitness effects caused by genomic incompatibilities can under some circumstances be “overruled” by positive fitness effects in hybrids. As will be discussed below, hybrids can also contain new advantageous allele combinations that contribute to new phenotypic characteristics and that allow them to explore niches not occupied by the parental species (Greig et al. 2002).

As mentioned above, the chromosome content and structure of hybrids have implications for the ability to backcross with parental species. Thereby, the evolutionary potential of heteroploid and homoploid hybrids is not the same and hybrid speciation is considered more likely in heteroploid hybrids (Yakimowski and Rieseberg 2014). This is because the altered genome structure of the hybrid can act as a reproductive barrier and prevent backcrossing to any of the parents. In contrast, backcrossing is a more probabilistic outcome in homoploid hybrids where chromosome number is retained. In other words, introgressive hybridization is a more probable outcome with homoploid hybrids that share the same chromosome composition as their parents (Yakimowski and Rieseberg 2014).

Several studies have in recent years documented that some fungal plant pathogens can tolerate and even benefit from high levels of genomic plasticity, including rearrangements, dynamics of repetitive islands, and accessory chromosomes (see reviews by Croll and McDonald 2012; Raffaele and Kamoun 2012; Seidl and Thomma 2014). To what extent hybrid incompatibilities can be overcome in genomes that already frequently undergo rearrangements should be a central question in future studies of hybrid plant pathogens. In this context, it may be considered that fungi typically produce millions of spores; therefore, many “hybrid experiments” can be tested by natural selection. One or few spores with the right combination of genes can be enough for the establishment of a fit hybrid lineage.

THE ECOLOGICAL NICHE OF HYBRIDS

Not only do genomic conflicts constrain hybrids, but also their level of fitness in a new niche or one co-occupied by parental species. The parents are expected to be considerably better adapted

in their respective environments relative to hybrids of intermediate genotypes and phenotypes. Hybrids may however contain new gene combinations that contribute new phenotypic characteristics, allowing them to exploit other environmental niches not occupied by the parental species (Le Gac et al. 2007a; Schirrmann and Leuchtmann 2015). In yeast, experimental evolution studies revealed that hybrid progenies of *S. cerevisiae* and *S. paradoxus* under certain intermediate environmental conditions can exhibit higher fitness than their parents (Greig et al. 2002). Similar hybrid fitness advantages in plant pathogens have, as mentioned above, been described for hybrids of *Ophiostoma*, *Verticillium*, and *Microbotryum* (Brasier and Kirk 2010; Gibson et al. 2014; Inderbitzin et al. 2011). The ability to colonize a new host niche can be the “key” to establishment without resource competition between parental and hybrid lineages.

Also host expansion through introgressive hybridization has been reported for a number of pathogens including several oomycete species in the genera *Phytophthora* and *Albugo* (Brasier et al. 2004; Goss et al. 2011; McMullan et al. 2015). *Albugo candida* is an obligate biotrophic parasite, comprising a number of races specialized to infect distinct *Brassicaceae* host species. While the genomes of these different races are clearly distinct, they also show signatures of recurrent introgression, including the transfer of host-specific effector determinants (McMullan et al. 2015). Introgressive hybridization is thereby a mechanism by which new host specificities of *A. candida* can rapidly emerge and be transferred between species. The role of introgressive hybridization in adaptation and ecological speciation of fungal plant pathogens is only studied in a few systems (Le Gac et al. 2007a; Schirrmann and Leuchtmann 2015). The acquisition of genome data from an increasing number of species will however provide new possibilities for future studies of introgression between plant pathogens species of wild and cultivated plants.

THE POPULATION GENETICS OF HYBRIDS

Different evolutionary mechanisms, including natural selection, gene flow, recombination and genetic drift shape the population genetic structure of populations of plant pathogens (McDonald and Linde 2002). The population genetic structure of a hybrid plant pathogen may differ considerably from the population genetic structure of the parental species. Consequently, populations of hybrids may evolve by mechanisms other than the parental species.

Hybridization can be associated with a substantial loss of genetic variation as observed in the grass pathogen *Zymoseptoria pseudotritici* (Stukenbrock et al. 2012a). This will be the case when hybrids are formed by a single cross or by few crosses. In the absence of gene flow and backcrossing, the build up of new genetic variation will be determined by the acquisition of new mutations. In this case, the hybrid population will resemble a founder population characterized by low levels of genetic variation and strong effects of genetic drift (Barton and Charlesworth 1984). Hybrids may also emerge as clones that do not undergo sexual recombination, e.g., *Verticillium longisporum* a pathogen of cruciferous plants (Inderbitzin et al. 2011). Comparative genome studies of fungal plant pathogens have shown that adaptive genetic diversity can be generated by other mechanisms than sexual recombination (Seidl and Thomma 2014). Adaptive genetic diversity can evolve in repeat-rich and fast evolving genome compartments (Klostermann et al. 2011; Rouxel et al. 2011). Clonal hybrids may likewise benefit from rapidly evolving repeat-rich regions.

Hybrids that undergo recurrent backcrossing will receive new genetic variation from the parental species. If the extent of backcrossing is high then the genetic differentiation between the hybrid and the parental species will decrease proportionally (Arnold 2004). Below, I briefly consider different population

genetics parameters in the context of hybridization of plant pathogens. Further detailed theoretical analyses of hybrid evolution are presented in reviews by Orr (1995), Baack and Rieseberg (2007), and Barton (2001).

Effective population size and genetic drift. The parameter “effective populations size” termed N_e reflects the amount of genetic diversity that can be passed on from one generation to the next in a population (Wright 1938). Depending on the “format” of the hybridization event, hybrids may emerge with small effective population sizes resembling that of a founder population. In species with small effective population sizes, the effect of genetic drift will be more severe. Genetic drift refers to the random loss of genetic variation and such an effect can potentially aid the negative effect of genomic incompatibilities. In populations with many alleles of a particular gene (high effective population size), the effect of genetic drift is small; however, in populations with fewer alleles (low effective population size), the effect is high (Barton and Charlesworth 1984).

Gene flow. Gene flow mediates the transfer of genetic variation between populations and may be the source of new genetic variation into the hybrid population. Introgressive hybridization is the extreme case in which the hybrid is a “transient” state in the exchange of genetic material between the two parental species (Arnold 2004; Mallet 2005). As discussed above, the extent of backcrossing will be determined by the nature of reproductive barriers between the hybrid and the parental species, which for example may be a difference in genome composition for heteroploid hybrids. Furthermore, the ecological niches occupied by the species will affect rates of gene flow (Giraud et al. 2010). Distinct host compatibilities can act as reproductive barriers between plant pathogens and thereby also isolate hybrids from gene flow (Giraud et al. 2010).

Lastly, gene flow between species can be facilitated by human mediated long distance dispersal of plant pathogens (Brown and Hovmøller 2002). This has for example been the case for the ascomycete fungus *O. ulmi*, the causal agent of Dutch elm disease (Brasier 2000). Intercontinental dispersal of the two species *O. ulmi* and *O. novo-ulmi* led to the emergence of rare hybrids that mediated the transfer of genetic material between *Ophiostoma* species by introgressive hybridization (Brasier 2001).

Natural selection. As already pointed out above, the hybrid genome, in contrast to the genomes of the parental species, was never tested in the same extent by natural selection and comprises gene combinations that did not evolve in parallel (Barton 2001). To which extent hybrids can overcome such genomic incompatibilities depends on multiple factors such as the genetic composition of the hybrid, the nature of negative epistatic interactions, and the interplay of environment, selection and genetics in the hybrid (Barton 2001). As shown experimentally for yeast hybrids (Greig et al. 2002), a particular intermediate “recombinant” environment can provide an optimal niche for a hybrid where it has a selective advantage compared with the parental species.

Recombination and mating systems. Reproduction of fungal plant pathogens ranges from purely clonal reproduction (e.g., *V. dahliae*) to obligate outcrossing (e.g., *Ustilago maydis*) (McDonald and Linde 2002). Furthermore sexually reproducing species can be heterothallic (mating between compatible strains of opposite mating types) or homothallic (when self-fertilization is possible) (Lin and Heitman 2007). In this wide range of reproductive modes and mating systems, hybrids can form in different ways and possess different types of mating systems similar or distinct to the mating systems of their parents.

In hybrids that are able to undergo meiosis, recombination may be important in combining the right parental alleles required for successful establishment and propagation. Recombination can also be instrumental in breaking down genomic incompatibilities between parental traits. Furthermore, as in all sexually reproducing

organisms, recombination in the hybrid can act to speed up adaptive evolution through the fixation of beneficial mutations and the removal of deleterious and nonadaptive variants from the genome (Goddard et al. 2005; Marais and Charlesworth 2003).

Sexual recombination has been a main driver in the emergence and evolution of the hybrid *Z. pseudotritici*, a close relative of the heterothallic ascomycete *Z. tritici* (Stukenbrock et al. 2012b). Consistent with the mating type structure in *Z. tritici*, we found both mating types present among individuals of the hybrid *Z. pseudotritici* (Waalwijk et al. 2002). Furthermore, the genome of this hybrid is shaped by recombination between conspecific strains suggesting that mating behavior resemble mating in *Z. tritici* and other nonhybrid relatives.

In contrast to *Z. pseudotritici*, hybrids of *Epichloe* are asexual while nonhybrids include both asexual and sexual species (Charlton et al. 2014). *Epichloe* hybrids are abundant and are considered to increase the diversity of alkaloids produced to enhance host protection against herbivores (Charlton et al. 2014).

Parasexuality and dikaryosis. Parasexuality refers to the transfer of genetic material between individuals without meiotic recombination and the formation of fruiting bodies. Some asexual pathogens can benefit from parasexual mating by the formation of new genotypes (Hickman et al. 2015). Parasexuality can also be instrumental in the formation of hybrids (Roach and Heitman 2014; Schardl and Craven 2003).

The first step of parasexual reproduction in fungi is the fusion of vegetative cells or hyphae. Normally, vegetative compatibility in both Ascomycetes and Basidiomycetes is governed by genetic programs that prevent fusion with non-conspecific individuals (Glass et al. 2000; Worrall 1997). In cases where vegetative compatibility systems fail to prevent interspecific fusions, a heterokaryon will be formed. Heterokaryons of incompatible strains are often unstable and revert into homokaryons or they show reduced growth and fitness (Kausserud et al. 2012). But in some cases the heterokaryon can convert into heteroploid or haploid cells or hyphae by parasexual nuclear fusion and mitotic cross-over of chromosomes. Recent studies of the human asexual pathogen *Candida albicans* show that parasexual mating can generate a diverse population of progenitors differing in ploidy and chromosome numbers (Hickman et al. 2015). In the same way, parasexual mating between non-conspecific partners can give rise to a diverse population of heteroploid hybrid individuals from which natural selection can “pick out” fit genotypes.

RECENT EXAMPLES OF PLANT-PATHOGENIC FUNGAL HYBRIDS

***Z. pseudotritici*: A one-time fit combination.** *Z. pseudotritici*, a close relative of the prominent wheat pathogen *Z. tritici*, was isolated from two distinct grass hosts *Elymus repens* and *Dactylis glomerata* in the north of Iran (Stukenbrock et al. 2012b). Population genomic sequence data of the haploid *Z. pseudotritici* revealed a peculiar mosaic genome structure with long segments completely lacking variation (Stukenbrock et al. 2012a). The nonvariable segments were found interspersed by regions of unusual high sequence variation. The variable segments consistently exhibited only two haplotypes differing by 30 single nucleotide polymorphisms per 1 kb on average, corresponding to 3% nucleotide variation between haplotypes.

A genomic region with comparatively low levels of genetic variation often indicates a selective sweep where a beneficial allele in a population of individuals through time has been fixed by natural selection. In the case of *Z. pseudotritici*, however, the nonvariable segments were interspersed between variable regions always comprising two haplotypes. This mosaic genome structure of *Z. pseudotritici* did not support a selective sweep model. We instead propose that hybrid speciation explains this peculiar genomic

mosaic in *Z. pseudotritici* and postulated that the species was formed by a single hybridization event between two individual from two closely related species or lineages about 3% divergent from each other. The nonvariable segments in the genome are chromosomal regions that were transmitted from one parent to all hybrid progeny. The variable segments comprising two haplotypes are chromosomal regions that were transmitted to the hybrid swarm from both parents. Using the correlation of the length distribution of variable segments with recombination rate and generation time of *Z. pseudotritici* it was possible to estimate the number of recombination events that have occurred since the hybridization event forming *Z. pseudotritici*. From the generation time of *Z. pseudotritici* we found that the mosaic genome of *Z. pseudotritici* results from approximately 400 recombination events between individuals of the hybrid swarm. The lack of additional polymorphism in the nonvariable segments indicates that backcrossing to the parental species did not occur after the initial hybridization event.

Z. pseudotritici was found to occur frequently on grasses collected at five plots along a 500 km transect in Iran. Individuals from the parental species were not collected from grasses at the same plots at subsequent collecting trips in the same area (M. Javan-Nikkhah, M. Zala, B. A. McDonald, and E. H. Stukenbrock, unpublished data), suggesting that the hybrid has invaded a completely new niche (in time or space) or outcompeted both parental species.

Z. pseudotritici is an excellent example of recent hybrid speciation of a fungal plant pathogen. A number of questions remain to be addressed in this hybrid system. Which hosts did the parental species colonize? Why did only one successful hybrid clone emerge from the parental species? How has successful adaptation over multiple generations been possible following extreme loss of genetic variation? Further dissection of the genome evolution in *Z. pseudotritici* as well as more extensive sampling at the center of origin of *Z. pseudotritici* including *Septoria* pathogens from non-*Poaceae* hosts will provide further insight into the origin and evolution of this hybrid.

Microbotryum: Incomplete reproductive barriers and introgression. The species complex of the anther smut fungus *M. violaceum* (Basidiomycota) provides an excellent model system to address the importance of introgressive hybridization among closely related plant pathogens (Giraud et al. 2008; Gladieux et al. 2011; Le Gac et al. 2007a,b). Gladieux and colleagues addressed the extent of hybridization between the two species *M. lychnidis-dioicae* and *M. silene-dioicae* using population genetic analyses and coalescent models (Gladieux et al. 2011). *M. lychnidis-dioicae* and *M. silene-dioicae* are ecologically isolated from reproduction as they each are specialized to distinct *Silene* hosts. In a large collection of *Microbotryum* isolates, the authors identified only a small number of hybrids (15 out of 1,028 isolates) based on microsatellite data and a Bayesian clustering algorithm. Few isolates were assigned as “cross-species” isolates, i.e., isolate assigned to either *M. lychnidis-dioicae* and *M. silene-dioicae* but isolated from the original host of the other *Microbotryum* species (Gladieux et al. 2011). These “cross-species” isolates show that host-defined prezygotic reproductive barriers can be permissive allowing isolates from distinct hosts to cross with each other. By applying an “isolation-with-migration” coalescence model to the data set, the authors were able to date the divergence of *M. lychnidis-dioicae* and *M. silene-dioicae* and to estimate when gene flow between species occurred (Gladieux et al. 2011). Consistent with the low frequency of hybrids, analyses of gene flow revealed recent divergence times and low rates of gene flow between the two *Microbotryum* species. Although the two pathogen-species co-exist in the same environment, reproductive barriers determined by the host may be sufficient to prevent the long-term establishment of hybrid individuals. Nevertheless, recurrent introgressive hybridization between *Microbotryum* species

may allow the transfer of new adaptive traits between species and eventually promote the evolution of new host specificities. Experimental back-crossings of *M. lychnidis-dioicae* and *M. silene-dioicae* hybrids have demonstrated a mating type effect on interspecific conjugation formation (Büker et al. 2013). This could suggest that not only host factors determine reproductive barriers but in addition, as also observed in *N. crassa* (Turner et al. 2011), mating type related factors.

Comparative genome analyses may in the future allow the identification of introgressed traits in *Microbotryum* species. Which and how many genes determine host specificity in the *Microbotryum* system still needs to be determined. An intriguing question will be whether these genes have been introgressed between species as predicted.

Epichloe: Hybrids and the evolution of host specificities. The ascomycete endophyte *Epichloe* is the main fungal model for studies of hybridization and the impact of hybridization on phenotypes in planta. Species of *Epichloe* colonize the apoplast of pooid grasses. The endophyte has little effect on the vegetative tissue of its host; however, during sexual reproduction *Epichloe* can negatively affect the development of flowering shoots by formation of fungal sexual structures in these plant parts, causing the so-called “choke” disease (Schardl et al. 2004). Sexual reproduction of the pathogenic *Epichloe* species is thereby tightly associated with reproductive success of the host. *Epichloe* also includes mutualistic endophytic species that confer a fitness advantage to their host, for example by the production of anti-herbivore alkaloids (Schardl 2001). Interestingly, several mutualistic *Epichloe* species have been identified as hybrids. These hybrids do not originate from in-terspecific sexual mating, as mating populations in the field can be clearly distinguished according to sexual compatibility and host specificities (Schirrmann and Leuchtman 2015). Rather, hybrids are asexual lineages that originate from vegetative fusions between hyphae of different *Epichloe* species (Shoji et al. 2015; Tsai et al. 1994).

Asexual *Epichloe* species were previously classified in the genus *Neotyphodium*, but they were recently attributed to the genus *Epichloe* (Leuchtman et al. 2014). Interestingly, phylogenetic analyses show that asexual species frequently are heteroploid hybrids emerged from interspecific hyphal fusions of other *Epichloe* species (Moon et al. 2004; Shoji et al. 2015).

In a recent paper, Shoji and coworkers assessed the fate of nuclei and organelles during vegetative hyphal fusions in different *Epichloe* species (Shoji et al. 2015). Cytological analyses of interspecific hyphal fusions showed that distinct types of nuclei never coexisted. Viable hybrids are therefore formed by nuclear fusion and the formation of allodiploid hyphae. The authors also addressed the frequency of interspecific hyphal fusions and could with quantitative analyses show that interspecific hyphal fusions are rare. Nevertheless, in nature interspecific hybrids of *Epichloe* are found frequently. The authors propose that established hybrids could have a fitness advantage through host specificities acquired by the hybridization event allowing them to rapidly increase in frequency. It is also possible that *in-planta* conditions promote hyphal fusions to an extent not reproduced experimentally.

Interestingly, nonhybrid and hybrid *Epichloe* endophytes differently affect growth and reproduction of host species. The evidence suggests that the performance of hybrid compared with nonhybrid endophytes depends on host genotype and environmental conditions. Hybrids were shown to enhance the competitive abilities of the grass *Festuca arizonica* by increased biomass production (Saari and Faeth 2012). But in another grass species *Hordelymus europaeus* nonhybrid *Epichloe* endophytes were found to have more positive influence, compared with hybrid endophytes, on host fitness measured as seed production of different accessions of *H. europaeus* (Oberhofer et al. 2014). As for *Z. pseudotritici* and *Microbotryum*, a deeper understanding of the underlying

host–fungus interaction is necessary to unravel differences in the biology of hybrid and nonhybrid *Epichloe* species.

***V. longisporum*: Multiple parents and multiple hybridizations.** Formation of the diploid hybrid *V. longisporum* occurred by vegetative fusion of other haploid *Verticillium* species. Phylogenetic analyses have shown that the hybrid originated from multiple hybridization events involving the unknown Species A1 and Species D1 and the D2 and D3 lineages of *V. dahliae* (Inderbitzin et al. 2011; Inderbitzin and Subbarao 2014).

Both genome structure and host range of *V. longisporum* differ from those of *V. dahliae*. While *V. dahliae* is known to have a broad host range, it is, with some exceptions, not infectious on *Brassicaceae* hosts. *V. longisporum* on the other hand is mainly a pathogen of *Brassicaceae* plants (Zeise and Tiedemann 2002), indicating that new host determinants were acquired from the unknown parental species (A1 or D1) or emerged by the combination of multiple host determinants from several parents. The capacity to infect new host species may however not only have resulted from a new combination of genes but also from the doubling of chromosomes in the diploid *V. longisporum*. Populations of other heteroploid hybrids have similarly been shown to comprise an increased phenotypic variability compared with the parental species allowing the exploitation of a broader range of host species (Brasier et al. 2004; Newcombe et al. 2001). To address the genomic source of new virulence traits in heteroploid hybrids as *V. longisporum*, the underlying determinants of host infections are still to be discovered.

STUDIES OF FUNGAL HYBRID SPECIES: FUTURE DIRECTIONS

Comparative genomics and population genomics provide new powerful approaches to recognize ongoing and past hybridization in plant pathogens (Menardo et al. in press; Stukenbrock et al. 2012a). Likewise, experimental evolution studies have shed light on the genetics of reproductive barriers in model ascomycetes (Dettman et al. 2007; Turner et al. 2011). Although hybridization may be an important (yet underestimated) force in the evolution of plant pathogens, we still know little about the biology and genetics of hybrid plant-pathogenic fungi. A key question in evolutionary genomics of hybrids is how negative epistatic interactions of distinct parental alleles are overcome. Future studies should combine experimental and genomics approaches to explore the genetics of hybrid incompatibilities and to assess phenotypic and evolutionary potentials of hybrid plant pathogens.

Population genomics. I consider at least four important applications of population genomics in hybrid studies. The first is the use of population genomic data to investigate the genome wide distribution of genetic variation in hybrids. As discussed above, this may deviate significantly from the parents, but nevertheless influence the evolutionary potential of the hybrid, including the potential for dispersal, introgression and adaptation. Secondly, population genomic data can be explored to identify signatures of selection in the hybrid genome reflecting novel adaptations and the selection against incompatible allele combinations (Stukenbrock 2013). Signatures of selection can be identified as outlier loci of increased divergence (Ellison et al. 2011), regions depleted of variation (selective sweeps) (e.g., Shapiro et al. 2012) or genes with an increased proportion of nonsynonymous variation (balancing selection) (Stukenbrock et al. 2011). Thirdly, introgressive hybridization will leave strong signatures of interspecific exchanges of genetic material (Arnold 2004). Genomic data from hybrids and their parents allow the identification of these transferred and putative adaptive traits (Arnold 2004). Introgressed regions are recognized in genome scans as segments with altered levels of divergence (Menardo et al. in press). For example, divergence will be reduced in regions affected by introgressive

hybridization. Lastly, genomic incompatibilities in hybrids may be recognized by in depth analyses of linkage disequilibrium or QTL mapping. As described above experimental evolution, QTL analyses and genomics have shed light on genetic incompatibilities in *S. cerevisiae* and *N. crassa* (Dettman et al. 2007; Turner et al. 2011).

Experimental studies in planta and in vitro. Few experimental studies have used in planta experiments to compare fitness of fungal hybrids relative to nonhybrid species (Büker et al. 2013; Saari and Faeth 2012; Oberhofer et al. 2014). It has been documented that hybridization can change the host range of plant pathogens but it is not known how (Inderbitzin et al. 2011). Is it the new combination of effectors that allow hybrids to explore novel host niches? Or do changes in ploidy and gene dosage alter virulence in the hybrid? Studies that aim to identify new hybrid host determinants are essential to understand the success of hybrid plant pathogens. Such traits may be identified from comparative genome or transcriptome studies and their functional relevance assessed by reverse genetics and experimental approaches. An intriguing question is whether such traits are inherited from one parental species and only confer a distinct fitness advantage in the genomic background of the hybrid, or if these traits are recombined and represent completely new variants.

Finally, a better understanding of the underlying genetics of reproductive barriers and self-versus-nonsel self recognition between species of fungal plant pathogens could improve predictions of hybridization events in agroecosystems. As shown in *N. crassa*, genetic determinants of reproductive barriers can be identified by experimental hybridization assays and QTL mappings (Turner et al. 2011). Cytological analyses of between-species, vegetative hyphal fusions can furthermore be applied to study nuclear behavior during interspecific fusions and the formation of heteroploid hybrids (Shoji et al. 2015).

A number of central questions relating to hybrid evolution can be addressed by experimental evolution approaches. In contrast to retrospective sequence-based analyses, experimental evolution of microbial species provides the opportunity to directly study rates of mutational changes and to identify the nature and order of these changes. It also allows fitness assays of all generations from progenitor to last evolved population and thereby a comparison of fitness effects of ancestral and derived alleles. Experimental evolution could be applied for example in studies of hybrid pathogens aiming at addressing the fate of genetic variation and genomic incompatibilities in hybrids following multiple generations of asexual or sexual propagation.

ACKNOWLEDGMENTS

I thank F. B. Christiansen and J. Y. Dutheil for fruitful discussions and helpful comments and two anonymous reviewers for comments to a previous version of the manuscript. Research conducted in the group of E. H. Stukenbrock is funded by the Max Planck Society, Germany and a personal grant from the State of Schleswig-Holstein, Germany to E. H. Stukenbrock.

LITERATURE CITED

- Anderson, J. B., Korhonen, K., and Ullrich, R. C. 1980. Relationships between European and North American biological species of *Armillaria mellea*. *Exp. Mycol.* 4:78-86.
- Arnold, M. L. 2004. Transfer and origin of adaptations through natural hybridization: Were Anderson and Stebbins right? *Plant Cell* 16:562-570.
- Baack, E. J., and Rieseberg, L. H. 2007. A genomic view of introgression and hybrid speciation. *Curr. Opin. Genet. Dev.* 17:513-518.
- Barton, N. H. 2001. The role of hybridization in evolution. *Mol. Ecol.* 10:551-568.
- Barton, N. H., and Charlesworth, B. 1984. Genetic revolutions, founder effects, and speciation. *Annu. Rev. Ecol. System.* 15:133-164.

- Brasier, C. 2000. Plant pathology: The rise of the hybrid fungi. *Nature* 405: 134-135.
- Brasier, C. M. 2001. Rapid evolution of introduced plant pathogens via interspecific hybridization is leading to rapid evolution of Dutch elm disease and other fungal plant pathogens. *Bioscience* 51:123-133.
- Brasier, C. M., and Kirk, S. A. 2010. Rapid emergence of hybrids between the two subspecies of *Ophiostoma novo-ulmi* with a high level of pathogenic fitness. *Plant Pathol.* 59:186-199.
- Brasier, C. M., Kirk, S. A., Delcan, J., Cooke, D. E., Jung, T., and Man In't Veld, W. A. 2004. *Phytophthora alni* sp. nov. and its variants: Designation of emerging heteroploid hybrid pathogens spreading on *Alnus* trees. *Mycol. Res.* 108:1172-1184.
- Brasier, C. M., Kirk, S. A., Pipe, N., and Buck, K. W. 1998. Rare hybrids in natural populations of the Dutch elm disease pathogen *Ophiostoma ulmi* and *O. novo-ulmi*. *Mycol. Res.* 102:45-57.
- Brown, J. K. M., and Hovmöller, M. S. 2002. Aerial dispersal of pathogens on the global and continental scales and its impact on plant disease. *Science* 297:537-41.
- Büker, B., Petit, E., Begerow, D., and Hood, M. E. 2013. Experimental hybridization and backcrossing reveal forces of reproductive isolation in *Microbotryum*. *BMC Evol. Biol.* 13:224.
- Charlton, N. D., Craven, K. D., Afkhami, M. E., Hall, B. A., Ghimire, S. R., and Young, C. A. 2014. Interspecific hybridization and bioactive alkaloid variation increases diversity in endophytic *Epichloë* species of *Bromus laevipes*. *FEMS Microbiol. Ecol.* 90:276-289.
- Croll, D., and McDonald, B. A. 2012. The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS Pathog.* 8:e1002608.
- Dettman, J. R., Anderson, J. B., and Kohn, L. M. 2008. Divergent adaptation promotes reproductive isolation among experimental populations of the filamentous fungus *Neurospora*. *BMC Evol. Biol.* 8:35.
- Dettman, J. R., Jacobson, D. J., Turner, E., Pringle, A., and Taylor, J. W. 2003. Reproductive isolation and phylogenetic divergence in *Neurospora*: Comparing methods of species recognition in a model Eukaryote. *Evolution* 57: 2721-2741.
- Dettman, J. R., Sirjusingh, C., Kohn, L. M., and Anderson, J. B. 2007. Incipient speciation by divergent adaptation and antagonistic epistasis in yeast. *Nature* 447:585-588.
- Ellison, C. E., Hall, C., Kowbel, D., Welch, J., Brem, R. B., Glass, N. L., and Taylor, J. W. 2011. Population genomics and local adaptation in wild isolates of a model microbial eukaryote. *Proc. Natl. Acad. Sci.* 108: 2831-2836.
- Gibson, A. K., Refrégier, G., Hood, M. E., and Giraud, T. 2014. Performance of a hybrid fungal pathogen on pure-species and hybrid host plants. *Int. J. Plant Sci.* 175:724-730.
- Giraud, T., Gladioux, P., and Gavrillets, S. 2010. Linking the emergence of fungal plant diseases with ecological speciation. *Trends Ecol. Evol.* 25: 387-395.
- Giraud, T., Yockteng, R., López-Villavicencio, M., Refrégier, G., and Hood, M. E. 2008. Mating system of the anther smut fungus *Microbotryum violaceum*: Selfing under heterothallism. *Eukaryot. Cell* 7:765-775.
- Gladioux, P., Vercken, E., Fontaine, M. C., Hood, M. E., Jonot, O., Couloux, A., and Giraud, T. 2011. Maintenance of fungal pathogen species that are specialized to different hosts: Allopatric divergence and introgression through secondary contact. *Mol. Biol. Evol.* 28:459-471.
- Glass, N. L., Jacobson, D. J., and Shiu, P. K. 2000. The genetics of hyphal fusion and vegetative incompatibility in filamentous ascomycete fungi. *Annu. Rev. Genet.* 34:165-186.
- Goddard, M. R., Godfray, H., and Burt, A. 2005. Sex increases the efficacy of natural selection in experimental yeast populations. *Nature* 434:636-640.
- Goss, E. M., Cardenas, M. E., Myers, K., Forbes, G. A., Fry, W. E., Restrepo, S., and Grünwald, N. J. 2011. The plant pathogen *Phytophthora andina* emerged via hybridization of an unknown *Phytophthora* species and the Irish potato famine pathogen, *P. Infestans*. *PLoS One* 6:e24543.
- Greig, D., Louis, E. J., Borts, R. H., and Travisano, M. 2002. Hybrid speciation in experimental populations of yeast. *Science* 298:1773-1775.
- Hickman, M. A., Paulson, C., Dudley, A. M., and Berman, J. 2015. Parasexual ploidy reduction drives population heterogeneity through random and transient aneuploidy. *Genetics* 200:781-794.
- Inderbitzin, P., Davis, R. M., Bostock, R. M., and Subbarao, K. V. 2011. The ascomycete *Verticillium longisporum* is a hybrid and a plant pathogen with an expanded host range. *PLoS One* 6:e18260.
- Inderbitzin, P., and Subbarao, K. V. 2014. *Verticillium* systematics and evolution: How confusion impedes *Verticillium* wilt management and how to resolve it. *Phytopathology* 104:564-574.
- Kausserud, H., Knudsen, H., Høggberg, N., and Skrede, I. 2012. Evolutionary origin, worldwide dispersal, and population genetics of the dry rot fungus *Serpula lacrymans*. *Fungal Biol. Rev.* 26:84-93.
- Klosterman, S. J., Subbarao, K. V., Kang, S., Veronese, P., Gold, S. E., Thomma, B. P. H. J., Chen, Z., Henrissat, B., Lee, Y.-H., Park, J., Garcia-Pedrajas, M. D., Barbara, D. J., Anchieta, A., de Jonge, R., Santhanam, P., Maruthachalam, K., Atallah, Z., Amyotte, S. G., Paz, Z., Inderbitzin, P., Hayes, R. J., Heiman, D. I., Young, S., Zeng, Q., Engels, R., Galagan, J., Cuomo, C. A., Dobinson, K. F., and Ma, L.-J. 2011. Comparative genomics yields insights into niche adaptation of plant vascular wilt pathogens. *PLoS Pathog.* 7:e1002137.
- Kondrashov, A. S., Sunyaev, S., and Kondrashov, F. A. 2002. Dobzhansky-Muller incompatibilities in protein evolution. *Proc. Natl. Acad. Sci.* 99: 14878-14883.
- Kohn, L. M. 2005. Mechanisms of fungal speciation. *Annu. Rev. Phytopathol.* 43:279-308.
- Kuehne, H. A., Murphy, H. A., Francis, C. A., and Sniegowski, P. D. 2007. Allopatric divergence, secondary contact, and genetic isolation in wild yeast populations. *Curr. Biol.* 17:407-411.
- Lamichhane, S., Berglund, J., Almén, M. S., Maqbool, K., Grabherr, M., Martinez-Barrio, A., Promerová, M., Rubin, C.-J., Wang, C., Zamani, N., Grant, B. R., Grant, P. R., Webster, M. T., and Andersson, L. 2015. Evolution of Darwin's finches and their beaks revealed by genome sequencing. *Nature* 518:371-375.
- Le Gac, M., Hood, M. E., Fournier, E., and Giraud, T. 2007a. Phylogenetic evidence of host-specific cryptic species in the anther smut fungus. *Evolution* 61:15-26.
- Le Gac, M., Hood, M. E., and Giraud, T. 2007b. Evolution of reproductive isolation within a parasitic fungal species complex. *Evolution* 61:1781-1787.
- Lee, H. Y., Chou, J. Y., Cheong, L., Chang, N. H., Yang, S. Y., and Leu, J. Y. 2008. Incompatibility of nuclear and mitochondrial genomes causes hybrid sterility between two yeast species. *Cell* 135:1065-1073.
- Leuchtmann, A., Bacon, C. W., Schardl, C. L., White, J. F., and Tadych, M. 2014. Nomenclatural realignment of *Neotyphodium* species with genus *Epichloë*. *Mycologia* 106:202-215.
- Lin, X., and Heitman, J. 2007. Mechanisms of homothallism in fungi and transitions between heterothallism and homothallism. Pages 35-57 in: *Sex in Fungi*. J. Heitman, J. W. Kronstad, J. W. Taylor, and L. A. Casselton, eds. ASM Press, Washington, D.C.
- Mallet, J. 2005. Hybridization as an invasion of the genome. *Trends Ecol. Evol.* 20:229-237.
- Mallet, J. 2007. Hybrid speciation. *Nature* 446:279-283.
- Marais, G., and Charlesworth, B. 2003. Genome evolution: recombination speeds up adaptive evolution. *Curr. Biol.* 13:R68-R70.
- McDonald, B. A., and Linde, C. 2002. Pathogen population genetics, evolutionary potential, and durable resistance. *Annu. Rev. Phytopathol.* 40:349-379.
- McMullan, M., Gardiner, A., Bailey, K., Kemen, E., Ward, B. J., Cevik, V., Robert-Saileniantz, A., Schultz-Larsen, T., Balmuth, A., Holub, E., van Oosterhout, C., and Jones, J. D. G. 2015. Evidence for suppression of immunity as a driver for genomic introgressions and host range expansion in races of *Albugo candida*, a generalist parasite. *eLife* 4:e04550.
- Menardo F., Praz, C., Wyder, S., Ben-David, R., Bourras, S. A., Matsumae, H., McNally, K. E., Parlange, F., Riba, A., Roffler, S., Schaefer, L. K., Shimizu, K. K., Valenti, L., Zbinden, H., Wicker, T., and Keller, B. Hybridization of powdery mildew strains gives rise to pathogens on novel agricultural crop species. *Nat. Genet.* (In press.)
- Moon, C. D., Craven, K. D., Leuchtmann, A., Clement, S. L., and Schardl, C. S. 2004. Prevalence of interspecific hybrids amongst asexual fungal endophytes of grasses. *Mol. Ecol.* 13:1455-1467.
- Newcombe, G., Stirling, B., and Bradshaw, H. D. 2001. Abundant pathogenic variation in the new hybrid rust *Melampsora × columbiana* on hybrid poplar. *Phytopathology* 91:981-985.
- Oberhofer, M., Güsewell, S., and Leuchtmann, A. 2014. Effects of natural hybrid and non-hybrid *Epichloë* endophytes on the response of *Hordelymus europaeus* to drought stress. *New Phytol.* 201:242-253.
- Orr, H. A. 1995. The population genetics of speciation: The evolution of hybrid incompatibilities. *Genetics* 139:1805-1813.
- Pardo-Diaz, C., Salazar, C., Baxter, S. W., Merot, C., Figueiredo-Ready, W., Joron, M., Owen McMillan, W., and Jiggins, C. D. 2012. Adaptive introgression across species boundaries in *Heliconius* butterflies. *PLoS Genet.* 8:e1002752.
- Parreiras, L. S., Kohn, L. M., and Anderson, J. B. 2011. Cellular effects and epistasis among three determinants of adaptation in experimental populations of *Saccharomyces cerevisiae*. *Eukaryot. Cell* 10:1348-1356.
- Raffaele, S., and Kamoun, S. 2012. Genome evolution in filamentous plant pathogens: Why bigger can be better. *Nat. Rev. Microbiol.* 10:417-430.
- Roach, K. C., and Heitman, J. 2014. Unisexual reproduction reverses Muller's ratchet. *Genetics* 198:1059-1069.
- Rouxel, T., Grandaubert, J., Hane, J. K., Hoede, C., van de Wouw, A. P., Couloux, A. V., Dominguez, V., Anthouard, V., Bally, P., Bourras, S., Cozijnsen, A. J., Ciuffetti, L. M., Degraeve, A., Dilmaghani, A., Duret, L.,

- Fudal, I., Goodwin, S. B., Gout, L., Glaser, N., Linglin, J., Kema, G. H. J., Lapalu, N., Lawrence, C. B., May, K., Meyer, M., Ollivier, B., Poulain, J., Schoch, C. L., Simon, A., Spatafora, J. W., Stachowiak, A., Turgeon, B. G., Tyler, B. M., Vincent, D., Weissenbach, J., Amselem, J., Quesneville, H., Oliver, R. P., Wincker, P., Balesdent, M.-H., and Howlett, B. J. 2011. Effector diversification within compartments of the *Leptosphaeria maculans* genome affected by repeat-induced point mutations. *Nat. Commun.* 2:202.
- Saari, S., and Faeth, S. H. 2012. Hybridization of Neotyphodium endophytes enhances competitive ability of the host grass. *New Phytol.* 195:231-236.
- Schardl, C. L. 2001. *Epichloë festucae* and related mutualistic symbionts of grasses. *Fungal Genet. Biol.* 33:69-82.
- Schardl, C. L., and Craven, K. D. 2003. Interspecific hybridization in plant-associated fungi and oomycetes: A review. *Mol. Ecol.* 12:2861-2873.
- Schardl, C. L., Leuchtman, A., and Spiering, M. J. 2004. Symbioses of grasses with seedborne fungal endophytes. *Annu. Rev. Plant Biol.* 55:315-340.
- Schirrmann, M. K., and Leuchtman, A. 2015. The role of host-specificity in the reproductive isolation of *Epichloë* endophytes revealed by reciprocal infections. *Fungal Ecol.* 15:29-38.
- Seidl, M. F., and Thomma, B. P. H. J. 2014. Sex or no sex: Evolutionary adaptation occurs regardless. *BioEssays* 36:335-345.
- Shapiro, B. J., Friedman, J., Cordero, O. X., Preheim, S. P., Timberlake, S. C., Szabó, G., Polz, M. F., and Alm, E. J. 2012. Population genomics of early events in the ecological differentiation of bacteria. *Science* 336:48-51.
- Shoji, J.-Y., Charlton, N. D., Yi, M., Young, C. A., and Craven, K. D. 2015. Vegetative hyphal fusion and subsequent nuclear behavior in *Epichloë* grass endophytes. *PLoS One* 10:e0121875.
- Stankowski, S., and Streisfeld, M. A. 2015. Introgressive hybridization facilitates adaptive divergence in a recent radiation of monkeyflowers. *Proc. R. Soc. B* 282:1814.
- Stenlid, J., and Karlsson, J.-O. 1991. Partial intersterility in *Heterobasidion annosum*. *Mycol. Res.* 95:1153-1159.
- Stukenbrock, E. H. 2013. Evolution, selection and isolation: A genomic view of speciation in fungal plant pathogens. *New Phytol.* 199:895-907.
- Stukenbrock, E. H., Bataillon, T., Dutheil, J. Y., Hansen, T. T., Li, R., Zala, M., and Schierup, M. H. 2011. The making of a new pathogen: Insights from comparative population genomics of the domesticated wheat pathogen *Mycosphaerella graminicola* and its wild sister species. *Genome Res.* 21:2157-2166.
- Stukenbrock, E. H., Christiansen, F. B., Hansen, T. T., Dutheil, J. Y., and Schierup, M. H. 2012a. Fusion of two divergent fungal individuals led to the recent emergence of a unique widespread pathogen species. *Proc. Natl. Acad. Sci.* 109:10954-10959.
- Stukenbrock, E. H., Quaedvlieg, W., Javan-Nikhah, M., Zala, M., Crous, P. W., and McDonald, B. A. 2012b. *Zymoseptoria ardabiliae* and *Z. pseudotritici*, two progenitor species of the Septoria tritici leaf blotch fungus *Z. tritici* (synonym: *Mycosphaerella graminicola*). *Mycologia* 104:1397-1407.
- Tsai, H. F., Liu, J. S., Staben, C., Christensen, M. J., Latch, G. C., Siegel, M. R., and Schardl, C. L. 1994. Evolutionary diversification of fungal endophytes of tall fescue grass by hybridization with *Epichloë* species. *Proc. Natl. Acad. Sci.* 91:2542-2546.
- Turner, E., Jacobson, D. J., and Taylor, J. W. 2011. Genetic architecture of a reinforced, postmating, reproductive isolation barrier between *Neurospora* species indicates evolution via natural selection. *PLoS Genet.* 7:e1002204.
- Waalwijk, C., Mendes, O., Verstappen, E. C., de Waard, M. A., and Kema, G. H. 2002. Isolation and characterization of the mating-type idiomorphs from the wheat Septoria leaf blotch fungus *Mycosphaerella graminicola*. *Fungal Genet. Biol.* 35:277-286.
- Worrall, J. J. 1997. Somatic incompatibility in basidiomycetes. *Mycologia* 89:24-36.
- Wright, S. 1938. Size of population and breeding structure in relation to evolution. *Science* 87:430-431.
- Yakimowski, S., and Rieseberg, L. H. 2014. The role of homoploid hybridization in evolution: A century of studies synthesizing genetics and ecology. *Am. J. Bot.* 101:1247-1258.
- Zeise, K., and Tiedemann, A. V. 2002. Host specialization among vegetative compatibility groups of *Verticillium dahliae* in relation to *Verticillium longisporum*. *J. Phytopathol.* 150:112-119.