Quantification of *Botrytis cinerea* in Grapevine Bunch Trash by Real-Time PCR

Melissa Si Ammour,¹ Giorgia Fedele,¹ Caterina Morcia,² Valeria Terzi,² and Vittorio Rossi^{1,†}

¹ Department of Sustainable Crop Production (DI.PRO.VE.S.), Università Cattolica del Sacro Cuore, 29122 Piacenza, Italy

² Research Centre for Genomics and Bioinformatics (CREA-GB), Council for Agricultural Research and Economics, 29017 Fiorenzuola d'Arda, Italy

Accepted for publication 18 February 2019.

ABSTRACT

Quantification of colonization of grape bunch trash by *Botrytis cinerea* is crucial for Botrytis bunch rot (BBR) control. A previously developed quantitative polymerase chain reaction (qPCR) method was adapted to quantify *B. cinerea* DNA in grape bunch trash, and a colonization coefficient (CC) was calculated as the ratio between the DNA concentrations of *B. cinerea* and of *Vitis vinifera*. CC values increased linearly with the number of conidia of *B. cinerea* or the quantity of mycelium of *B. cinerea* added to the bunch trash increased. CC values also increased linearly in bunch trash samples containing increasing percentages of *B. cinerea*-colonized bunch trash; in the latter samples, CC values were correlated with subsequent assessments of *B. cinerea*

colonization of trash (as determined by plating on agar) and sporulation on the trash (as determined by spore counts after incubation in humid chambers). The qPCR assay was also validated using trash collected from bunches treated or not treated with fungicides in three vineyards in two seasons. CC values reflected the reduction in sporulation and in latent infections of mature berries caused by fungicide application. The qPCR assay enables rapid, specific, sensitive, and reliable quantification of the degree of colonization of bunch trash by *B. cinerea*, which makes it a useful tool for studies of the epidemiology and management of BBR.

Keywords: colonization rate, gray mold, sporulation potential

Botrytis bunch rot (BBR) is an economically important disease of grapevines (*Vitis vinifera* L.) and is caused by the fungus *Botrytis cinerea* Pers.: Fr (Elmer and Michailides 2007). Control of BBR is challenging for the following reasons: (i) *B. cinerea* can function as a saprophyte, necrotroph, or parasite; (ii) the fungus can overwinter and sporulate on multiple inoculum sources (including bunch trash, leaf trash, and rotted berries); (iii) grapevines are susceptible at multiple growth stages; (iv) multiple infection pathways exist; and (v) infections can occur under a wide range of environmental conditions, which differ among infection pathways (Ciliberti et al. 2015a, 2015b, 2016; Elad et al. 2007; Elmer and Michailides 2007; Hill et al. 2014; Nair et al. 1995).

After flowering, the pathogen is able to saprophytically colonize the “bunch trash” (i.e., the dead stamens, aborted flowers, aborted berries, calyptras, tendrils, and leaf pieces retained within developing bunches) (Seyb et al. 2000). Under favorable conditions, the fungus produces abundant conidia on the colonized bunch trash, and these conidia are a source of inoculum for berry infection, mainly after veraison. Elmer and Michailides (2007) referred to these phenomena as infection pathways III (conidial infection and extensive colonization of floral debris in grape bunches), IV (conidial accumulation within the developing bunch), and V (conidial infection of ripening fruit), respectively.

Bunch trash colonized early by *B. cinerea* is a major source of berry infection (Elmer and Michailides 2007) and has been related to the severity of BBR at harvest (Holz et al. 2003; Keller et al. 2003; Nair et al. 1995; Viret et al. 2004). Thus, reducing the quantity of bunch trash, as well as reducing colonization of bunch trash by *B. cinerea* at flowering and postflowering, should contribute to

control of BBR (Calvo-Garrido et al. 2014; González-Domínguez et al. 2015). Reduction of available bunch trash has been explored by removal of floral debris from clusters either at early or late fruit set using compressed air or leaf blowers (Wolf et al. 1997) and fungicide sprays (González-Domínguez et al. 2015) and the application of biocontrol agents and other natural products (Calvo-Garrido et al. 2014; Pertot et al. 2017) has been investigated to reduce colonization of bunch trash by *B. cinerea*.

Evaluation of the effectiveness of these interventions requires methods to quantify the colonization of bunch trash by *B. cinerea* and the subsequent production of spores. Traditionally, *B. cinerea* colonization of bunch trash has been quantified by plating on selective media (Abdelwahab and Younis 2012; Edwards and Seddon 2001) or by microscopic assessment (Calvo-Garrido et al. 2014). Sporulation on bunch trash was measured using a sporulation index on a 0 to 5 scale (Calvo-Garrido et al. 2014) or by microscope counts of conidia (Jaspers et al. 2013; Keller et al. 2003; Mundy et al. 2012; Nair et al. 1995). As alternatives to traditional methods, molecular tools may offer rapid, specific, and accurate estimation of the quantity of *B. cinerea* in bunch trash (Abdelwahab and Younis 2012; Diguta et al. 2010).

A direct polymerase chain reaction (PCR) assay has been described for the detection of *B. cinerea* in pea-sized berries and receptacles (Gindro et al. 2005). Real-time quantitative PCR (qPCR) assays have been developed for the detection and quantification of *B. cinerea* inoculum (conidia and/or mycelium) from air samples and the surface of ripe berries (Carisse et al. 2014; Diguta et al. 2010) and for the quantification of the colonization of *B. cinerea* in developing grape berries and receptacles (Cadle-Davidson 2008; Saito et al. 2013), grape stamens, and ripe berries (Celik et al. 2009; Hill et al. 2014; Sanzani et al. 2012).

The objectives of the current study were as follows: (i) to optimize a hydrolysis probe-based qPCR assay for the quantification of *B. cinerea* DNA in bunch trash; (ii) to investigate the relationships between the quantity of *B. cinerea* DNA measured by qPCR, and the colonization measured based on the plating method,

[†]Corresponding author: V. Rossi; vittorio.rossi@unicatt.it

The author(s) declare no conflict of interest.

and as measured by sporulation potential (SP) of bunch trash using microscope counts of conidia; and (iii) to evaluate the qPCR assay under vineyard conditions.

MATERIALS AND METHODS

Real-time qPCR optimization. *Fungal isolates.* Strains of *B. cinerea* belonging to the transposon genotypes *transposa* (isolate 213 T) and *vacuma* (isolate 351 V) (Ciliberti et al. 2016) and other fungal isolates (Table 1) were obtained from the culture collection of the Department of Sustainable Crop Production of the Università Cattolica del Sacro Cuore (UCSC) in Piacenza, Italy. The biotrophic pathogens *Plasmopara viticola* and *Erysiphe necator* were collected from symptomatic leaves (cultivar Merlot) in the vineyard of UCSC in 2017 and 2018 and were maintained on container-grown grape plants (Merlot) in a greenhouse kept at a temperature of $24 \pm 3^\circ\text{C}$ with a 12-h photoperiod.

Plant material. Bunch trash was obtained from plants grown in a greenhouse to minimize natural colonization by *B. cinerea*. Woody cuttings were collected in winter from an experimental vineyard (*V. vinifera* 'Merlot') at UCSC. Cuttings were grown in a greenhouse maintained at $24 \pm 3^\circ\text{C}$ with a 12-h photoperiod, and flowers were obtained following the technique of Mullins and Rajasekaran (1981). At full flowering, bunch trash was collected by gently shaking the inflorescences inside paper bags. Bunch trash samples were desiccated at 35 to 40°C for 72 h, and the dry weights were determined. Samples were stored at room temperature until use.

DNA extraction. Genomic DNA was obtained from 15 fungal species (Table 1) and bunch trash samples. Except in the case of *P. viticola* and *E. necator*, genomic DNA was extracted from fresh mycelium (obtained by scraping the surface of 10-day-old-colonies grown on potato dextrose agar [PDA], at 20°C with a 18-h photoperiod). DNA of *P. viticola* and *E. necator* was obtained from leaf discs with lesions showing abundant and fresh sporulation (100 mg of leaf material). In brief, mycelium (100 mg fresh weight) or bunch trash (100 mg dry weight) was placed in 2-ml microcentrifuge tubes containing 100 mg of glass sand (425 to 600 μm in diameter), two glass beads (5 mm in diameter), and 500 μl of cetyl trimethylammonium bromide (CTAB) extraction buffer (2% CTAB, 100 mM of Tris-HCl, pH 8.0, 20 mM of ethylenediaminetetraacetic acid, 1.4 M of NaCl, and 1% polyvinylpyrrolidone). The tubes were placed in a Mixer Mill MM200 (Retsch GmbH, Haan, Germany) for 1 min at 30 cycles/s. Subsequently, a CTAB DNA extraction procedure was conducted as described by Saito et al. (2013). The yield and purity of the extracted DNA was determined using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA). The extracts were adjusted to 10 ng/ μl of DNA for fungal samples and to 20 ng/ μl of DNA for bunch trash samples.

Primers and hydrolysis probes. The qPCR assay was based on two specific primers and a hydrolysis probe (Bc3) designed to target the intergenic spacer region (IGS) of the nuclear ribosomal DNA (Suarez et al. 2005) of *B. cinerea*. To normalize the quantification DNA of *B. cinerea* in plant tissues, two specific primers and a hydrolysis probe (Res) designed to target the *V. vinifera* resveratrol synthase gene I (Valseesia et al. 2005) were used as an internal control, with the fluorescent reporter JOE as a substitute for FAM (6-carboxyfluorescein). The sequences for Bc3 were as follows: 5'-GCT GTA ATT TCA ATG TGC AGA ATC C-3' (forward [Bc3F]), 5'-GGA GCA ACA ATT AAT CGC ATT TC-3' (reverse [Bc3R]); and 5'-6-FAM-TCA CCT TGC AAT GAG TGG-BHQ-1-3' (probe [Bc3P]). The sequences for Res were as follows: 5'-CGA GGA ATT TAG AAA CGC TCA AC-3' (forward [ResF]), 5'-GCT GTG CCA ATG GCT AGG A-3' (reverse [ResR]), and 5'-JOE-TGC CAA GGG TCC GGC CAC C-TAMRA-3' (probe [ResP]).

Singleplex and duplex reactions. Singleplex reaction mixtures contained the 1 $\times$ QuantiTect Multiplex PCR Kit (Qiagen, Milano,

Italy), 150 nM of probe (Bc3P or ResP), 500 nM of each primer (Bc3F/R or ResF/R), and 2 μl of DNA template in a final volume of 10 μl . Duplex reaction mixtures contained the 1 $\times$ QuantiTect Multiplex PCR Kit, 150 nM of the *V. vinifera* probe ResP, 150 nM of the *B. cinerea* probe Bc3P, 100 nM of each *V. vinifera* primer (Res F/R), 500 nM of each *B. cinerea* primer (Bc3F/R), and 2 μl of DNA template in a final volume of 10 μl . Both singleplex and duplex assays were performed using an Applied Biosystems 7300 Real-Time PCR System (Thermo Fisher Scientific Inc.) with an initial incubation at 95°C for 15 min followed by 40 cycles of 95°C for 15 s and 60°C for 45 s.

Specificity. The specificity of the qPCR assay for the detection of *B. cinerea* in grape tissue was determined in a test that included the most common grape pathogens and other fungal species frequently found in grapevines and in air samples (Table 1); several of these species were not assessed by Suarez et al. (2005).

Standard curves, calibration DNA, and qPCR optimization. Standard curves were obtained from two singleplex qPCR assays: the Res assay with DNA of *V. vinifera* from bunch trash as a template in a 10-fold dilution series (from 20 to 0.02 ng/ μl), and the Bc3 assay with DNA of *B. cinerea* extracted from the mycelium (isolate 213T, Ciliberti et al. 2016) as a template in a 10-fold dilution series (from 10 to 0.001 ng/ μl). Singleplex qPCR assays were carried out twice; for each assay, each dilution was replicated three times. A water control was included in triplicate in each assay. To detect any potential inhibition of amplification of *B. cinerea* DNA by *V. vinifera* DNA, duplex qPCR assays were performed with DNA of *B. cinerea* mixed with DNA of *V. vinifera*, following the approach described by Saito et al. (2013). In brief, 1 μl from each of the previously described DNA dilutions for *B. cinerea* was mixed with 1 μl of *V. vinifera* DNA (20 ng/ μl), yielding a 10-fold dilution series from 1:2 to 1:20,000 w/w of *B. cinerea*/*V. vinifera* DNA. The duplex qPCR assay was performed twice with three replicates for each dilution. Standard curves of both singleplex and duplex qPCR assays were produced by linear regression, and the coefficient of determination (R^2) was calculated. The amplification efficiency (E) of all assays was determined from the slope of the standard curves (Bustin et al. 2009). To allow comparisons among the results of different tests, all duplex qPCR plates contained a calibration DNA template in triplicate consisting of 1 ng/ μl of *B. cinerea* DNA diluted in 20 ng/ μl of *V. vinifera* DNA (1:20, wt/wt, of *B. cinerea*/*V. vinifera* DNA).

Colonization coefficient. To quantify DNA of *B. cinerea* in bunch trash, known numbers of conidia or known weights of fresh mycelium were added to noncolonized bunch trash; the trash was obtained from the plants grown under isolation, and the absence

TABLE 1. List of isolates screened during specificity tests of the real-time quantitative polymerase chain reaction (qPCR) assay used to quantify *Botrytis cinerea* in grape bunch trash

Genus and species	Isolate code	qPCR result ^a
<i>Alternaria alternata</i>	5	–
<i>Alternaria</i> sp.	23	–
<i>Aspergillus flavus</i>	4	–
<i>Aspergillus niger</i>	A1	–
<i>B. cinerea</i>	213T and 351V	+
<i>Erysiphe necator</i>	FP 2017 and FP 2018 ^b	–
<i>Guignardia bidwellii</i>	Q15 and C14	–
<i>Monilia laxa</i>	11	–
<i>Penicillium</i> sp.	2	–
<i>Phomopsis viticola</i>	Pho-1 and Pho-6	–
<i>Plasmopara viticola</i>	FP 2017 and FP 2018	–
<i>Rhizopus</i> sp.	26	–
<i>R. stolonifer</i>	MUCL38013	–
<i>Sclerotinia sclerotiorum</i>	22	–
<i>Stemphylium</i> sp.	14	–

^a Minus signs indicate not amplified, and the plus sign indicates amplified.

^b FP = field population and year of collection.

of *B. cinerea* was confirmed by a Bc3 qPCR assay. Conidial suspensions (10 µl containing from 1 to 1,250 conidia/µl) or fresh mycelia (5.0, 1.0, 0.5, or 0.1 mg) were added to 0.1-g bunch trash samples in 2-ml microcentrifuge tubes. The total DNA from two biological replicates of each sample was then extracted as described earlier; noncolonized bunch trash without the addition of *B. cinerea* was used as a negative control. Duplex qPCR assays were performed twice for each experiment with two technical replicates of each template DNA. A water control and calibration DNA were included in each assay.

The quantification of DNA of *B. cinerea* in the presence of DNA of *V. vinifera* was expressed in terms of a colonization coefficient (CC), which was the ratio between DNA concentrations of *B. cinerea* and *V. vinifera*, corrected by a correction coefficient (ΔCC) (Gusberti et al. 2012). CC values were calculated as follows:

$$CC = \text{DNA}_{B. cinerea} / \text{DNA}_{V. vinifera} + \Delta CC \quad (1)$$

DNA amounts were obtained by transforming the quantification cycles (C_q) values of both targets (*B. cinerea* and *V. vinifera*) according to the standard curves obtained from the serial dilution assays (Table 2) as follows:

$$\text{DNA (ng/}\mu\text{l)} = 10^{[(C_q \text{ value} - y\text{-axis intercept})/\text{slope}]} \quad (2)$$

ΔCC was calculated as the difference between the average CC value of calibrator DNA calculated in the standard curve (CC_{St} = 5.15) and the average CC value of calibrator DNA of the assay (CC_A) (i.e., ΔCC = CC_{St} - CC_A).

Evaluation of the qPCR method with inoculated bunch trash. *Plant material.* Bunch trash samples were collected in 2017 in a vineyard located in Castell'Arquato (CA) in the Emilia-Romagna region of Northern Italy (44°51'26.1"N 9°51'20.7"E, 400 m above sea level). The CA vineyard was planted to Merlot, which is highly susceptible to *B. cinerea* (Bisiach et al. 1996; Corvi and Tullio 1980). The vines were 10 years old and were trained using the Guyot system. The within and between-row spacings were 1.0 and 2.3 m, respectively. The vineyard was managed following an integrated pest management (IPM) program, with between-row grass, branches pruned to 10 to 12 buds per cane, and no irrigation. Vines were not treated for control of *B. cinerea*. At full flowering (stage 65 of Lorenz et al. 1994), bunch trash was collected from the vines by gently shaking grape bunches inside paper bags. Bunch trash samples were transported to the laboratory and were immediately desiccated at 35 to 40°C for 72 h, and the dry weights were determined. Samples were stored at room temperature.

Preparation of inoculum of B. cinerea and inoculation of bunch trash. Conidia of *B. cinerea* (isolate 213T) were obtained from 10-day-old cultures grown on PDA. The conidial suspensions were prepared by flooding the dishes with sterile-distilled water and gently scraping the agar surface with a sterile rod. The suspension was filtered through two layers of autoclaved gauze and quantified using a hemocytometer. The inoculum concentration was adjusted to 10⁵ conidia/ml.

The samples of bunch trash (0.1 g) collected in the vineyard were placed on autoclaved filter paper discs in Petri dishes (60 mm in diameter) and inoculated with 1 ml of the conidial suspension of *B. cinerea* by using a micropipette. The samples were incubated at

20°C for 18 h in the dark to favor conidial germination and bunch trash colonization. The colonized samples were dried in a laminar flow hood at room temperature for 2 h. Bunch trash samples with different degrees of colonization by *B. cinerea* (0, 25, 50, 75, and 100%) were obtained by mixing colonized and noncolonized bunch trash; for example, 75% colonization comprised 0.75 g of colonized bunch trash and 0.25 g of noncolonized bunch trash. Three replicate 1.0-g samples were prepared for each colonization level.

Colonization of inoculated bunch trash as determined by qPCR. In the first assay, genomic DNA was extracted from 0.1 g of two replicate samples for each of the five bunch trash colonization levels. The extracted DNA was quantified by the duplex qPCR assay described earlier. A water control and calibration DNA were included in each assay. The quantity of DNA of *B. cinerea* in the presence of DNA of *V. vinifera* was expressed as a CC value.

Colonization of inoculated bunch trash as determined by plating. In the second assay, colonization of inoculated bunch trash by *B. cinerea* was quantified for three replicate samples of each of the five colonization levels by randomly and individually plating 50 pieces (stamens, aborted flowers, aborted berries, calyptras, tendrils, or leaf fragments) on PDA in Petri dishes (90 mm in diameter). The dishes were incubated at 20°C with an 18-h photoperiod for 3 days. The dishes were examined using a stereomicroscope, and the colonization rate (CR) was expressed as the percentage of pieces with characteristic grayish sporulation indicating the growth of *B. cinerea*.

SP on inoculated bunch trash as determined by incubation and spore enumeration. In the third assay, the SP of *B. cinerea* on inoculated bunch trash was determined for three replicate samples (0.05 g each) at each colonization level. The bunch trash was placed on a disc of autoclaved filter paper in Petri dishes (60 mm in diameter); sterile water (0.5 ml per dish) was used to moisten the filter paper to maintain a saturated atmosphere. The dishes were sealed with Parafilm and incubated at 20°C with an 18-h photoperiod to induce sporulation of *B. cinerea*. After 3 days of incubation, the bunch trash was suspended in 5 ml of sterile water in a 15-ml Falcon tube and mixed with a vortex apparatus for 10 s. Conidia of *B. cinerea* were counted using a hemocytometer and expressed as the number of conidia per gram of dry bunch trash.

The bunch trash inoculation experiment and quantification of *B. cinerea* by qPCR, plating, and sporulation was performed three times.

Evaluation of the qPCR method with naturally inoculated bunch trash. *Vineyards and treatments.* Experiments were conducted in the CA vineyard and in two additional vineyards (designated MA and CO) located in Northern Italy in 2016 and 2017. The MA vineyard (44°41'57"N 12°19'66"E, at sea level) is located in Mandriole in the Emilia-Romagna region, and the CO vineyard (45°57'05"N 13°27'19"E, 1 m at sea level) is located in Cormons in the Friuli-Venezia Giulia region of Italy. The MA vineyard was planted with cultivar Trebbiano Romagnolo, which were trained using the Casarsa system. Vines were 12 years old in 2016. The CO vineyard was planted with cultivar Merlot, which were trained using the Guyot system. Vines were 7 years old in 2016. Within- and between-row spacing in the MA and CO vineyards was 1.0 and 3.0 m and 0.8 and 2.4 m, respectively. The MA vineyard was managed according to standard IPM practice in

TABLE 2. Linear regression results and reaction efficiencies (E) for the relationship between serially diluted DNA concentrations (log transformed) of *Botrytis cinerea* and *Vitis vinifera* and corresponding C_q values obtained in singleplex and duplex quantitative polymerase chain reaction (qPCR) assays

qPCR assay	DNA template	Linear equation ^a	R ^{2b}	P value ^c	E (%)
Singleplex Bc3	<i>B. cinerea</i>	y = -3.42x + 22.5	0.99	<0.001	96
Singleplex Res	<i>V. vinifera</i>	y = -3.37x + 23.7	0.99	<0.001	98
Duplex Bc3/Res	<i>B. cinerea</i> + <i>V. vinifera</i>	y = -3.56x + 24.0	0.99	<0.001	91

^a In the equations, y refers to the C_q value, and x refers to the DNA concentration.

^b R² is the coefficient of determination of the regression.

^c P value indicates fit of the regression model.

this region (Ministero delle politiche agricole alimentari, forestali e del turismo 2017), with between-row grass, vines pruned to 10 to 12 buds per cane, and emergency irrigation. The CO vineyard followed a conventional pest management strategy, with between-row grass, vines long pruned, and irrigation. Like Merlot in the CA and CO vineyards, Trebbiano Romagnolo in the MA vineyard is highly sensitive to *B. cinerea* (Bisiach et al. 1996; Corvi and Tullio 1980).

In each of the three vineyards, fungicide treatments were applied to obtain a range of colonization of bunch trash by *B. cinerea*. There were two treatments: (i) nontreated control (NT), and (ii) fungicide applied at full flowering (stage 65) (T). Treatments were arranged in a complete randomized block design with four replicate plots per treatment and with six plants per plot. The T treatment was a commercial mixture of fludioxonil (25%) and cyprodinil (37.5%) (Switch; Syngenta Crop Protection) applied at 0.8 g/liter of water until run-off using a 15-liter Elettropius knapsack sprayer (Davide e Luigi Volpi S.p.a, Casalromano, Italy). Seven days after the treatment, bunch trash was collected from five random bunches per plot; these were combined to yield four replicate trash bunches per treatment.

Colonization of naturally inoculated bunch trash as determined by qPCR. Genomic DNA in the naturally inoculated bunch trash was extracted from each of the four replicate samples per treatment (0.1 g each) and was quantified by the duplex qPCR assay as described previously; there were two technical replicates of each template DNA. A water control and calibration DNA were included in each assay. The quantities of DNA of *B. cinerea* in the presence of DNA of *V. vinifera* were expressed as CC values.

SP on naturally inoculated bunch trash as determined by incubation and spore enumeration. The SP of *B. cinerea* in the naturally inoculated bunch trash was determined for each of the four replicate samples per treatment by wrapping the bunch trash in three layers of sterile filter paper to which 5 ml of sterile water was added. The bunch trash in filter papers was sealed in polyethylene bags and incubated at 20°C for 5 days to induce sporulation in *B. cinerea*. Each sample of bunch trash was suspended in 15 ml of sterile water in a 50-ml Falcon tube and vortexed. The conidia of *B. cinerea* were counted using a hemocytometer and the quantity was expressed as the number of conidia per gram of dry bunch trash.

Latent infection of naturally inoculated berries. The incidence of latent infection (ILI) of berries by *B. cinerea* was assessed in the three vineyards; 25 randomly selected, symptomless berries with the pedicel attached were collected at maturity (stage 89) in each replicate plot in both years. The berries were rinsed in tap water, surface sterilized by immersion for 1 min in a 30% sodium hypochlorite solution, and rinsed in sterile-distilled water for 1 min. Berries were positioned individually over a metal grid that was placed in a sterile metal box, the bottom of which was covered with wet, sterile paper. The boxes were sealed in plastic bags to maintain a saturated atmosphere and were incubated for 7 days at 25°C. The ILI was visually assessed as the percentage of berries showing typical sporulation of *B. cinerea*.

Data analysis. All statistical analyses were performed using SPSS (version 24; IBM SPSS Statistics, IBM Corp., Armonk, NY). For experiments with inoculated bunch trash, regression analysis was used to investigate the relationships between the number of conidia of *B. cinerea* or the quantity of mycelium added and the CC, and between the CC and the CR of bunch trash or the SP on bunch trash. Both linear and nonlinear regression functions were used to explore these relationships:

$$Y = a + bX \quad (3)$$

$$Y = Y_{\max}/(1 + \exp[a - bX]) \quad (4)$$

in which a and b are intercept and slope parameters and $Y_{\max}$ is the maximum value of Y in the experiments. The coefficient of

determination (R^2) was used to assess the strength of the relationship.

The data from the field experiment were subject to a factorial analysis of variance (ANOVA) to determine whether the quantity of DNA of *B. cinerea* in bunch trash, SP, and ILI were affected by the main effects of year (2016 and 2017), vineyard (CA, CO, and MA), treatment (T and NT), and their interactions. Prior to the ANOVA, the SP and ILI values were transformed by natural logarithm and arcsine functions, respectively, to ensure homogeneity of variances.

RESULTS

qPCR specificity and standard curves. The Bc3 probe/primer set did not amplify the purified DNA of nontarget organisms but did amplify the purified DNA of *B. cinerea* (Table 1), demonstrating excellent specificity. In the singleplex qPCR reaction, the *B. cinerea* standard curve revealed a high reaction efficiency of 96%, with a close relationship between the Cq values and the concentrations of DNA of *B. cinerea* obtained by dilution (Table 2). The Bc3 assay was able to amplify the lowest concentration of DNA of *B. cinerea* tested (0.001 ng/μl), demonstrating excellent sensitivity. In the duplex assay, the presence of grape DNA did not influence the sensitivity or coefficient of determination (R^2 value), whereas the reaction efficiency of the Bc3 set was slightly reduced (Table 2). Similar results were obtained for the *V. vinifera* standard curve. The Res assay was able to amplify the lowest concentration of DNA of *V. vinifera* tested (0.02 ng/μl).

Evaluation of the qPCR assay using inoculated bunch trash. The CC values, which represented the quantity of DNA of *B. cinerea* detected in the presence of DNA of *V. vinifera*, were proportional to the number of conidia of *B. cinerea* added to the bunch trash (Fig. 1A) and to the quantity of mycelium of *B. cinerea* added to the bunch trash (Fig. 1B); the coefficients of determination indicated a strong linear regression between these variables ($R^2 = 0.92$ and 0.97 , respectively). When the qPCR assay was used with bunch trash samples containing different proportions of non-colonized bunch trash and *B. cinerea*-colonized bunch trash, the CC values were strongly related to the percentage of bunch trash colonized by *B. cinerea* (Fig. 2A) ($R^2 = 0.93$).

For inoculated bunch trash, the relationship between the CC value and the SP (as determined by incubation followed by microscopic counting of spores) and between the CC value and the CR (as determined by plating bunch trash pieces) was nonlinear (Fig. 2B and C). As CC increased, both SP and CR increased, but the rate of increase of SP and CR declined while CC continued to increase, resulting in a logistic relationship. The coefficients of determination indicated a strong relationship these variables ($R^2 = 0.92$ and 0.97 , respectively).

Evaluation of the qPCR assay using naturally colonized bunch trash. The application of fungicides to control *B. cinerea* at flowering significantly reduced the CC (Fig. 3A), SP (Fig. 3B), and ILI on berries (Fig. 3C) ($P < 0.001$ for all three variables; Table 3). Year had a significant effect on CC ($P \leq 0.001$): the degree of colonization of bunch trash was greater in 2017 ($CC = 1.76 \pm 0.45$) compared with colonization in 2016 ($CC = 0.23 \pm 0.09$). Main effects of year also affected SP ($P < 0.0001$), but not ILI ($P = 0.3$). The main effect of vineyard affected SP ($P = 0.02$) and ILI ($P \leq 0.0001$), but not CC ($P = 0.7$). In addition, SP was significantly influenced by the interaction of year $\times$ vineyard, and ILI by the interactions of year $\times$ vineyard and year $\times$ treatment. Thus, the fungicide treatment reduced CC, SP, and ILI in all of the vineyards, irrespective of the initial degree of colonization of bunch trash by *B. cinerea*, the SP of the bunch trash, or the ILI of berries.

The interaction of year $\times$ vineyard $\times$ treatment had no significant effect on CC ($P = 0.6$), SP ($P = 0.5$), or ILI ($P = 0.8$), demonstrating that beyond the two-way interactions of main effects, there were no

more complex associations in this study. These results indicate that the CC values reflected the reduction in sporulation of *B. cinerea* and the latent infection of berries caused by fungicide application at flowering. The qPCR method was able to detect differences between treated and nontreated plots in vineyards whether the CC of bunch trash by *B. cinerea* was <0.1 (Fig. 4A), <3 (Fig. 4B), or <5 (Fig. 4C).

DISCUSSION

We evaluated a qPCR assay for the quantification of *B. cinerea* DNA in grape bunch trash. The qPCR assay was based on the procedure developed by Saito et al. (2013), with a few adaptations regarding the handling of plant material prior to DNA extraction and the qPCR reaction mixture and conditions. The qPCR assay was compared with traditional mycological techniques for quantifying

Fig. 1. Relationship between the colonization coefficient (the ratio between DNA concentrations of *Botrytis cinerea* and *Vitis vinifera* as determined by the quantitative polymerase chain reaction assay) and **A**, the number of conidia added per gram of bunch trash, and **B**, the weight of fresh mycelium added to the bunch trash in inoculation experiments. Markers indicate means, whiskers indicate standard errors, and dotted lines indicate the linear relationships. In A, $Y = 0.011 + 0.001X$ ($R^2 = 0.92$); and in B, $Y = 1.75 + 710.74X$ ($R^2 = 0.97$).

Fig. 2. Relationship between the colonization coefficient (the ratio between DNA concentrations of *Botrytis cinerea* and *Vitis vinifera* as determined by the quantitative polymerase chain reaction assay) of grape bunch trash and **A**, the proportion (%) of bunch trash that had been inoculated with and colonized by *B. cinerea*, **B**, the sporulation potential of the bunch trash (expressed as the number of conidia produced per gram of bunch trash as determined by incubating the bunch trash in humid chambers), and **C**, bunch trash colonization (expressed as the proportion [%] of bunch trash pieces that were colonized by *B. cinerea* as determined by plating the bunch trash on potato dextrose agar). The grape bunch trash was inoculated with *B. cinerea* prior to being assayed for DNA of *B. cinerea*, sporulation potential, and colonization rate. Markers indicate means, whiskers indicate standard errors, and dotted lines indicate linear and nonlinear relationships. In A, $Y = 1.79 + 13.81X$ ($R^2 = 0.93$); in B, $Y = 95.5/(1 + \exp[3.38 - 1.19X])$ ($R^2 = 0.99$); and in C, $Y = 1.17E+07/(1 + \exp[4.31 - 1.41X])$ ($R^2 = 0.99$).

B. cinerea based on colonization of and sporulation on bunch trash. Our results indicate that the qPCR assay and the CC calculation provide a sensitive and reliable method for quantifying colonization by *B. cinerea* of the trash materials (stamens, aborted flowers, aborted berries, calyptas, tendrils, and leaf pieces) remaining in grape bunches after flowering.

Colonized bunch trash serves as one of the primary sources of inoculum for the infection of ripening berries (Elmer and Michailides 2007; Holz et al. 2003; Nair et al. 1995). Thus, quantitative assessment of bunch trash colonization by *B. cinerea* is important for both research purposes and practical disease management, in order to make decisions regarding control of BBR. When the bunch trash has a low incidence of *B. cinerea*, subsequent development of BBR during berry ripening is likely to be low too (Keller et al. 2003; McClellan and Hewitt 1973; Pezet et al. 2003), and the number of fungicide applications can therefore be reduced (González-Domínguez et al. 2019). Visual assessment, plating on agar media, and microscope counts of spores have been commonly used to evaluate the colonization and the SP of *B. cinerea* in bunch trash (Abdelwahab and Younis 2012; Calvo-Garrido et al. 2014; Jaspers et al. 2013; Mundy et al. 2012); these traditional techniques are time-consuming and require expertise for the identification of *B. cinerea* colonies and/or conidia.

The results of Suarez et al. (2005) and those of our study indicate that the qPCR assay is highly specific to *B. cinerea*. We found that the Bc3 system amplified the DNA of different *B. cinerea* strains, including strains that belong to the transposon genotypes *transposa* (T) or *vacuma* (V) (Ciliberti et al. 2016), but did not amplify the DNA of phylogenetically related species (*B. fabae*, *Monilia laxa*, and *Sclerotinia sclerotiorum*). Also, the Bc3 system did not amplify the DNA of other common grapevine pathogens (*E. necator*, *Guignardia bidwellii*, *Phomopsis viticola*, and *Plasmopara viticola*) or of other fungal species frequently present in vineyards (*Alternaria* spp., *Aspergillus* spp., *Penicillium* spp., and *Rhizopus* spp.).

In addition to its specificity, the qPCR assay is also sensitive because the targeted IGS region is a multicopy gene (Bruns et al. 1991). The lowest DNA concentration of *B. cinerea* tested in this study (0.001 ng/μl) was amplified in both the singleplex Bc3 assay and the duplex Bc3/Res assay in the presence of grape DNA, which is consistent with the results obtained by Saito et al. (2013) and Hill et al. (2014). Suarez et al. (2005) showed that the Bc3 assay is able to detect DNA concentrations as low as 20 fg/μl.

The results of the qPCR assay were used to calculate the CC (i.e., the ratio of the pathogen and host DNA concentrations) (Gusberti et al. 2012). In duplex qPCR analyses, researchers have described several methods for calculating the quantity of DNA of a pathogen in host tissue. To account for variation among samples and qPCR runs in terms of tissue weight, pipetting volumes, and efficiencies of DNA extraction and amplification, these methods account for amounts of host plant DNA in order to provide internal normalization. Sanzani et al. (2012), for example, normalized the DNA concentration of *B. cinerea* according to the quantity of host DNA by using a host DNA correction factor for each grape sample. Valsesia et al. (2005) developed the infection coefficient (IC), which is based on the ratio between Cq values of the pathogen and host generated by the qPCR assay. The IC approach was also used to determine the pathogen coefficient (PC) of *B. cinerea* in grape berries and receptacles (Hill et al. 2014; Saito et al. 2013). In a preliminary analysis, we found that the PC values based on the current data increased as the number of *B. cinerea* conidia or quantity of mycelium added to bunch trash increased. However, in the inoculation experiments, the PC values did not increase linearly as the level of colonization increased from 0 to 100% (data not shown). These preliminary results generally agree with those of Saito et al. (2013) and Hill et al. (2014), who found that the accuracy of PC decreased as the

Fig. 3. Effect of the application of fungicides (a mixture of fludioxonil [25%] and cyprodinil [37.5%]) to control *Botrytis cinerea* at flowering on **A**, the colonization coefficient (the ratio between DNA concentrations of *B. cinerea* and *Vitis vinifera* in naturally colonized grape bunch trash as determined by the quantitative polymerase chain reaction assay), **B**, the sporulation potential of bunch trash (expressed as the number of conidia produced per gram of naturally inoculated bunch trash after incubation in humid chambers), and **C**, the incidence of grape berries with latent infection (expressed as the percentage of berries showing the characteristic sporulation of *B. cinerea*). Bars indicate means of treated (T) and nontreated (NT) plots in three vineyards from 2 years of data, and whiskers indicate standard errors ($n = 24$; three vineyards, 2 years, four replicates). In each panel, means are significantly different ($P < 0.001$).

severity of BBR increased. For this reason, we used the CC rather than the PC in the current study.

The results of the qPCR assay were comparable to those obtained with the traditional methods used to estimate *B. cinerea* colonization of and sporulation on bunch trash. Therefore, the qPCR assay described in this work is a valuable alternative to the traditional methods. Traditional methods and qPCR require expertise in mycology and molecular biology, respectively. The traditional methods are time-consuming, whereas the qPCR assay takes 3 to 4 h. They also have limitations that potentially reduce their accuracy. Plating of trash pieces on agar can lead to the growth of other fast-growing fungal species that may result in lower estimates of the number of *B. cinerea* colonies, while the accurate enumeration of spores from incubated trash in humid chambers is dependent on the operator's expertise to correctly identify conidia. The qPCR assay, on the contrary, provides sensitive and specific results.

The validity of the qPCR assay was confirmed in the field using bunch trash naturally colonized with *B. cinerea* that had been treated or not treated with fungicides during flowering. In the field,

the CC values were consistent with the reduction of the SP caused by fungicide treatment and revealed differences between fungicide-treated plants and nontreated plants under different environmental conditions, even in situations where the incidence of colonization of bunch trash by *B. cinerea* was very low. The CC values of bunch trash were also consistent with the reduction in the ILI of berries caused by fungicide application at flowering. Although the latter result requires confirmation, it suggests that when the colonization of bunch trash is low, the ILI is also low as a result of unfavorable conditions for reproduction of *B. cinerea* during flowering.

In conclusion, the qPCR methodology described here is a sensitive, specific, and reliable tool for quantifying *B. cinerea* in bunch trash in vineyards. The qPCR assay can be used as an alternative to traditional methods for the quantification of *B. cinerea* during the early-season period (as an indicator of inoculum potential) and thus BBR severity at harvest; it can also be used as a tool in other epidemiological studies and to determine the effect of disease management methods on the reduction of inoculum of *B. cinerea*.

TABLE 3. Results of the analysis of variance performed to explore main effects of fungicide treatment, vineyard and year, and main effect interactions on the quantity of DNA of *Botrytis cinerea* in bunch trash, the sporulation potential of bunch trash, and the incidence of grape berries with latent infection^a

Main effects and interactions	df ^b	Colonization coefficient		Sporulation potential		Incidence of latent infections	
		F value	P value	F value	P value	F value	P value
1. Year	1	17.4	<0.001	2,309.3	<0.001	1.3	0.3
2. Vineyard	2	0.4	0.7	4.4	0.02	78.5	<0.001
3. Treatment	1	17.8	<0.001	50.6	<0.001	54.9	<0.001
1 × 2	2	1.7	0.2	21.8	<0.001	8.4	0.001
1 × 3	1	2.7	0.08	1.8	0.2	4.8	0.04
2 × 3	2	0.2	0.8	2.1	0.1	2.6	0.09
1 × 2 × 3	2	0.6	0.6	0.8	0.5	0.2	0.8

^a The quantity of DNA of *B. cinerea* in bunch trash was defined as the colonization coefficient determined by the quantitative polymerase chain reaction assay, the sporulation potential of bunch trash was defined as the natural logarithm of the number of conidia produced per gram of naturally inoculated bunch trash after incubation in humid chambers, and the incidence of grape berries with latent infection was defined as the arcsin transformation of the percentage of berries showing the characteristic sporulation of *B. cinerea*. Samples were collected from a field experiment comparing grape bunches either treated with a commercial mixture of fludioxonil (25%) and cyprodinil (37.5%) (Switch; Syngenta Crop Protection) applied at 0.8 g/liter of water or not treated, with the experiment conducted in 2016 and 2017 in three vineyards in Italy.

^b df = degrees of freedom.

Fig. 4. Ability of the quantitative polymerase chain reaction (qPCR) assay to detect differences in the natural colonization by *Botrytis cinerea* of fungicide-treated (T) and nontreated (NT) vines when the colonization coefficient of grape bunch trash was A, <0.1, B, <3, and C, <5. The data for A, B, and C were from the Castell'Arquato vineyard in 2017, the Cormons vineyard in 2016, and the Mandriole vineyard in 2016, respectively. Fungicides (fludioxonil [25%] and cyprodinil [37.5%]) were applied at flowering. Bars indicate means of the colonization coefficient (the ratio between DNA concentrations of *B. cinerea* and *Vitis vinifera* in naturally inoculated grape bunch trash as determined by the qPCR assay), and whiskers indicate standard errors ($n = 4$; based on four replicates). In all panels, treatments are significantly different at $P < 0.001$. The interaction of year $\times$ vineyard $\times$ treatment was not significant ($P = 0.6$).

ACKNOWLEDGMENTS

G. Fedele conducted this study within the Doctoral School of the Agro-Food System (Agrisystem) at UCSC.

LITERATURE CITED

- Abdelwahab, H., and Younis, R. A. 2012. Early detection of gray mold in grape using conventional and molecular methods. *Afr. J. Biotechnol.* 11: 15241-15245.
- Bisiach, M., Zerbetto, F., and Cortesi, P. 1996. Attività fungicida della miscela ciprodinil + fludioxonil contro *Botrytis cinerea* su vite da vino. *ATTI Giornate Fitopatologiche* 2:363-368.
- Bruns, T. D., White, T. J., and Taylor, J. W. 1991. Fungal molecular systematics. *Annu. Rev. Ecol. Syst.* 22:525-564.
- Bustin, S. A., Benes, V., Garson, J. A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M. W., Shipley, G. L., Vandesompele, J., and Wittwer, C. T. 2009. The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* 55:611-622.
- Cadle-Davidson, L. 2008. Monitoring pathogenesis of natural *Botrytis cinerea* infections in developing grape berries. *Am. J. Enol. Vitic.* 59:387-395.
- Calvo-Garrido, C., Usall, J., Viñas, I., Elmer, P. A., Cases, E., and Teixidó, N. 2014. Potential secondary inoculum sources of *Botrytis cinerea* and their influence on bunch rot development in dry Mediterranean climate vineyards. *Pest Manag. Sci.* 70:922-930.
- Carisse, O., Tremblay, D. M., and Lefebvre, A. 2014. Comparison of *Botrytis cinerea* airborne inoculum progress curves from raspberry, strawberry and grape plantings. *Plant Pathol.* 63:983-993.
- Celik, M., Kalpulov, T., Zutahy, Y., Ish-shalom, S., Lurie, S., and Lichter, A. 2009. Quantitative and qualitative analysis of *Botrytis* inoculated on table grapes by qPCR and antibodies. *Postharvest Biol. Technol.* 52:235-239.
- Ciliberti, N., Fermaud, M., Languasco, L., and Rossi, V. 2015a. Influence of fungal strain, temperature, and wetness duration on infection of grapevine inflorescences and young berry clusters by *Botrytis cinerea*. *Phytopathology* 105:325-333.
- Ciliberti, N., Fermaud, M., Roudet, J., Languasco, L., and Rossi, V. 2016. Environmental effects on the production of *Botrytis cinerea* conidia on different media, grape bunch trash, and mature berries. *Aust. J. Grape Wine Res.* 22:262-270.
- Ciliberti, N., Fermaud, M., Roudet, J., and Rossi, V. 2015b. Environmental conditions affect *Botrytis cinerea* infection of mature grape berries more than the strain or transposon genotype. *Phytopathology* 105:1090-1096.
- Corvi, F., and Tullio, V. 1980. Un biennio di prove di lotta contro la muffa grigia dell'uva (*Botrytis cinerea* Pers.) nelle marche (II contributo). *Giornate Fitopatol.* 2:553-560.
- Diguta, C. F., Rousseaux, S., Weidmann, S., Bretin, N., Vincent, B., Guilloux-Benatier, M., and Alexandre, H. 2010. Development of a qPCR assay for specific quantification of *Botrytis cinerea* on grapes. *FEMS Microbiol. Lett.* 313:81-87.
- Edwards, S. G., and Seddon, B. 2001. Selective media for the specific isolation and enumeration of *Botrytis cinerea* conidia. *Lett. Appl. Microbiol.* 32:63-66.
- Elad, Y., Williamson, B., Tudzynski, P., and Delen, N., eds. 2007. *Botrytis: Biology, Pathology and Control*. Springer, Dordrecht, The Netherlands.
- Elmer, P. A. G., and Michailides, T. J. 2007. Epidemiology of *Botrytis cinerea* in orchard and vine crops. Pages 243-272 in: *Botrytis: Biology, Pathology and Control*. Y. Elad, B. Williamson, P. Tudzynski, and N. Delen, eds. Springer, Dordrecht, The Netherlands.
- Gindro, K., Pezet, R., Viret, O., and Richter, H. 2005. Development of a rapid and highly sensitive direct-PCR assay to detect a single conidium of *Botrytis cinerea* Pers.: Fr in vitro and quiescent forms in planta. *Vitis Geilweilerhof* 44:139-142.
- González-Domínguez, E., Caffi, T., Ciliberti, N., and Rossi, V. 2015. A mechanistic model of *Botrytis cinerea* on grapevines that includes weather, vine growth stage, and the main infection pathways. *PLoS One* 10:e0140444.
- González-Domínguez, E., Fedele, G., Caffi, T., Delière, L., Sauris, P., Gramaje, D., Ramos-Saez de Ojer, J. L., Díaz-Losada, E., Díez-Navajas, A. M., Bengoa, P., and Rossi, V. 2019. A network meta-analysis provides new insight into fungicide scheduling for the control of *Botrytis cinerea* in vineyards. *Pest Manag. Sci.* 75:324-332.
- Gusberti, M., Patocchi, A., Gessler, C., and Broggin, G. A. L. 2012. Quantification of *Venturia inaequalis* growth in *Malus × domestica* with quantitative real-time polymerase chain reaction. *Plant Dis.* 96:1791-1797.
- Hill, G. N., Evans, K. J., Beresford, R. M., and Damberg, R. G. 2014. Comparison of methods for the quantification of *Botrytis* bunch rot in white wine grapes. *Aust. J. Grape Wine Res.* 20:432-441.
- Holz, G., Gütschow, M., Coertze, S., and Calitz, F. J. 2003. Occurrence of *Botrytis cinerea* and subsequent disease expression at different positions on leaves and bunches of grape. *Plant Dis.* 87:351-358.
- Jaspers, M. V., Seyb, A. M., Trought, M. C. T., and Balasubramaniam, R. 2013. Overwintering grapevine debris as an important source of *Botrytis cinerea* inoculum. *Plant Pathol.* 62:130-138.
- Keller, M., Viret, O., and Cole, F. M. 2003. *Botrytis cinerea* infection in grape flowers: Defense reaction, latency, and disease expression. *Phytopathology* 93:316-322.
- Lorenz, D., Eichhorn, K., Bleiholder, H., Klose, R., Meier, U., and Weber, E. 1994. Phänologische Entwicklungsstadien der Weinrebe (*Vitis vinifera* L. ssp. *vinifera*). *Vitic. Enol. Sci.* 49:66-70.
- McClellan, W. D., and Hewitt, W. B. 1973. Early *Botrytis* rot of grapes: Time of infection and latency of *Botrytis cinerea* Pers. in *Vitis vinifera* L. *Phytopathology* 63:1151-1157.
- Ministero delle politiche agricole alimentari, forestali e del turismo. 2017. Linee guida nazionali di produzione integrata delle colture, 2018. <https://www.reterurale.it/flex/cm/pages/ServeBLOB.php/L/IT/IDPagina/17765>
- Mullins, M. G., and Rajasekaran, K. 1981. Fruiting cuttings: Revised method for producing test plants of grapevine cultivars. *Am. J. Enol. Vitic.* 32: 35-40.
- Mundy, D. C., Agnew, R. H., and Wood, P. N. 2012. Grape tendrils as an inoculum source of *Botrytis cinerea* in vineyards - a review. *N. Z. Plant Prot.* 65:218-227.
- Nair, N., Guilbaud-Oulton, S., Barchia, I., and Emmett, R. 1995. Significance of carry over inoculum, flower infection and latency on the incidence of *Botrytis cinerea* in berries of grapevines at harvest in New South Wales. *Aust. J. Exp. Agric.* 35:1177-1180.
- Pertot, I., Caffi, T., Rossi, V., Mugnai, L., Hoffmann, C., Grando, M. S., Gary, C., Lafond, D., Duso, C., Thiery, D., Mazzoni, V., and Anfora, G. 2017. A critical review of plant protection tools for reducing pesticide use on grapevine and new perspectives for the implementation of IPM in viticulture. *Crop Prot.* 97:70-84.
- Pezet, R., Viret, O., Perret, C., and Tabacchi, R. 2003. Latency of *Botrytis cinerea* Pers.: Fr. and biochemical studies during growth and ripening of two grape berry cultivars, respectively susceptible and resistant to grey mould. *J. Phytopathol.* 151:208-214.
- Saito, S., Dunne, K. J., Evans, K. J., Barry, K., Cadle-Davidson, L., and Wilcox, W. F. 2013. Optimisation of techniques for quantification of *Botrytis cinerea* in grape berries and receptacles by quantitative polymerase chain reaction. *Aust. J. Grape Wine Res.* 19:68-73.
- Sanzani, S. M., Schena, L., De Cicco, V., and Ippolito, A. 2012. Early detection of *Botrytis cinerea* latent infections as a tool to improve postharvest quality of table grapes. *Postharvest Biol. Technol.* 68:64-71.
- Seyb, A., Gaunt, R., Trought, M., Frampton, C., Balasubramaniam, R., and Jaspers, M. 2000. Relationship between debris within grape bunches and *Botrytis* infection of berries. Page 451 in: *Proceedings of the 1998 New Zealand Plant Protection Conference*. New Zealand Plant Protection Society, Auckland, New Zealand.
- Suarez, M. B., Walsh, K., Boonham, N., O'Neill, T., Pearson, S., and Barker, I. 2005. Development of real-time PCR (TaqMan®) assays for the detection and quantification of *Botrytis cinerea* in planta. *Plant Physiol. Biochem.* 43: 890-899.
- Valsesia, G., Gobbin, D., Patocchi, A., Vecchione, A., Pertot, I., and Gessler, C. 2005. Development of a high-throughput method for quantification of *Plasmopara viticola* DNA in grapevine leaves by means of quantitative real-time polymerase chain reaction. *Phytopathology* 95:672-678.
- Viret, O., Keller, M., Jaudzems, V. G., and Cole, F. M. 2004. *Botrytis cinerea* infection of grape flowers: Light and electron microscopical studies of infection sites. *Phytopathology* 94:850-857.
- Wolf, T. K., Baudoin, A. B. A. M., and Martinez-Ochoa, N. 1997. Effect of floral debris removal from fruit clusters on *Botrytis* bunch rot of Chardonnay grapes. *Vitis* 16:27-33.