Real-Time Fluorescent Polymerase Chain Reaction Detection of *Phytophthora ramorum* and *Phytophthora pseudosyringae* Using Mitochondrial Gene Regions

Paul W. Tooley, Frank N. Martin, Marie M. Carras, and Reid D. Frederick

First, third, and fourth authors: U.S. Department of Agriculture-Agricultural Research Service (USDA-ARS), Foreign Disease-Weed Science Research Unit, Ft. Detrick, MD 21702; and second author: USDA-ARS, 1636 East Alisal St., Salinas, CA 93905.
Accepted for publication 24 November 2005.

ABSTRACT

Tooley, P. W., Martin, F. N., Carras, M. M., and Frederick, R. D. 2006. Real-time fluorescent polymerase chain reaction detection of *Phytophthora ramorum* and *Phytophthora pseudosyringae* using mitochondrial gene regions. *Phytopathology* 96:336-345.

A real-time fluorescent polymerase chain reaction (PCR) detection method for the sudden oak death pathogen *Phytophthora ramorum* was developed based on mitochondrial DNA sequence with an ABI Prism 7700 (TaqMan) Sequence Detection System. Primers and probes were also developed for detecting *P. pseudosyringae*, a newly described species that causes symptoms similar to *P. ramorum* on certain hosts. The species-specific primer-probe systems were combined in a multiplex assay with a plant primer-probe system to allow plant DNA present in extracted samples to serve as a positive control in each reaction. The lower limit of detection of *P. ramorum* DNA was 1 fg of genomic DNA, lower

than for many other described PCR procedures for detecting *Phytophthora* species. The assay was also used in a three-way multiplex format to simultaneously detect *P. ramorum*, *P. pseudosyringae*, and plant DNA in a single tube. *P. ramorum* was detected down to a 10^{-5} dilution of extracted tissue of artificially infected rhododendron 'Cunningham's White', and the amount of pathogen DNA present in the infected tissue was estimated using a standard curve. The multiplex assay was also used to detect *P. ramorum* in infected California field samples from several hosts determined to contain the pathogen by other methods. The real-time PCR assay we describe is highly sensitive and specific, and has several advantages over conventional PCR assays used for *P. ramorum* detection to confirm positive *P. ramorum* finds in nurseries and elsewhere.

Additional keywords: *coxI*, *coxII*.

Phytophthora ramorum (Werres, De Cock & Man in't Veld) sp. nov. causes sudden oak death, a serious disease of California oak species such as coast live oak (*Quercus agrifolia*) and tan oak (*Lithocarpus densiflorus*) (44). The pathogen also is widespread in Europe primarily as a pathogen of ornamentals (14,28,40,59, 60). Because of concern that *P. ramorum* may spread eastward and threaten the vast oak forests of the Eastern United States, state, federal, and Canadian regulations were drafted in 2001 that restricted movement of *P. ramorum* hosts out of infested areas of California (7,8,42).

In 2003, new *P. ramorum* outbreaks were reported in nursery stock found in nurseries from Oregon, Washington, Canada, and additional areas of California (22,41; J. Jones, *personal communication*). Also in 2003, a national *P. ramorum* survey was initiated (12). In 2004, several large west coast production nurseries and some smaller nurseries were confirmed to be infested with *P. ramorum*. These facilities shipped over 2 million host plants, of which only a small portion were infected, to 49 states and the District of Columbia (51; J. Jones, *personal communication*). Efforts were made on the part of several agencies including the USDA Animal and Plant Health Inspection Service (APHIS), U.S. Forest Service, and State Departments of Agriculture to track and test the

shipments, monitor for presence of *P. ramorum* in Eastern states, and educate the public about sudden oak death. By the end of 2004, 171 locations (wholesale nurseries and retail outlets) in 20 states were found to contain plants infected with *P. ramorum*. On 22 April 2004, APHIS issued an amended Emergency Order that implemented new restrictions on interstate movement of host nursery stock and associated articles from all commercial nurseries in California that are outside the quarantined area. Nurseries in Oregon and Washington that ship interstate were added to this regulatory oversight on 10 January 2005. This order also listed 31 confirmed hosts of *P. ramorum* (those for which Koch's postulates had been performed) and a list of 37 additional plant species associated with *P. ramorum* because results of culture or polymerase chain reaction (PCR) tests had returned results positive for the pathogen. The host range of *P. ramorum* continues to increase as the pathogen is identified on an ever-widening group of plant species (13,24,31,41; J. Jones, *personal communication*).

In light of the recent movement of *P. ramorum* to the Eastern United States through shipment of nursery stock, the availability of rapid, sensitive and specific *P. ramorum* detection methods are needed. Unequivocal identification of *P. ramorum* is the goal of survey workers, as false identification and/or confusion of *P. ramorum* with other *Phytophthora* species could lead to the development of improper quarantine measures and/or rejection of plant shipments by state inspectors. *P. ramorum* has several distinguishing morphological characters that may be used for identification. It is characterized by semipapillate, deciduous sporangia with short pedicels and high length to width ratios, large chlamydospores, relatively slow growth, and low cardinal temperatures for growth (60). To accurately assess morphological features, however, requires experience in *Phytophthora* identification as

Corresponding author: P. W. Tooley; E-mail address: paul.tooley@ars.usda.gov

* The e-Xtra logo stands for "electronic extra" and indicates that the online version contains supplemental material not included in the print edition. The online version contains a link to F. Martin's website where additional data are presented.

DOI: 10.1094/PHYTO-96-0336

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 2006.

some characteristics often show a continuum among different species. It can also be time consuming, especially when a number of samples have to be processed. Furthermore, it can be difficult to culture the pathogen from infected tissue at certain times of the year (23).

As an adjunct to morphological identification, several molecular procedures for identification and detection of *P. ramorum* have been developed and are in use in various laboratories and state and federal agencies. These include classical PCR methods based on internal transcribed spacer (ITS) regions of ribosomal DNA (13,23,61) and mitochondrial gene regions (39), PCR–single-strand conformation polymorphism analysis (32), and PCR–restriction fragment length polymorphism (RFLP) analysis (38). In 2004, a single nucleotide polymorphism procedure was also developed to allow differentiation among *P. ramorum* isolates from Europe and North America (33). In 2003, APHIS adopted the ITS-based conventional nested PCR method (13) as an accepted protocol for identification of *P. ramorum* and has stated in an amended order dated 22 April 2004 that positive (conventional) nested PCR tests alone may be used to confirm presence of *P. ramorum* and prohibit movement of affected nursery stock, without requiring confirmatory culturing of the pathogen (54).

Real-time PCR is based on the labeling of primers, probes, or amplicon with fluorogenic molecules and allows detection of the target fragment to be monitored while the amplification is in progress (35,46). In 5' fluorogenic real-time PCR (TaqMan), a sequence-specific oligonucleotide probe labeled with a fluorescent reporter and a quencher generates fluorescence at a rate directly proportional to the amount of product amplified in the reaction (26). The method is now being applied to a range of organisms in many different research applications (30,34,35,43,50), including detection and quantification of fungal plant pathogens (1,3,18,19, 21,46,47,48,55,56). For *Phytophthora* species, real-time PCR has been used in studies detecting and quantifying levels of various species in host plants and soil (4,29,47,56).

Several real-time PCR assays have been described for detection of *P. ramorum*. Bilodeau et al. (2) described an assay based on the ITS, β -tubulin, and elicitor regions using TaqMan and SYBR green assays. Hughes et al. (27) described an ITS-based real-time PCR assay for *P. ramorum* that uses TaqMan chemistry and has been adapted for field use with a SmartCycler (Cepheid, Inc., Sunnyvale, CA) instrument. A real-time PCR procedure for detection of *P. ramorum* based on the ITS region using SYBR green has been described by Hayden et al. (23).

Here, we describe the development of a real-time PCR assay for the sudden oak death pathogen *P. ramorum* based upon mitochondrial sequences. In previous work, we characterized the *coxI* and II genes in *Phytophthora* and described a conventional PCR assay for *P. ramorum* (36–39). In this study, we utilize the same primers as the conventional PCR method previously described (39), except with the addition of TaqMan probes specially designed for *P. ramorum*, *P. pseudosyringae*, and plant DNA. Plant primers were used as a positive control to insure that PCR amplification always occurs with DNA extracted from symptomatic samples. The real-time PCR assay we describe provides a sensitive, specific tool for detection of *P. ramorum*, based on a genomic region not used in other *P. ramorum* assays. It offers advantages over conventional PCR procedures as a stand-alone method or confirmatory procedure for workers monitoring for the presence of *P. ramorum* in new geographic regions.

MATERIALS AND METHODS

Cultures and DNA extraction. *Phytophthora* isolates (Table 1) were maintained on rye A agar (9) at 20°C in darkness and all were used to test primer and probe specificity. Genomic DNA was extracted according to Goodwin et al. (20) from 60 mg of

lyophilized mycelium grown on a synthetic medium (63). DNA concentrations were determined with a spectrophotometer (Model ND-1000; Nanodrop Technologies, Wilmington, DE) and by comparison with known DNA standards using agarose gel electrophoresis. Plant genomic DNA was extracted from noninoculated leaves with a Qiagen DNeasy Plant Maxi Kit (Qiagen Inc., Valencia, CA). Leaves of rhododendron 'Cunningham's White' were inoculated with sporangia of *P. ramorum* isolate 0-217 as described by Tooley et al. (53). California bay laurel (*Umbellularia californica*) was artificially inoculated with *P. ramorum*, *P. pseudosyringae*, or both pathogens by placing a 6-mm-diameter agar plug of mycelium on a wound on the leaf and incubating it in a moist chamber for 7 days. Total DNA was extracted by homogenizing two 6-mm-diameter leaf disks from lesions on infected leaves in a Fastprep FP120 instrument (Qbiogene, Inc., Carlsbad, CA) and using a Qbiogene FastDNA Kit according to the manufacturer's instructions.

Field samples from California. Samples of total DNA from symptomatic plants collected from the field were processed at the California Department of Food and Agriculture as described previously (39). The presence of *Phytophthora* spp. was confirmed by plating tissue on differential medium, and DNA was extracted from diseased tissue and tested with the ITS marker system (13) to determine if *P. ramorum* was present. Samples were also assayed with the mitochondrial marker system described in Martin et al. (39). Real-time PCR assays were conducted on 53 samples from 11 hosts in blind fashion; the samples were numbered randomly and the results of culturing and/or conventional PCR were not known until real-time PCR analyses were completed. DNA samples were also diluted 1:10 with sterile water prior to use because undiluted samples some times amplified poorly.

Primers, probes, and PCR conditions. The nucleotide sequences of the gene regions from which primer and probe sequences were designed are as described previously (39). Plant primers FMPI-2b and FMPI-3b (Table 2) were constructed from the mitochondrially encoded cytochrome oxidase I gene and generated a target fragment of 143 bp (39). Species-specific primers for *P. ramorum* (FMPr-1a and FMPr-7) and *P. pseudosyringae* (FMPps-1c and FMPps-2c) amplified spacer sequences between the *coxII* and *coxI* genes and produced amplicons of 134 and 158 bp, respectively (39) (Table 2). Primers were synthesized by Qiagen. The TaqMan probes were labeled at the 5' end with either the fluorescent reporter dye 6-carboxyfluorescein (FAM) or CAL Fluor Orange 560 (CAL Orange) and labeled at the 3' end with the black hole quencher dye (BHQ, Biosearch Technologies, Novato, CA) (Table 2). In multiplex PCR experiments, the plant probe was labeled at the 5' end with TAMRA (*N,N,N'*-tetramethyl-5-carboxyrhodamine) as a reporter dye instead of CAL Orange.

Real-time PCR was performed using an ABI Prism 7700 Sequence Detection System (Perkin Elmer/Applied Biosystems, Foster City, CA) in a total volume of 25 μ l containing 100 pg of DNA template and 1 $\times$ TaqMan Universal Master Mix (Perkin Elmer/Applied Biosystems) with an additional 0.5 mM MgCl₂. Annealing temperature and magnesium concentration were varied to determine optimum levels for amplification (data not shown). For duplex reactions incorporating both *P. ramorum* and plant primers and probes, an additional 75 μ M dNTPs was added, while for single reactions using *P. pseudosyringae* primers, an additional 1.5 mM MgCl₂ was added. Cycling conditions were 50°C for 2 min, 95°C for 10 min, and 60 cycles of 95°C for 15 s and 55°C for 1 min. The FMPr-1a/FMPr-7 and FMPps-1c/FMPps-2c primer combinations were used at a final primer concentration of 1,000 nM and probe concentration of 400 nM, whereas the FMPI2b/FMPI3b (plant) primers were used at a final primer concentration of 100 nM and probe concentration of 80 nM. For multiplex reactions, we used conditions identical to those for

duplex reactions except that 50- μ l reaction volumes were used and the plant probe was at a concentration of 400 nM. A water blank was included as a negative control in each experiment.

Dilution series experiments. Three repeated experiments with two replications each were performed using spectrophotometrically quantified DNA of *P. ramorum* isolate 288 or *P. pseudosyringae* isolate 471 diluted in sterile distilled water. To determine whether the presence of plant DNA affected the DNA dilution series for *P. ramorum*, experiments were performed using a *P. ramorum* DNA dilution series “spiked” with DNA extracted from uninfected azalea cv. Gloria. Two 6-mm-diameter leaf disks were extracted with the Qbiogene FastDNA Kit in a final volume of 100 μ l and diluted 1:10. Two microliters of extract was added to a dilution series of *P. ramorum* DNA from isolate 288 ranging from 10 ng down to 100 ag, and real-time PCR was performed using only the *P. ramorum* primers and probe as well as a two-way multiplex reaction with the *P. ramorum* primers and probe plus the plant primers and probe (three replications each). In addition, dilution series were made from total DNA extracted from infected rhododendron ‘Cunningham’s White’ inoculated as described above. Individual dilution series were constructed from three separate extractions and two experiments were conducted each using dilution series from all three extractions.

Data analysis. Data acquisition and analysis were performed using the TaqMan data worksheet and software according to the manufacturer’s instructions (Applied Biosystems). The cycle threshold (Ct) values for each reaction were calculated automatically by the ABI Prism sequence detection software (version 1.6.3) by determining the PCR cycle number at which the reporter fluorescence exceeded background.

RESULTS

***P. ramorum*-specific primers and probe.** A high level of *P. ramorum* specificity was observed using the primers FMPr-1a and FMPr-7 and the Pr-FAM probe (Table 2) when tested against 45 other species of *Phytophthora* (multiple isolates tested for some species) at a concentration of 100 pg of DNA with an annealing temperature of 55°C (Table 1). Only *P. ramorum* showed a Ct value of less than 30 cycles with other species exhibiting no detection after 60 cycles (Fig. 1A; Table 3). Twenty-five diverse isolates of *P. ramorum* were amplified at a concentration of 100 pg of DNA using primers FMPr-1a and FMPr-7 and the Pr-FAM probe, with Ct values ranging from 22.56 to 28.91 (Table 3). Primers FMPr-1a and FMPr-7 and the FAM probe worked successfully at 55°C, but at 57°C, amplification became inconsistent (data not shown).

TABLE 1. Isolates of *Phytophthora* spp. used in this study

Species	Group ^a	Isolate number ^b	Host	Origin
<i>Phytophthora arecae</i>	II	441 ^{PT} , IMI348342	<i>Theobroma cacao</i>	Indonesia
<i>Phytophthora boehmeriae</i>	II	325 ^{PT} , P1257 ^{MC}	<i>Boehmeria nivia</i>	Papua New Guinea
<i>Phytophthora cactorum</i>	I	384 ^{PT} , NY577	<i>Fragaria</i> $\times$ <i>ananassa</i>	New York
		385 ^{PT} , NY568	<i>Malus sylvestris</i>	New York
<i>Phytophthora cambivora</i>	VI	443 ^{PT} , 33-4-8	<i>Prunus dulcis</i>	California
<i>Phytophthora capsici</i>	II	306 ^{PT} , Pc-m1	<i>Capsicum annuum</i>	New Jersey
<i>Phytophthora cinnamomi</i>	VI	Cn-2 ^{DJM} (A-2 mating type)	<i>Vaccinium</i> spp.	Florida
		446 ^{PT} , 3210 ^{GB}	<i>Castanea</i>	California
		447 ^{PT} , 3267 ^{GB}	<i>Jugulands californica</i>	California
<i>Phytophthora citricola</i>	III	422 ^{PT} , CR4	<i>Cornus</i>	UNK
<i>Phytophthora citrophthora</i>	II	461 ^{PT}	<i>Rhododendron</i> sp.	Oregon
<i>Phytophthora clandestine</i>	I	IMI287317 ^{DC}	<i>Trifolium subterranean</i>	Australia
<i>Phytophthora colocasiae</i>	IV	345 ^{PT} , 1696 ^{MC}	<i>Colocasia esculenta</i>	China
<i>Phytophthora cryptogea</i>	VI	310 ^{PT} , 620 ^{PH}	<i>Pinus lambertiana</i>	Oregon
		389 ^{PT} , NY508 ^{WW}	<i>Prunus avium</i>	California
<i>Phytophthora drechsleri</i>	VI	401 ^{PT} , ATCC 64494	<i>Solanum tuberosum</i>	Egypt
<i>Phytophthora erythroseptica</i>	VI	374 ^{PT}	<i>Solanum tuberosum</i>	Maine
<i>Phytophthora fragariae fragariae</i>	V	398 ^{PT} , 94-96 ^{JIM}	<i>Fragaria</i> $\times$ <i>ananassa</i>	Oregon
<i>Phytophthora gonapodyides</i>	VI	392 ^{PT} , NY414 ^{WW}	<i>Prunus persica</i>	New York
<i>Phytophthora heveae</i>	II	462 ^{PT} , 97-251 ^{PC}	<i>Rhododendron</i> sp.	Oregon
<i>Phytophthora hibernalis</i>	IV	338 ^{PT} , ATCC 56353, 3822 ^{MC}	Citrus	Australia
<i>Phytophthora humicola</i>	V	IMI302303 ^{DC}	Soil from citrus	Taiwan
<i>Phytophthora idaei</i>	I	IDA3 ^{DC} (Type)	<i>Rubus idaeus</i>	Scotland
<i>Phytophthora ilicis</i>	IV	344 ^{PT} , P3939 ^{MC} , ATCC 56615	<i>Ilex aquifolium</i>	Canada
<i>Phytophthora inflata</i>	III	IMI342898 ^{DC}	<i>Syringa</i> sp.	
<i>Phytophthora infestans</i>	IV	561 ^{PT} , P30 ^{JG}	<i>Solanum cardiophyllum</i>	Mexico
<i>Phytophthora iranica</i>	I	IMI158964 ^{DC}	<i>Solanum melongera</i>	Iran
<i>Phytophthora katusrae</i>	II	IMI360596 ^{DC}	<i>Cocos nucifera</i>	Ivory Coast
<i>Phytophthora lateralis</i>	V	451 ^{PT} , 91/11/1-5 ^{MG}	<i>Chamaecyparis lawsoniana</i>	Oregon
<i>Phytophthora medii</i>	II	IMI129185 ^{DC}	<i>Hevea brasiliensis</i>	India
<i>Phytophthora megasperma</i>	V	309 ^{PT} , 336 ^{PH}	<i>Pseudotsuga menziesii</i>	Washington
		437 ^{PT} , IMI133317	<i>Malus sylvestris</i>	Australia
<i>Phytophthora megakarya</i>	II	327 ^{PT} , P132 ^{CB}	<i>Theobroma cacao</i>	Nigeria
		328 ^{PT} , P184 ^{CB}	<i>Theobroma cacao</i>	Cameroon
<i>Phytophthora melonis</i>	VI	IMI325917 ^{DC}	<i>Cucumis</i> sp.	China
<i>Phytophthora mirabilis</i>	IV	340 ^{PT} , ATCC 64070, P3007 ^{MC}	<i>Mirabilis jalapa</i>	Mexico
<i>Phytophthora nemorosa</i>	IV	482 ^{PT} , P-13 ^{EH} Type	<i>Lithocarpus densiflorus</i>	California
<i>Phytophthora nicotianae</i>	II	360 ^{PT}	<i>Solanum tuberosum</i>	Delaware
<i>Phytophthora parasitica</i>	II	332 ^{PT} , P1751 ^{MC}	<i>Nicotiana tabacum</i>	Australia
		334 ^{PT} , P3118 ^{MC}	<i>Lycopersicon esculentum</i>	Australia

(Continued on next page)

^a Waterhouse morphological group (58).

^b CB = Clive Brasier, DC = DNA supplied by David Cooke, MC = Michael Coffey, KD = Ken Deahl, PH = Phil Hamm (E. Hansen), DJM = Dave Mitchell, DS = Dave Shaw, PT = Paul Tooley, UCR = University of California at Riverside, SW = Sabine Werres, WW = Wayne Wilcox, DR = Dave Rizzo, CDFA = Cheryl Blomquist, California Department of Food and Agriculture, PC = Plant Clinic identification by Paul Reeser, and JG = J. Galindo.

^c Species groupings of Brasier et al. (6).

Results from real-time PCR based on a DNA dilution series showed that amplification with the *P. ramorum* primers and probe occurred down to 1 fg of template DNA, which had a Ct value of 42 (Fig. 2A). A standard curve was calculated based on three replicate serial dilutions of DNA extracted from *P. ramorum* isolate 288 and demonstrated the linearity in response of the assay to DNA concentrations (Fig. 2B). Data for the 100-ag quantity was omitted from the standard curve analysis since detection was variable at that low level. Addition of plant DNA in amounts similar to those that would likely be added when assaying field samples slightly reduced the amplification efficiency of *P. ramorum* template amplification (slope of -4.14 compared with -3.68); the regression equation for the spiked DNA standard curve was $y = -4.14 \log(x) + 21.96$ with an r^2 value of 0.984.

***P. pseudosyringae*-specific primers and probe.** Primers FMPps1c and FMPps2c and the PpsCALOrange probe (Table 2) specifically detected all six isolates of *P. pseudosyringae* when tested at an annealing temperature of 55°C and did not amplify any of the other 45 *Phytophthora* species (including the closely related *P. nemorosa*) when tested at a concentration of 100 pg of DNA, including 25 isolates of *P. ramorum* (Table 3). Results of a DNA dilution series showed that amplification with the *P. pseudosyringae* primers and probe occurred down to 10 fg of template DNA, which had an average Ct value (based on six replications) of 39.94 (data not shown). A standard curve was calculated

based on three replicate serial dilutions of *P. pseudosyringae* isolate 471 each containing two replications, and the regression demonstrated the linearity in response of the assay to DNA concentrations (Fig. 2C). Data for the 1-fg quantity was omitted from the standard curve analysis since detection was variable at that low level.

Testing primers and probes with plant DNA. Both the *P. ramorum* and *P. pseudosyringae* primers and probes were tested with DNA of the following plant species using an annealing temperature of 55°C and no amplification was observed: *Rhododendron* sp. (cv. Cunningham's White), *Glycine max* cv. Williams, *Solanum demissum*, *S. cardiophyllum*, *S. tuberosum* cv. Russet Burbank, *Lycopersicon esculentum*, coast live oak (*Q. agrifolia*), laurel oak (*Q. laurifolia*), *Kalmia latifolia* cv. Olympic Wedding, California bay laurel (*U. californica*), *Pieris japonica*, highbush blueberry (*Vaccinium corymbosum*), tan oak (*Lithocarpus densiflorus*), *Citrus* sp., *Zauschneria californica*, *Fragaria × ananassa*, and *Juniperus* sp.

Sensitivity of detection of real-time PCR assay with infected tissue. We performed a dilution series from rhododendron leaf disks artificially inoculated with *P. ramorum* to determine the approximate limits of pathogen detection in infected tissue (Table 4). Even at dilutions of 10^{-6} , pathogen detection was observed, albeit with a Ct value of 55.34. The amount of DNA at each serial dilution of the infected plant extract was estimated using the

TABLE 1. (Continued from preceding page)

Species	Group ^a	Isolate number ^b	Host	Origin
<i>Phytophthora palmivora</i>	II	329 ^{PT} , P131 ^{CB}	<i>Theobroma cacao</i>	Nigeria
<i>Phytophthora phaseoli</i>	IV	352 ^{PT} , ATCC 60171, CBS556.88 373 ^{PT}	<i>Phaseolus lunatus</i> <i>Phaseolus lunatus</i>	Unknown Delaware
<i>Phytophthora porri</i>	III	CBS782.97 ^{DC}	<i>Brassica chinensis</i>	The Netherlands
<i>Phytophthora primulae</i>	III	CBS620.97 ^{DC}	<i>Primula acaulis</i>	Germany
<i>Phytophthora pseudosyringae</i>	IV	470 ^{PT} , P193907A ^{CDFA} 471 ^{PT} 472 ^{PT} 473 ^{PT} 484 ^{PT} , PSEU16 ^{TJ} , NFV-BU97-15 485 ^{PT} , P96 ^{EH}	<i>Manzanita</i> sp. <i>Umbellularia californica</i> <i>Umbellularia californica</i> <i>Umbellularia californica</i> <i>Fagus sylvatica</i> <i>Umbellularia californica</i>	Royal Oaks, CA Napa, CA Calistoga, CA Yountville, CA Germany Contra Costa Co., CA
<i>Phytophthora pseudotsugae</i>	I	308 ^{PT} , H270 ^{PH}	<i>Pseudotsugae menziesii</i>	Oregon
<i>Phytophthora quercina</i>	V	IMI340618 ^{DC}	<i>Quercus robur</i>	Germany
<i>Phytophthora ramorum</i>	IV	Prn-1 ^{PT} , PD93/844 ^{sw} Prn-2 ^{PT} , PD94/844 ^{sw} Prn-3 ^{PT} , PD98/8/6743 ^{sw} Prn-4 ^{PT} , PD98/8/6285 ^{sw} Prn-5 ^{PT} , PD98/8/2627 ^{sw} Prn-6 ^{PT} , PD98/8/5233 ^{sw} Prg-1 ^{PT} , BBA 69082 ^{sw} Prg-2 ^{PT} , BBA 9/95 ^{sw} , CBS101553 (Type) Prg-3 ^{PT} , BBA 14/98-a ^{sw} Prg-4 ^{PT} , BBA 12/98 ^{sw} Prg-5 ^{PT} , BBA 13/99-1 ^{sw} Prg-6 ^{PT} , BBA 16/99 ^{sw} Prg-7 ^{PT} , BBA 9/3 ^{sw} Prg-8 ^{PT} , BBA 104 ^{sw} 288 ^{MG} 73101 ^{CDFA} 044519 ^{CDFA} 044522 ^{CDFA} P072648 ^{CDFA} 201C ^{DR} 0-217, Pr-52 ^{DR} Coen ^{MG} 0-13, Pr-5 ^{DR} 0-16, Pr-6 ^{DR} C ^{MG}	<i>Rhododendron</i> sp. <i>Rhododendron</i> sp. <i>Rhododendron</i> sp. <i>Rhododendron</i> sp. <i>Rhododendron</i> sp. <i>Viburnum</i> sp. <i>Rhododendron</i> sp. <i>Rhododendron catawbiense</i> <i>Rhododendron catawbiense</i> <i>Rhododendron catawbiense</i> <i>Rhododendron catawbiense</i> <i>Viburnum bodnantense</i> Water Water <i>Rhododendron</i> sp. <i>Lithocarpus densiflorus</i> <i>Umbellularia californica</i> <i>Lithocarpus densiflorus</i> <i>Quercus agrifolia</i> <i>Rhododendron</i> sp. <i>Rhododendron</i> sp. <i>Rhododendron</i> sp. <i>Lithocarpus densiflorus</i> <i>Quercus agrifolia</i> <i>Umbellularia californica</i>	Netherlands Netherlands Netherlands Netherlands Netherlands Netherlands Germany Germany Germany Germany Germany Germany Germany California California California California California California California California California California California The Netherlands
<i>Phytophthora richardiae</i>	VI	ATCC 46538 ^{DC}	<i>Zantedeschia</i> sp. root	The Netherlands
<i>Phytophthora sojae</i>	V	312 ^{PT} , ATCC 48068	<i>Glycine max</i>	Wisconsin
<i>Phytophthora syringae</i>	III	442 ^{PT} , P1023 ^{CB} , IMI296829 469 ^{PT}	<i>Rubus idaeus</i> <i>Kalmia latifolia</i>	Scotland Oregon
<i>Phytophthora tentaculata</i>	I	CBS552.96 ^{DC}	<i>Chrysanthemum leucanth</i>	Germany
<i>Phytophthora</i> sp. "O" group ^c		P246 ^{DC} , IMI389751	<i>Salix</i> roots	U.K.
<i>P. taxon Raspberry</i> ^c		P896 ^{DC} , IMI389744	Soil	Tasmania

standard dilution series curve (Fig. 2) with the 10⁻⁵ dilution extrapolated to have 1.7 fg of *P. ramorum* DNA.

Use of two-way multiplex real-time PCR assay with field samples from California. Samples from naturally infected plant hosts in California received from the California Department of Food and Agriculture were evaluated with the *P. ramorum*, *P. pseudosyringae*, and plant primers (Table 5). We performed a two-way multiplex real-time PCR using *P. ramorum* and plant primers and probes. For samples negative for *P. ramorum*, we then performed a second real-time PCR using the *P. pseudosyringae* primers and probe. Results for all 53 samples showed good agreement between the real-time PCR and the results of prior analysis (Table 5). All 14 samples previously determined to be infected with *P. ramorum* were correctly identified with the real-time assay, as were all six of the samples infected with *P. pseudosyringae*. Cross reactivity between these two species or with several other *Phytophthora* spp. colonizing the tissue was not observed. Importantly, no examples of false positives were obtained. Use of plant primers and probe allowed confirmation that amplifiable DNA was present in all samples and was of high quality and did not contain PCR inhibitors that would prevent amplification and result in false negatives.

Three-way multiplex real-time PCR assay. Experiments were conducted using California bay laurel (*U. californica*) artificially infected with *P. ramorum*, *P. pseudosyringae*, or both pathogens using their respective primers and probes and plant primers and probes in three-way multiplex reactions. Initial studies were performed to determine optimum concentrations of dNTPs, magnesium, and primers/probes and optimum probe–fluorochrome combinations to prevent competitive interference between the three components in the multiplex reactions (data not shown). Two multiplex experiments were performed at an annealing temperature of 55°C, with two replications each. Ct values (Table 6) revealed specificity for each pathogen or for plants with each respective primer–probe combination. For the *P. ramorum* primer–probe combination, amplification from samples containing DNA of both pathogens had the same Ct (Table 6) and amplification curve (Fig. 3) to that obtained with *P. ramorum* alone. For the *P. pseudosyringae* primer–probe combination, amplification from samples containing both pathogens not only had a reduced Ct (Table 6), but the amplification curve was substantially reduced compared with that containing *P. pseudosyringae* alone (Fig. 3). Use of the plant primer–probe combination in multiplex PCR resulted in similar levels of amplification with individual pathogen samples as well as the combined sample (Fig. 3).

DISCUSSION

We have described a real-time PCR protocol based on mitochondrial gene regions that offers advantages over conventional PCR procedures and will provide a useful and rapid tool in nation-

wide efforts to detect the sudden oak death pathogen, *P. ramorum*. The need for such a test, which combines ease of use along with the specificity of conventional PCR and DNA hybridization (due to the inclusion of a specific TaqMan probe sequence), is especially pressing in light of the recent spread of the pathogen to the Eastern United States via shipments of nursery stock (51). The PCR method we describe can differentiate *P. ramorum* from other *Phytophthora* spp., some of which can cause similar looking lesions on the same hosts as *P. ramorum*. Using a multiplex format, additional *Phytophthora* species could be added to the assay as well. The described method uses mitochondrial gene regions rather than nuclear regions for detection, and thus offers the advantage of targeting a different region of the pathogen genome than in other tests. Several other real-time PCR assays for *P. ramorum* have targeted nuclear genes such as the ITS regions (2,13,23,27,61) and β -tubulin and elicitor genes (2). When used in combination, assays based on different genomic regions are more

Fig. 1. Real-time amplification profiles for A, *Phytophthora ramorum* and B, *P. pseudosyringae* using primers and probes described in Table 2.

TABLE 2. Polymerase chain reaction primer and fluorescent probe sequences used to develop species-specific assays for *Phytophthora ramorum* and *P. pseudosyringae*

Target	Primer/probe	Sequence (5' to 3')	Length	T _m ^a	%GC ^b
<i>P. ramorum</i>	FMPPr-1a	GTATTTAAAAATCATAGGTGTAATTTG	26	50.0	23.1
<i>P. ramorum</i>	FMPPr-7	TGGTTTTTTTAAATTATATTATCAATG	27	51.9	14.8
<i>P. ramorum</i>	PrFAM probe	6-FAM d(CAGATATTAACAAATATATATAAAATCAAACAA) BHQ-1 ^c	35	56.2	14.3
Plant	FMPI-2b	GCGTGGACCTGGAATGACTA	20	57.2	55.0
Plant	FMPI-3b	AGGTTGTATTAAAGTTTCGATCG	23	53.5	34.8
Plant	Plant CALOrange probe	CAL Orange d(CTTTTATTATCACTTCCGGTACTGGCAGG) BHQ-1	29	64.5	44.8
<i>P. pseudosyringae</i>	FMPps1c	AGTTTCATTAGAAGATTATTTAC	23	52.1	21.7
<i>P. pseudosyringae</i>	FMPps2c	AAAATTGTTTGATTTTATTAAGTATC	26	52.0	15.4
<i>P. pseudosyringae</i>	PpsCALOrange probe	CAL Orange d(TTAATAAAAAAATTATGATATTTAACTAATTGGT) BHQ-1	35	56.3	11.4

^a Melting temperature, T_m, was calculated at 50 nM primer and 50 nM salt using the program Primer Express (Applied Biosystems).

^b Percentage of guanine and cytosine.

^c TaqMan probes were labeled at the 5' end with either the fluorescent reporter dye 6-carboxy-fluorescein (FAM) or CAL Fluor Orange (CAL Orange) and labeled at the 3' end with the black hole quencher dye (BHQ, Biosearch Technologies, Novato, CA).

powerful and reliable than either test used alone, particularly in cases where one test may result in faint positive reactions and the pathogen cannot be cultured on selective agar medium. The fact that mitochondrial sequences are high copy also aids with the sensitivity of the assay.

However, the high AT/CG ratio and abundance of A and T in mitochondrial DNA offer a challenge to development of molecular detection methods. Methods such as increasing the ratio of dATP and dTTP versus dGTP and dCTP in PCR and/or reducing extension temperatures can enhance amplification of mitochondrial A+T-rich DNAs (45,52). A possible explanation for the reduced sensitivity we observed in multiplex PCR may be the A+T-rich nature of primers and probes we designed for use with our mitochondrial target region. Our primers and probes have a G/C base composition that is far below the 50% composition considered optimum (Table 2). However, it is known that low G/C content can be compensated for by an increase in primer length (10). In spite of such potential difficulties, mitochondrial gene regions have proven useful in identification and detection studies with a number of different fungi (11,16,38,64).

The specificity of our assay was determined by evaluating 45 different *Phytophthora* species (for some species multiple isolates were examined). In contrast, the specificity of the PCR assay based on the ITS region has been tested with 20 species, some of which (*P. lateralis* and *P. cambivora*) cross reacted at certain DNA concentrations (13,23). The real-time PCR assay described

here also detected a variety of *P. ramorum* isolates, including those from Europe. U.S. and European populations have been shown to be different for several characters including mating type (5,59) and our assay is able to detect *P. ramorum* from either population. The assay also exhibited a linear response between DNA concentration and detection limit and was sensitive enough to detect *P. ramorum* when present at a concentration of 1 fg of culture extracted DNA. The presence of plant extracts in the amplification mix in the amount equal to what would be used in assays of field samples did not alter the sensitivity of the assay. In fact, DNA extractions from infected leaves from a *Rhododendron* sp. could be diluted to 10^{-5} and the pathogen could still be detected. This marker system was initially developed for conventional nested PCR with the first round amplification done using a genus-specific primer pair followed by nested amplification with the species-specific primer pair (39). While it has not been experimentally verified, conducting conventional PCR with the genus-specific primers followed by the described nested real-time PCR

TABLE 3. Cycle threshold (Ct) values for 25 isolates of *Phytophthora ramorum*, *P. pseudosyringae*, and other *Phytophthora* species subjected to real-time polymerase chain reaction (PCR) analysis

	Ct value ^a	
	<i>P. ramorum</i> primers and probe	<i>P. pseudosyringae</i> primers and probe
<i>P. ramorum</i>		
Coen	28.91 ± 0.44	>60 ± 0 ^b
201C	26.40 ± 0.13	>60 ± 0
0-13	24.84 ± 0.46	>60 ± 0
0-16	26.83 ± 0.56	>60 ± 0
0-217	25.23 ± 0.11	>60 ± 0
288	28.81 ± 0.05	>60 ± 0
C	27.40 ± 0.75	>60 ± 0
73101	25.41 ± 0.55	>60 ± 0
044519	25.26 ± 0.13	>60 ± 0
044522	25.28 ± 0.44	>60 ± 0
Prn-1	25.66 ± 0.15	>60 ± 0
Prn-2	28.45 ± 0.69	>60 ± 0
Prn-3	28.87 ± 0.14	>60 ± 0
Prn-4	27.25 ± 0.01	>60 ± 0
Prn-5	26.73 ± 1.12	>60 ± 0
Prn-6	26.68 ± 0.18	>60 ± 0
Prg-1	26.88 ± 0.21	>60 ± 0
Prg-2	22.56 ± 0.11	>60 ± 0
Prg-3	24.86 ± 0.14	>60 ± 0
Prg-4	27.07 ± 0.18	>60 ± 0
Prg-5	27.49 ± 0.27	>60 ± 0
Prg-6	25.02 ± 0.15	>60 ± 0
Prg-7	28.53 ± 0.42	>60 ± 0
Prg-8	24.37 ± 0.52	>60 ± 0
P72648	25.66 ± 0.76	>60 ± 0
<i>P. pseudosyringae</i>		
470	>60 ± 0	25.41 ± 0.03
471	>60 ± 0	25.01 ± 0.40
472	>60 ± 0	24.52 ± 0.64
473	>60 ± 0	24.11 ± 0.06
484	>60 ± 0	27.74 ± 0.33
485	>60 ± 0	24.93 ± 0.25
Other <i>Phytophthora</i> species ^c		
Negative control	>60 ± 0	>60 ± 0

^a Data are mean values of two replicated experiments ± standard error.
^b No fluorescence was detected at 60 cycles of PCR amplification when tested at a concentration of 100 pg of DNA.
^c Other species listed in Table 1.

Fig. 2. A, Real-time polymerase chain reaction amplification profile for representative dilution series of DNA extracted from *Phytophthora ramorum* isolate 288. B, Standard curve of cycle threshold (Ct) values calculated from serial dilutions of DNA from *P. ramorum* isolate 288 with standard error bars indicated. C, Standard curve of Ct values calculated from serial dilutions of DNA from *P. pseudosyringae* isolate 471 with standard error bars indicated.

procedure would be expected to enhance the sensitivity of pathogen detection.

Hayden et al. (23) reported detection of *P. ramorum* down to 12 fg of DNA in an ITS-based PCR assay using SYBR green

TABLE 4. Amount of DNA estimated to be present in dilutions of DNA extracted from *Rhododendron* sp. (cv. Cunningham's White) leaf disks infected with *Phytophthora ramorum*

Dilution from Bio101 kit ^a	Ct average $\pm$ SE ^b	Amount of DNA calculated from standard curve
1:10	27.75 $\pm$ 0.32	20.9 pg
1:100	32.06 $\pm$ 0.53	1.4 pg
1:1000	35.37 $\pm$ 0.62	177 fg
1:10,000	39.57 $\pm$ 0.33	13 fg
1:100,000	42.81 $\pm$ 0.58	1.7 fg
1:1,000,000	55.34 $\pm$ 2.95	ND ^c

^a DNA was extracted from two 6-mm-diameter leaf disks using a Qbiogene Fast DNA Extraction Kit (Qbiogene, Inc., Carlsbad, CA) according to manufacturer's instructions.

^b Cycle threshold (Ct) values are means of six observations, plus or minus the standard error. Three separate extractions were performed (each using two 6-mm-diameter leaf disks), and two replicate real-time polymerase chain reaction experiments were conducted, each containing sample from all three extractions diluted as indicated ($n = 6$).

^c ND = not determined because out of range of the standard curve.

detection but several other *Phytophthora* species cross-reacted in the assay at DNA template concentrations above 0.7 ng. SYBR green binds indiscriminately to double-stranded DNA, so false positives caused by detection of primer-dimers and nonspecific amplification are possible (49). Vandemark and Barker (56) reported a detection limit of 1 pg of DNA for *P. medicaginis* using a fluorescent real-time PCR primer-probe set based on a

TABLE 6. Cycle threshold (Ct) values for multiplex experiments with California bay laurel (*Umbellularia californica*) artificially infected with *Phytophthora ramorum*, *P. pseudosyringae*, or both pathogens using primers and probes specific for *P. ramorum*, *P. pseudosyringae*, and plant DNA^a

Sample	<i>P. ramorum</i> primers and probe	<i>P. pseudosyringae</i> primers and probe	Plant primers and probe
<i>P. ramorum</i> 0-217	28.6	>60 ^b	30.6
<i>P. pseudosyringae</i> 470	>60	27.5	32.3
0-217 plus 470	28.5	34.2	29.1
Negative control	>60	>60	>60
MSD ^c	0.8	5.0	1.2

^a Data are means of four observations (two experiments with two replications each).

^b No fluorescence was detected at 60 cycles of PCR amplification.

^c Minimum significant difference, K ratio = 100 for Waller-Duncan K ratio t test for Ct value.

TABLE 5. Real-time polymerase chain reaction (PCR) results for symptomatic plant samples collected from the field in California and processed by the California Department of Food and Agriculture to determine which *Phytophthora* spp. were present

Host species	Pathogen identification ^a	Real-time PCR result (Ct value) ^b	
		<i>P. ramorum</i>	<i>P. pseudosyringae</i>
<i>Acer macrophyllum</i> (6 samples)	None detected	>60 ^c	>60
<i>Aesculus californica</i> (3 samples)	None detected	>60	>60
<i>Arbutus menziesii</i> (2 samples)	None detected	>60	>60
<i>Heteromeles arbutifolia</i> (2 samples)	None detected	>60	>60
<i>Pseudotsuga menziesii</i>	None detected	>60	>60
<i>Rhamnus californica</i>	<i>Phytophthora</i> sp.	>60	>60
<i>Rhododendron</i> sp.	<i>P. ramorum</i>	34	>60
<i>Rhododendron</i> sp.	<i>P. pseudosyringae</i>	>60	30
<i>Rhododendron</i> sp.	<i>Phytophthora</i> sp.	>60	>60
<i>Rhododendron</i> sp.	<i>Phytophthora</i> sp.	>60	>60
<i>Rhododendron</i> sp. (2 samples)	<i>P. syringae</i>	>60	>60
<i>Rhododendron</i> sp. (2 samples)	None detected	>60	>60
<i>Salal</i> sp.	None detected	>60	>60
<i>Sambucus</i> sp.	None detected	>60	>60
<i>Sequoia sempervirens</i> (2 samples)	None detected	>60	>60
<i>Umbellularia californica</i> (8 samples)	<i>P. nemorosa</i>	>60	>60
<i>Umbellularia californica</i>	<i>P. pseudosyringae</i>	>60	30
<i>Umbellularia californica</i>	<i>P. pseudosyringae</i>	>60	34
<i>Umbellularia californica</i>	<i>P. pseudosyringae</i>	>60	37
<i>Umbellularia californica</i>	<i>P. pseudosyringae</i>	>60	32
<i>Umbellularia californica</i>	<i>P. pseudosyringae</i>	>60	39
<i>Umbellularia californica</i>	<i>P. ramorum</i>	38	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	35	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	41	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	41	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	40	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	44	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	39	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	32	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	35	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	38	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	40	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	37	>60
<i>Umbellularia californica</i>	<i>P. ramorum</i>	33	>60
<i>Umbellularia californica</i> (4 samples)	None detected	>60	>60

^a Plant samples from the field were the same as discussed previously (39). They were processed at the California Department of Food and Agriculture by plating on selective medium and confirming species identification based on morphological criteria and/or amplification of DNA extracted from infected tissue with the *P. ramorum*-specific internal transcribed spacer primers. These same samples were evaluated in a prior publication with the *Phytophthora* genus-specific, *P. ramorum*, *P. nemorosa*, and *P. pseudosyringae* species-specific primer pairs (39).

^b Real-time PCR was performed following 1:10 dilution of DNA extract for multiplex amplifications using the plant and the indicated species-specific primers and probe. Results using plant primers and probe were positive for all samples, with cycle threshold (Ct) values ranging from 23 to 34.

^c No fluorescence was detected at 60 cycles of PCR amplification.

sequence characterized DNA marker. Boehm et al. (4) reported a linear standard curve for detection of *P. infestans* using real-time PCR that ranged from 10^{-6} to 1 µg of template DNA per ml. This would place the lower limit of detection in the femtogram range similar to the results obtained with our real-time PCR assay.

Multiplex PCR allows for increased sample throughput and lower operating costs since multiple pathogens can be detected within the same plant extract by using different primer-probe combinations in the same reaction. Multiplex real-time PCR assays have been used previously for detecting both host and pathogen in the same reaction (25,62), and conventional (non-real-time) multiplex PCR was used to detect *P. lateralis* in Port-Orford-cedar (61) and multiple fungal pathogens of wheat (17). We evaluated a real-time duplex assay with markers for *P. ramorum* and the plant using infected plant samples from the greenhouse and field samples from California and found a high correlation between the results of the real-time PCR assay and those of culturing and other detection methods. Perhaps due to the presence of PCR inhibitors in the samples with the extraction procedure that was used, a 10-fold dilution of field sample DNA was necessary to obtain consistent amplification. Multiplexing ampli-

fication had a limited effect on the sensitivity of detection by the *P. ramorum* markers.

In an effort to simultaneously detect two pathogens causing similar foliar symptoms on some hosts, a three-way multiplex amplification was evaluated using markers for *P. ramorum*, *P. pseudosyringae*, and the plant to serve as a positive control. While multiplexing had no effect on the sensitivity of the *P. ramorum* and plant markers, there was a reduction in the detection sensitivity for the *P. pseudosyringae* markers (Fig. 3). However, the Ct values obtained were sufficient to determine whether the target pathogen was present or not in the assay. It is known that PCR efficiencies may be decreased when multiple primer sets are present in a single tube. Also, there exist many variables within PCR assays that can affect the efficiency of multiplexing, including the sequence of the oligonucleotides, concentrations of primers and probes, and concentrations of other PCR components (10,15). One or more of these variables may have been responsible for the observed results.

In the future, we plan to extend the utility of this assay by developing primer-probe combinations for *P. nemorosa*, a pathogen present in California that is often isolated from material also infected with *P. ramorum*. We also plan to adapt the assay for use in other PCR machines such as the portable SmartCycler (Cepheid, Inc.) platform for more broad use by other laboratories and federal and state regulatory agencies.

ACKNOWLEDGMENTS

We thank A. Sechler for assistance with development of protocols for real-time PCR; C. Blomquist from the California Department of Agriculture for providing extracts from infected plant materials for these studies; D. Cooke for providing extracted DNA for some *Phytophthora* spp.; and M. Benson for helpful editorial suggestions. We acknowledge the U.S. Forest Service, Pacific Southwest Research Station, for providing grant support for some of this work from the Sudden Oak Death Research Program.

LITERATURE CITED

1. Bates, J. A., Taylor, E. J. A., Kenyon, D. M., and Thomas, J. E. 2001. The application of real-time PCR to the identification, detection and quantification of *Pyrenophora* species in barley seed. *Mol. Plant Pathol.* 2:49-57.
2. Bilodeau, G., Levesque, C. A., de Cock, A. W. A. M., Duchaine, C., Kristjansson, G., and Hamelin, R. C. 2005. Molecular detection of *Phytophthora ramorum* by real-time PCR using TaqMan, SYBR green and molecular beacons with three genes. Proceedings of the Sudden Oak Death Second Science Symposium: The State of Our Knowledge. M. Haverty, P. Shea, and S. J. Frankel, eds. U.S. Dep. Agric. For. Serv., Gen. Tech. Rep., Pacific Southwest Research Station, Albany, CA. (In press)
3. Blum, B. H., Cousin, M. A., and Woloshuk, C. P. 2004. Multiplex real-time PCR detection of fumonisin-producing and trichothecene-producing groups of *Fusarium* species. *J. Food Prot.* 67:536-543.
4. Boehm, J., Hahn, A., Schubert, R., Bahnweg, G., Adler, N., Nechwatal, J., Oehlmann, R., and Osswald, W. 1999. Real-time quantitative PCR: DNA determination in isolated spores of the mycorrhizal fungus *Glomus mosseae* and monitoring of *Phytophthora infestans* and *Phytophthora citricola* in their respective host plants. *J. Phytopathol.* 147:409-416.
5. Brasier, C. 2003. Sudden oak death: *Phytophthora ramorum* exhibits transatlantic differences. *Mycol. Res.* 107:258-259.
6. Brasier, C. M., Cooke, D. E., Duncan, J. M., and Hansen, E. M. 2003. Multiple new phenotypic taxa from trees and riparian ecosystems in *Phytophthora gonapodyides*-*P. megasperma* ITS clade 6, which tend to be high-temperature tolerant and either inbreeding or sterile. *Mycol. Res.* 107:277-290.
7. California Department of Food and Agriculture. 2003. CDFA Plant Quarantine Manual. Pages 224.1-224.5 in: Section 301.92 *Phytophthora ramorum*. CDFA Pest Exclusion Branch, Sacramento, CA.
8. Canadian Food Inspection Agency. 2003. Phytosanitary Requirements to Prevent the Entry of *Phytophthora ramorum* Associated with Sudden Oak Death into Canada. Canadian Food Inspection Agency Directive D-01-01.
9. Caten, C. C., and Jinks, J. L. 1968. Spontaneous variability of single isolates of *Phytophthora infestans*. I. Cultural variation. *Can. J. Bot.* 46:329-348.

Fig. 3. Amplification profiles from multiplex real-time polymerase chain reaction analysis of leaf samples of California bay laurel (*Umbellularia californica*) artificially infected with *Phytophthora ramorum*, *P. pseudosyringae*, or both pathogens. Multiple experiments were performed; these amplification profiles represent results of a single run. The dye used for the *P. ramorum* probe was A, FAM, that for the *P. pseudosyringae* probe was B, CAL Orange, and that for the plant probe was C, TAMRA. Table 6 provides cycle threshold values associated with multiplex analysis.

10. Chamberlain, J. S., and Chamberlain, J. R. 1994. Optimization of multiplex PCRs. Pages 38-46 in: *The Polymerase Chain Reaction*. K. B. Mullis, F. Ferre, and R. A. Gibbs, eds. Birkhauser, Boston, MA.
11. Cohen, S., Allasia, V., Venard, P., Notter, S., Verniere, C., and Panabieres, F. 2003. Intraspecific variation in *Phytophthora citrophthora* from citrus trees in Eastern Corsica. *Eur. J. Plant Pathol.* 109:791-805.
12. Cooperative Agriculture Pest Survey program. 2002. Sudden oak death 2002/2003 pilot national survey. 04 March 2003. Center for Environmental and Regulatory Systems (CERIS), Purdue University, West Lafayette, IN. Published online.
13. Davidson, J. M., Werres, S., Garbelotto, M., Hansen, E. M., and Rizzo, D. M. 2003. Sudden oak death and associated diseases caused by *Phytophthora ramorum*. Online. *Plant Health Progress* doi:10.1094/PHP-2003-0707-01-DG.
14. De Merlier, D., Chandelier, A., and Cavelier, M. 2003. First report of *Phytophthora ramorum* on *Viburnum bodnantense* in Belgium. *Plant Dis.* 87:203.
15. Exner, M. M., and Lewinski, M. A. 2002. Sensitivity of multiplex real-time PCR reactions, using the LightCycler and the ABI prism 7700 sequence detection system, is dependent on the concentration of the DNA polymerase. *Mol. Cell. Probes* 16:351-357.
16. Ferreira, M. A. S. V., Tooley, P. W., Hatziloukas, E., Castro, C., and Schaad, N. W. 1996. Isolation of a species-specific mitochondrial DNA sequence for identification of *Tilletia indica*, the Karnal bunt of wheat fungus. *Appl. Environ. Microbiol.* 62:87-93.
17. Fraaije, B. A., Lovell, D. J., Coelho, J. M., Baldwin, S., and Hollomon, D. W. 2001. PCR-based assays to assess wheat varietal resistance to blotch (*Septoria tritici* and *Stagonospora nodorum*) and rust (*Puccinia striiformis* and *Puccinia recondita*) diseases. *Eur. J. Plant Pathol.* 107:905-917.
18. Frederick, R. D., Snyder, C. L., Peterson, G. L., and Bonde, M. R. 2002. Polymerase chain reaction assays for the detection and discrimination of the soybean rust pathogens *Phakopsora pachyrhizi* and *P. meibomia*. *Phytopathology* 92:217-227.
19. Frederick, R. D., Snyder, K. E., Tooley, P. W., Berthier-Schaad, Y., Peterson, G. L., Bonde, M. R., Schaad, N. W., and Knorr, D. A. 2000. Identification and differentiation of *Tilletia indica* and *T. walkeri* using the polymerase chain reaction. *Phytopathology* 90:951-960.
20. Goodwin, S. B., Drenth, A., and Fry, W. E. 1992. Cloning and genetic analysis of two highly polymorphic, moderately repetitive nuclear DNAs from *Phytophthora infestans*. *Curr. Genet.* 22:107-115.
21. Guillemette, T., Iacomini-Vasilescu, B., and Simoneau, P. 2004. Conventional and real-time PCR-based assay for detecting pathogenic *Alternaria brassicae* in cruciferous seed. *Plant Dis.* 88:490-496.
22. Hansen, E. M., Reeser, P. W., Sutton, W., Winton, L. M., and Osterbauer, N. 2003. First report of A1 mating type of *Phytophthora ramorum* in North America. *Plant Dis.* 87:1267.
23. Hayden, K. J., Rizzo, D., Tse, J., and Garbelotto, M. 2004. Detection and quantification of *Phytophthora ramorum* from California forests using a real-time polymerase chain reaction assay. *Phytopathology* 94:1075-1083.
24. Henricot, B., and Prior, C. 2004. *Phytophthora ramorum*, the cause of sudden oak death or ramorum leaf blight and dieback. *Mycologist* 18:151-156.
25. Hietala, A. M., Eikenes, M., Kvaalen, H., Solheim, H., and Fossdal, C. G. 2003. Multiplex real-time PCR for monitoring *Heterobasidion annosum* colonization in Norway spruce clones that differ in disease resistance. *Appl. Environ. Microbiol.* 69:4413-4420.
26. Holland, P. M., Abramson, R. D., Watson, R., and Gelfand, D. H. 1991. Detection of specific polymerase chain reaction product by utilizing the 5' to 3' exonuclease activity of *Thermus aquaticus* DNA polymerase. *Proc. Natl. Acad. Sci. USA* 88:7276-7280.
27. Hughes, K., Griffin, R., Tomlinson, J., Boonham, N., Barton, V., Giltrap, P., Hobden, E., Walker, L., Humphries, G., Barnes, A., Beales, P., Inman, A., and Lane, C. 2005. Comparative evaluation of real-time PCR (TaqMan) with isolation for diagnosis of *Phytophthora ramorum*. Poster no. 43 in: *Proceedings of the Sudden Oak Death Second Science Symposium: The State of Our Knowledge*. M. Haverly, P. Shea, and S. J. Frankel, eds. U.S. Dep. Agric. For. Serv., Gen. Tech. Rep., Pacific Southwest Research Station, Albany, CA. (In press)
28. Inman, A. J., Townend, V. C., Barnes, A. V., Lane, C. R., Hughes, K. J. D., Griffin, R. L., and Eales, S. J. 2003. First report of Ramorum dieback (*Phytophthora ramorum*) on *Pieris* in England. *New Dis. Rep.* 7. Published online by the British Society for Plant Pathology.
29. Ippolito, A., Schena, L., Nigro, F., Ligorio, V.-S., and Yaseen, T. 2004. Real-time detection of *Phytophthora nicotianae* and *P. citrophthora* in citrus roots and soil. *Eur. J. Plant Pathol.* 110:833-843.
30. Isayenkova, S., Fester, T., and Hause, B. 2004. Rapid determination of fungal colonization and arbuscule formation in roots of *Medicago truncatula* using real-time (RT) PCR. *J. Plant Physiol.* 161:1379-1383.
31. Jones, J. 2004. APHIS List of plants regulated and associated with *Phytophthora ramorum*. Published online by USDA-APHIS.
32. Kong, P., Hong, C. X., Tooley, P. W., Ivors, K., Garbelotto, M., and Richardson, P. A. 2004. Rapid identification of *Phytophthora ramorum* using PCR-SSCP analysis of ribosomal DNA ITS-1. *Lett. Appl. Microbiol.* 38:433-439.
33. Kroon, L. P. N. M., Verstappen, E. C. P., Kox, L. F. F., Flier, W. G., and Bonants, P. J. M. 2004. A rapid diagnostic test to distinguish between American and European populations of *Phytophthora ramorum*. *Phytopathology* 94:613-620.
34. Mackay, I. M., Arden, K. E., and Nitsche, A. 2002. Real-time PCR in virology. *Nucleic Acids Res.* 30:1292-1305.
35. Malorny, B., Paccassoni, E., Fach, P., Bunge, C., Martin, A., and Helmuth, R. 2004. Diagnostic real-time PCR for detection of *Salmonella* in food. *Appl. Environ. Microbiol.* 70:7046-7052.
36. Martin, F. N., and Tooley, P. W. 2003. Phylogenetic relationships among *Phytophthora* species inferred from sequence analysis of mitochondrially encoded cytochrome oxidase I and II genes. *Mycologia* 95:269-284.
37. Martin, F. N., and Tooley, P. W. 2003. Phylogenetic relationships of *Phytophthora ramorum*, *P. nemorosa*, and *P. pseudosyringae*, three species recovered from areas in California with sudden oak death. *Mycol. Res.* 12:1379-1391.
38. Martin, F. N., and Tooley, P. W. 2004. Identification of *Phytophthora* isolates to species level using restriction fragment length polymorphism analysis of a polymerase chain reaction-amplified region of mitochondrial DNA. *Phytopathology* 94:983-991.
39. Martin, F. N., Tooley, P. W., and Blomquist, C. 2004. Molecular detection of *Phytophthora ramorum*, the causal agent of sudden oak death in California, and two additional species commonly recovered from diseased plant material. *Phytopathology* 94:621-631.
40. Moralejo, E., and Werres, S. 2002. First report of *Phytophthora ramorum* on *Rhododendron* sp. in Spain. *Plant Dis.* 86:1052.
41. Parke, J. L., Linderman, R. G., Osterbauer, N. K., and Griesbach, J. A. 2004. Detection of *Phytophthora ramorum* blight in Oregon nurseries and completion of Koch's postulates on *Pieris*, *Rhododendron*, *Viburnum*, and *Camellia*. *Plant Dis.* 88:87.
42. *Phytophthora ramorum*. Code of Federal Regulations. 2003 ed. Title 7, Pt. 301.92. Pages 142-148. U.S. Gov. Printing Office, Washington, DC.
43. Rintila, T., Kassinen, A., Malinen, E., Krogius, L., and Palva, A. 2004. Development of an extensive set of 16S rDNA-targeted primers for quantification of pathogenic and indigenous bacteria in faecal samples by real-time PCR. *J. Appl. Microbiol.* 97:1166-1177.
44. Rizzo, D. M., Garbelotto, M., Davidson, J. M., Slaughter, G. W., and Koike, S. T. 2002. *Phytophthora ramorum* as the cause of extensive mortality of *Quercus* spp. and *Lithocarpus densiflorus* in California. *Plant Dis.* 86:205-214.
45. Rondan-Duenas, J. C., Panzetta-Dutari, G. M., and Gardenal, C.-N. 1999. Specific requirements for PCR amplification of long mitochondrial A+T-rich DNA. *Biotechniques* 27:258-260.
46. Schaad, N. W., and Frederick, R. D. 2002. Real-time PCR and its application for rapid plant disease diagnostics. *Can. J. Plant Pathol.* 24:250-258.
47. Schena, L., Nigro, F., and Ippolito, A. 2004. Real time PCR detection and quantification of soilborne fungal pathogens: The case of *Rosellinia necatrix*, *Phytophthora nicotianae*, *P. citrophthora*, and *Verticillium dahliae*. *Phytopathol. Mediterr.* 43:273-280.
48. Schena, L., Nigro, F., Ippolito, A., and Gallitelli, D. 2004. Real-time quantitative PCR: A new technology to detect and study phytopathogenic and antagonistic fungi. *Eur. J. Plant Pathol.* 110:893-908.
49. Schnerr, H., Niessen, L., and Vogel, R. F. 2001. Real time detection of the *Trt5* gene in *Fusarium* species by LightCycler PCR using SYBR Green I for continuous fluorescence monitoring. *Int. J. Food Microbiol.* 71:53-61.
50. Smith, M. S., Yang, R. K., Knapp, C. W., Niu, Y. F., Peak, N., Hanfelt, M. M., Galland, J. C., and Graham, D. W. 2004. Quantification of tetracycline resistance genes in feedlot lagoons by real-time PCR. *Appl. Environ. Microbiol.* 70:7372-7377.
51. Stokstad, E. 2004. Nurseries may have shipped sudden oak death pathogen nationwide. *Science* 303:1959.
52. Su, X.-Z., Wu, Y., Sifri, C. D., and Wellems, T. E. 1996. Reduced extension temperatures required for PCR amplification of extremely A+T-rich DNA. *Nucleic Acids Res.* 24:1574-1575.
53. Tooley, P. W., Kyde, K. L., and Englander, L. 2004. Susceptibility of selected ericaceous ornamental host species to *Phytophthora ramorum*. *Plant Dis.* 88:993-999.
54. USDA Animal Plant Health Inspection Service. 2004. Sudden oak death: *Phytophthora ramorum*. Pest Detection and Management Programs. Published online by USDA Animal Plant Health Inspection Service.
55. van de Graaf, P., Lees, A. K., Cullen, D. W., and Duncan, J. M. 2003. Detection and quantification of *Spongopora subterranea* in soil, water and plant tissue samples using real-time PCR. *Eur. J. Plant Pathol.* 109:589-597.
56. Vandemark, G. J., and Barker, B. M. 2003. Quantifying *Phytophthora medicaginis* in susceptible and resistant alfalfa with a real-time fluorescent PCR assay. *J. Phytopathol.* 151:577-583.

57. Wang, P. H., Boo, L. M., Lin, Y. S., and Yeh, Y. 2002. Specific detection of *Pythium aphanidermatum* from hydroponic nutrient solution by booster PCR with DNA primers developed from mitochondrial DNA. *Phytoparasitica* 30:473-485.
58. Waterhouse, G. M. 1963. Key to the species of *Phytophthora* de Bary. Mycological Paper 92. Commonwealth Mycological Institute, Kew, UK.
59. Werres, S., and De Merlier, D. 2003. First detection of *Phytophthora ramorum* mating type A2 in Europe. *Plant Dis.* 87:1266.
60. Werres, S., Marwitz, R., Man In't Veld, W. A., DeCock, A. W. A. M., Bonants, P. J. M., De Weerd, M., Themann, K., Ilieva, E., and Baayen, R. P. 2001. *Phytophthora ramorum* sp. nov., a new pathogen on *Rhododendron* and *Viburnum*. *Mycol. Res.* 105:1155-1165.
61. Winton, L. M., and Hansen, E. M. 2001. Molecular diagnosis of *Phytophthora lateralis* in trees, water, and foliage baits using multiplex polymerase chain reaction. *For. Pathol.* 31:275-283.
62. Winton, L. M., Stone, J. K., Watrud, L. S., and Hansen, E. M. 2002. Simultaneous one-tube quantification of host and pathogen DNA with real-time polymerase chain reaction. *Phytopathology* 92:112-116.
63. Xu, D.-Y., Huang, H., and Wang, C.-P. 1982. Polyacrylamide gel disc electrophoresis of proteins from species of *Phytophthora*. *Acta Mycol. Sin.* 1:40-47.
64. Zeng, Q.-Y., Rasmuson-Lestander, A., and Wang, X.-R. 2004. Extensive set of mitochondrial LSR rDNA-based oligonucleotide probes for the detection of common airborne fungi. *FEMS Microbiol. Lett.* 237:79-87.