Systematic Numbering of Vegetative Compatibility Groups in the Plant Pathogenic Fungus *Fusarium oxysporum*

H. C. Kistler, C. Alabouvette, R. P. Baayen, S. Bentley, D. Brayford, A. Coddington, J. Correll, M.-J. Daboussi, K. Elias, D. Fernandez, T. R. Gordon, T. Katan, H. G. Kim, J. F. Leslie, R. D. Martyn, Q. Migheli, N. Y. Moore, K. O'Donnell, R. C. Ploetz, M. A. Rutherford, B. Summerell, C. Waalwijk, and S. Woo

First author: Plant Pathology Department, University of Florida, Gainesville 32611-0680; second author: Laboratoire de Recherches sur la Flore Pathogène du Sol, INRA, 17 rue Sully, B.V. 1540 21034 Dijon Cedex, France; third and twenty-second authors: DLO Research Institute for Plant Protection, P.O. Box 9060, 6700 GW Wageningen, the Netherlands; fourth author: CRC for Tropical Plant Pathology, Fifth Level, John Hines Building, The University of Queensland, Brisbane, Queensland 4072, Australia; fifth and twentieth authors: International Mycological Institute, Bakeham Lane, Englefield Green, Egham, Surrey TW20 9TY, United Kingdom; sixth author: School of Biological Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom; seventh author: Department of Plant Pathology, University of Arkansas, Fayetteville 72701; eighth author: Génétique Moléculaire des Champignons Phytopathogènes, Institut de Génétique et Microbiologie, Bât 400, Université Paris-Sud 91405 Orsay Cedex, France; ninth author: USDA, ARS, Systematic Botany and Mycology Laboratory, B011A, Room 304, 10300 Baltimore Avenue, Beltsville, MD 20705; tenth author: Laboratoire de Phytopathologie, ORSTOM, B.P. 5045, 34032 Montpellier Cedex 1, France; eleventh author: Department of Plant Pathology, University of California, Davis 95616; twelfth author: Department of Plant Pathology, ARO, Volcani Center, P.O. Box 6, Bet Dagan 50250, Israel; thirteenth author: Department of Agricultural Biology, Chungnam National University, Taejon 305-764, Korea; fourteenth author: Department of Plant Pathology, Throckmorton Hall, Kansas State University, Manhattan 66506-5502; fifteenth author: Department of Botany & Plant Pathology, Purdue University, West Lafayette, IN 47907-1155; sixteenth author: Dipartimento di Valorizzazione e Protezione delle Risorse Agroforestali, Università di Torino, Via Leonardo da Vinci 44, I-10095 Grugliasco (Torino), Italy; seventeenth author: Plant Protection Unit, Department of Primary Industries, Indooroopilly Sciences Centre, 80 Meiers Road, Indooroopilly, Queensland 4068 Australia; eighteenth author: National Center for Agricultural Utilization Research, USDA/ARS, Peoria, IL 61604; nineteenth author: Tropical Research and Education Center, 18905 SW 280th Street, Homestead, FL 33031-3314; twenty-first author: Royal Botanic Gardens, Mrs. Macquaries Road, Sydney, New South Wales 2000, Australia; and twenty-third author: Istituto di Patologia Vegetale, Via Università, 100, 80055 Portici (Napoli), Italy.

Accepted for publication 2 October 1997.

Fusarium oxysporum Schlechtend.:Fr. is a cosmopolitan fungal pathogen (4) responsible for wilt and cortical rot diseases of more than 100 economically important plant hosts (3). Genetic diversity within *F. oxysporum* has been categorized extensively by vegetative compatibility grouping in laboratories around the world. Detailed studies indicate that isolates belonging to the same vegetative compatibility group (VCG) typically possess very similar or identical multilocus haplotypes and belong to the same clonal lineage (19). VCGs, therefore, can be good predictors of genetic similarity, clonal lineage, or both. In this communication, we propose a standardization of the system currently used for categorizing genetic diversity within this taxon. Our primary objective is to make the system more understandable and useful by reducing ambiguity and increasing the rigor with which the current system is applied. A secondary goal of this undertaking is to promote international communication and cooperation in this endeavor.

In the mid-1980s, Puhalla (27) proposed a method by which various strains of *F. oxysporum* could be identified and classified based on VCGs. Since that time, numerous investigators have classified isolates of *F. oxysporum* in this manner and have adopted Puhalla's numerical system for classification. In this system, vegetatively compatible isolates are given a VCG code consisting of a four- (or five-) digit number with the first three digits corresponding to host specialization, or forma specialis, and the last digit(s) corresponding to individual VCGs within that forma specialis. Puhalla categorized 16 VCGs in his original publication and provided

specific numbers for these VCGs. Since Puhalla's original contribution, over 125 VCGs have been identified among plant pathogenic isolates of *F. oxysporum*. Currently described VCGs are summarized in Table 1 (T. Katan, *unpublished data*). This list will be updated periodically on the worldwide web and can be accessed by contacting H. Corby Kistler (Plant Pathology Department, University of Florida, Gainesville 32611-0680; Phone: 352/392-9241; E-mail address: hck@gnv.ifas.ufl.edu). While care has been taken to continue the system of Puhalla, problems involving duplication have arisen as VCGs have been described for newly characterized formae speciales. For example, isolates from formae speciales *basilici* (16) and *phaseoli* (35) each have been given the VCG numerical code 016-.

To improve standardization, we endorse the following system for numbering formae speciales not specifically mentioned in the original publication of Puhalla and for putative nonpathogenic isolates of *F. oxysporum*. For the first described VCG in a forma specialis, the first three digits given to that VCG should be assigned by a VCG numbering coordinator (described below) and the last digit should be designated "0." VCGs described within that forma specialis should be numbered consecutively. To prevent overproliferation of VCG categories, we recommend numbering VCGs only if several independent isolates are found to be vegetatively compatible. Single-member VCGs or VCGs that are extremely localized geographically should not be numbered unless they possess other noteworthy attributes such as unusual cultural characteristics or pathogenicity, or can be shown by molecular markers to represent distinct lineages.

As more is learned about phylogenetic relationships within *Fusarium*, some VCGs may need to be transferred to related species and, thus, removed from the current listing. Baayen and coworkers (1), for example, recently discovered that some VCGs of the carnation wilt fungus originally described as *F. oxysporum* f. sp. *dianthi*

Corresponding author: H. C. Kistler; E-mail address: hck@gnv.ifas.ufl.edu

Publication no. P-1997-1031-01O

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 1998.

TABLE 1. Vegetative compatibility groups (VCGs) of *Fusarium oxysporum* listed by forma specialis

forma specialis	Host	forma specialis code	Assigned numbers for VCG	Number of VCGs	Literature citations
<i>albedinis</i>	<i>Phoenix dactylifera</i>	017-	0170	1	32
<i>apii</i>	<i>Apium</i>	001-	0010 to 0012	3	5
<i>asparagi</i>	<i>Asparagus</i>	100-	1001 to 1008	8 + ^a	7
<i>basilici</i>	<i>Ocimum</i>	016-	0160	1	16
<i>betae</i>	<i>Beta</i>	... ^b	...	7 +	12
<i>chrysanthemi</i>	<i>Chrysanthemum</i>	005-	0050 to 0051	2	27
<i>ciceris</i>	<i>Cicer</i>	...	...	1	25
<i>conglutinans</i>	<i>Brassica</i>	010-	0101	1	20
<i>cubense</i>	<i>Musa</i>	012-	0120 to 0126, 0128 to 01220	21 ? ^c	26
<i>cucumerinum</i>	<i>Cucumis sativus</i>	018-	0180	1 +	15
<i>cyclaminis</i>	<i>Cyclamen</i>	015-	0151 to 0153	3	36
<i>dianthi</i>	<i>Dianthus</i>	002-	0020 to 0022, 0025, 0027, 0028	6	1
<i>elaeidis</i>	<i>Elaeis</i>	001-* ^d , 014-	0011 to 0015, 0140 to 0141	5	6, 10
<i>erythroxyli</i>	<i>Erythroxylum</i>	...	...	1	31
<i>gladioli</i>	<i>Gladiolus</i>	034-	0340 to 0343	4	24
<i>lactucum</i>	<i>Lactuca</i>	...	...	1	13
<i>lilii</i>	<i>Lilium</i>	...	...	4	22
<i>lupini</i>	<i>Lupinus</i>	...	...	2 ?+	28, 29
<i>lycopersici</i>	<i>Lycopersicon</i>	003-	0030 to 0033	4 +	23
<i>mathioli</i>	<i>Mathiola</i>	010-	0103	1	20
<i>medicaginis</i>	<i>Medicago</i>	004-	0040 to 0041	2	27
<i>melongenae</i>	<i>Solanum melongena</i>	017-	0170	1	15
<i>melonis</i>	<i>Cucumis melo</i>	013-	0130 to 0138	8 +	14, 17
<i>niveum</i>	<i>Citrullus</i>	008-	0080 to 0082	3	21
<i>phaseoli</i>	<i>Phaseolus</i>	016-	0161 to 0165	5	35
<i>pisi</i>	<i>Pisum</i>	007-	0070	6 ?+	2, 34
<i>radicis-lycopersici</i>	<i>Lycopersicon</i>	009-	0090 to 0098	8 +	18, 30
<i>raphani</i>	<i>Raphanus</i>	010-	0102	1	20
<i>spinaciae</i>	<i>Spinacia</i>	...	...	3	9
<i>tracheiphilum</i>	<i>Vigna</i>	006-	0060	1	27
<i>tuberosi</i>	<i>Solanum tuberosum</i>	...	...	6	33
<i>vasinfectum</i>	<i>Gossypium</i>	011-	0111 to 01110	10	8

^a + = Additional isolates are incompatible with the established VCGs.

^b ... = VCG code and numbers not assigned.

^c ? = Two or more of these VCGs may overlap.

^d * = forma specialis code previously given to forma specialis *apii* by Puhalla (27).

are actually members of *F. redolens* f. sp. *dianthi*. In cases such as this, VCG numbers for isolates no longer considered to belong to *F. oxysporum* should be removed from use without replacement. A similar four-digit code as used for VCGs in *F. oxysporum* has been proposed for VCGs and formae speciales in *F. redolens* (1). The latter species causes wilt and rot diseases in numerous host plants including pea, carnation, and spinach (11). Other VCGs currently placed in *F. oxysporum* may eventually prove to belong to *F. redolens* or other species of *Fusarium*.

As a rule, nonpathogenic isolates of *F. oxysporum* should not be given four-digit VCG codes. Since tests for vegetative compatibility are unlikely to be done among the many nonpathogenic isolates from different laboratories, we discourage the routine assignment of formal VCG numbers for these isolates to prevent an overabundance of VCG categories. In some cases, however, these isolates will be sufficiently important to warrant classification of their VCG. VCG numbers may be assigned when (i) isolates belonging to the same VCG are recovered from multiple locations, (ii) isolates belonging to the same VCG compose a large fraction (>5%) of the population at a single site (sample size > 100 independent isolates), or (iii) isolates belonging to the same VCG are of particular utility, e.g., biological control isolates. Nonpathogenic isolates vegetatively compatible with pathogenic isolates also should be given the VCG number of their pathogenic counterparts. Four-digit VCG codes for nonpathogens will be assigned by the coordinator as done for pathogenic VCGs. If the following recommendations for numbering are followed, duplications and inconsistencies in the system can be kept to a minimum.

To address the problems outlined here, and with the endorsement of the *Fusarium* Committee of the International Society of Plant Pathology, we hereby propose the creation of a "VCG numbering coordinator." This person should be contacted by investi-

gators prior to submission of new VCG numbers for publication. If requested by the investigator, communications with the VCG numbering coordinator will be kept strictly confidential prior to publication. Talma Katan (Department of Plant Pathology, ARO, Volcani Center, P.O. Box 6, Bet Dagan 50250, Israel; fax: 972-3-968-3543) has accepted our nomination to serve as the initial VCG numbering coordinator until the 6th International *Fusarium* Workshop (to be held at the International Mycological Institute from 17 to 21 August 1998), at which time a new VCG coordinator can be selected or T. Katan can be renominated to continue as coordinator.

In the coming year, representative isolates of all known VCGs of *F. oxysporum* should be deposited for storage at two or more internationally recognized fungal culture collections. Individual investigators will be contacted shortly to assemble this collection. The dual utility of such a centralized collection is that it will be able to provide reference cultures of all of the major phytopathogenic VCGs and insure long-term strain maintenance. Puhalla (27) noted that "(to) standardize the assignment of VCG, a great deal of cooperation between *Fusarium* workers will be necessary." It is our hope that the efforts begun here will promote the cooperation required to make VCG standardization possible.

LITERATURE CITED

1. Baayen, R. P., van Dreven, F., Krijger, M. C., and Waalwijk, C. 1997. Genetic diversity in *Fusarium oxysporum* f.sp. *dianthi* and *Fusarium redolens* f.sp. *dianthi*. Eur. J. Plant Pathol. 103:395-408.
2. Bodker, L., Lewis, B. G., and Coddington, A. 1993. The occurrence of a new genetic variant of *Fusarium oxysporum* f.sp. *pisi*. Plant Pathol. 42: 833-838.
3. Booth, C. 1971. The Genus *Fusarium*. Commonwealth Mycological Institute, Kew, Surrey, United Kingdom.

4. Correll, J. C. 1991. The relationship between formae speciales, races, and vegetative compatibility groups in *Fusarium oxysporum*. *Phytopathology* 81:1061-1064.
5. Correll, J. C., Klittich, C. J. R., and Leslie, J. F. 1987. Nitrate nonutilizing mutants of *Fusarium oxysporum* and their use in vegetative compatibility tests. *Phytopathology* 77:1640-1646.
6. Dossa, C., Pando-Bahuon, A., Renard, J. L., and Boisson, C. 1991. Determination of vegetative compatibility groups in African *Fusarium oxysporum* strains isolated from vascular wilt-infected oil palms. *Oleagineux* 46:145-147.
7. Elmer, W. H., and Stephens, C. T. 1989. Classification of *Fusarium oxysporum* f. sp. *asparagi* into vegetatively compatible groups. *Phytopathology* 79:88-93.
8. Fernandez, D., Assigbetse, K., Dubois, M. P., and Geiger, J. P. 1994. Molecular characterization of races and vegetative compatibility groups in *Fusarium oxysporum* f.sp. *vasinfectum*. *Appl. Environ. Microbiol.* 60: 4039-4046.
9. Fiely, M. B., Correll, J. C., and Morelock, T. E. 1995. Vegetative compatibility, pathogenicity, and virulence diversity of *Fusarium oxysporum* recovered from spinach. *Plant Dis.* 79:990-993.
10. Flood, J., Whitehead, D. S., and Cooper, R. M. 1992. Vegetative compatibility and DNA polymorphisms in *Fusarium oxysporum* f.sp. *elaeidis* and their relationship to isolate virulence and origin. *Physiol. Mol. Plant Pathol.* 41:201-215.
11. Gerlach, W., and Nirenberg, H. I. 1982. The genus *Fusarium*—A pictorial atlas. *Mitt. Biol. Bundesanst. Land- Forstwirtschaft. Berlin-Dahlem.* 209:1-406.
12. Harveson, R. M., and Rush, C. M. 1997. Genetic variation among *Fusarium oxysporum* isolates from sugar beet as determined by vegetative compatibility. *Plant Dis.* 81:85-88.
13. Hubbard, J. C., and Gerik, J. S. 1993. A new wilt disease of lettuce incited by *Fusarium oxysporum* f. sp. *lactucum* forma specialis nov. *Plant Dis.* 77:750-754.
14. Jacobson, D. J., and Gordon, T. R. 1990. Further investigations of vegetative compatibility within *Fusarium oxysporum* f.sp. *melonis*. *Can. J. Bot.* 68:1245-1248.
15. Katan, T. 1996. Vegetative-compatibility groups in populations of *Fusarium oxysporum* in Israel. (Abstr.) *Phytoparasitica* 24:139.
16. Katan, T., Gamliel, A., and Katan, J. 1996. Vegetative compatibility of *Fusarium oxysporum* from sweet basil in Israel. *Plant Pathol.* 45:656-661.
17. Katan, T., Katan, J., Gordon, T. R., and Poznaniak, D. 1994. Physiologic races and vegetative compatibility groups of *Fusarium oxysporum* f. sp. *melonis* in Israel. *Phytopathology* 84:153-157.
18. Katan, T., Zamir, D., Sarfatti, M., and Katan, J. 1991. Vegetative compatibility groups and subgroups in *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Phytopathology* 81:255-262.
19. Kistler, H. C. 1997. Genetic diversity in the plant-pathogenic fungus *Fusarium oxysporum*. *Phytopathology* 87:474-479.
20. Kistler, H. C., Momol, E. A., and Benny, U. 1991. Repetitive genomic sequences for determining relatedness among strains of *Fusarium oxysporum*. *Phytopathology* 81:331-336.
21. Larkin, R. P., Hopkins, D. L., and Martin, F. N. 1990. Vegetative compatibility within *Fusarium oxysporum* f.sp. *niveum* and its relationship to virulence, aggressiveness, and race. *Can. J. Microbiol.* 36:352-358.
22. Löffler, H. J. M., and Rumine, P. 1991. Virulence and vegetative compatibility of Dutch and Italian isolates of *Fusarium oxysporum* f.sp. *lilii*. *J. Phytopathol.* 132:12-20.
23. Marlatt, M. L., Correll, J. C., Kaufmann, P., and Cooper, P. E. 1996. Two genetically distinct populations of *Fusarium oxysporum* f. sp. *lycopersici* race 3 in the United States. *Plant Dis.* 80:1336-1342.
24. Mes, J. J., van Doorn, J., Roebroek, E. J. A., van Egmond, E., van Aartrijk, J., and Boonekamp, P. M. 1994. Restriction fragment length polymorphisms, races and vegetative compatibility groups within a worldwide collection of *Fusarium oxysporum* f.sp. *gladioli*. *Plant Pathol.* 43: 362-370.
25. Nogales-Moncada, A. M., Pérez-Artés, E., and Jiménez-Díaz, R. M. 1993. VCG diversity within *Fusarium oxysporum* f.sp. *ciceris*. (Abstr.) Page 169 in: *Abstr. Int. Congr. Plant Pathol.*, 6th. National Research Council Canada, Ottawa.
26. Ploetz, R. C. 1994. Panama disease: Return of the first banana menace. *Int. J. Pest Manage.* 40:326-336.
27. Puhalla, J. E. 1985. Classification of strains of *Fusarium oxysporum* on the basis of vegetative compatibility. *Can. J. Bot.* 63:179-183.
28. Rataj-Guranowska, M. 1992. *Fusarium oxysporum* f.sp. *lupini* in Wielkopolska-genetic interrelationships. Pages 133-139 in: *IHAR Radzikow* (Eur. *Fusarium* Semin., 3rd), Poznan, Poland.
29. Rataj-Guranowska, M., Paszewski, A., and Walkowiak, I. 1991. *Fusarium oxysporum* forms heterokaryons with *Fusarium redolens*. *J. Phytopathol.* 132:294-302.
30. Rosewich, U. L., Pettway, R. E., Kistler, H. C., and Katan, T. 1997. Morphological and molecular characterization of isolates of *Fusarium oxysporum* f.sp. *radicis-lycopersici* from Florida. (Abstr.) Page 83 in: *Abstr. Fungal Genet. Conf.*, 19th. Fungal Genetics Stock Center, Kansas City, KS.
31. Sands, D. C., Ford, E. J., Miller, R. V., Sally, B. K., McCarthy, M. K., Anderson, T. W., Weaver, M. B., Morgan, C. T., Pilgeram, A. L., and Darlington, L. C. 1997. Characterization of a vascular wilt of *Erythroxylum coca* caused by *Fusarium oxysporum* f. sp. *erythroxyl* forma specialis nova. *Plant Dis.* 81:501-504.
32. Tantaoui, A., Ouinten, M., Geiger, J.-P., and Fernandez, D. 1996. Characterization of a single clonal lineage of *Fusarium oxysporum* f. sp. *albedinis* causing Bayoud disease of date palm in Morocco. *Phytopathology* 86:787-792.
33. Venter, S. L., Theron, D. J., Steyn, P. J., Ferreira, D. I., and Eicker, A. 1992. Relationship between vegetative compatibility and pathogenicity of isolates of *Fusarium oxysporum* f. sp. *tuberosi* from potato. *Phytopathology* 82:858-862.
34. Whitehead, D. S., Coddington, A., and Lewis, B. G. 1992. Classification of races by DNA polymorphism analysis and vegetative compatibility grouping in *Fusarium oxysporum* f. sp. *pisi*. *Physiol. Mol. Plant Pathol.* 41:295-305.
35. Woo, S. L., Zoina, A., Del Sorbo, G., Lorito, M., Nanni, B., Scala, F., and Noviello, C. 1996. Characterization of *Fusarium oxysporum* f. sp. *phaseoli* by pathogenic races, VCGs, RFLPs, and RAPD. *Phytopathology* 86:966-973.
36. Woudt, L. P., Neuvel, A., Sikkema, A., van Grinsven, M. Q. J. M., de Milliano, W. A. J., Campbell, C. L., and Leslie, J. F. 1995. Genetic variation in *Fusarium oxysporum* from cyclamen. *Phytopathology* 85:1348-1355.