Interactions of *Pratylenchus thornei* and *Fusarium oxysporum* f. sp. *ciceris* on Chickpea

Pablo Castillo, María P. Mora-Rodríguez, Juan A. Navas-Cortés, and Rafael M. Jiménez-Díaz

Instituto de Agricultura Sostenible, Consejo Superior de Investigaciones Científicas (CSIC), Apdo. 4084, 14080 Córdoba, Spain.
Accepted for publication 27 April 1998.

ABSTRACT

Castillo, P., Mora-Rodríguez, M. P., Navas-Cortés, J. A., and Jiménez-Díaz, R. M. 1998. Interactions of *Pratylenchus thornei* and *Fusarium oxysporum* f. sp. *ciceris* on chickpea. *Phytopathology* 88:828-836.

Fusarium oxysporum f. sp. *ciceris* and the root-lesion nematode *Pratylenchus thornei* coinfect chickpeas in southern Spain. The influence of root infection by *P. thornei* on the reaction of *Fusarium* wilt-susceptible (CPS 1 and PV 61) and wilt-resistant (UC 27) chickpea cultivars to *F. oxysporum* f. sp. *ciceris* race 5 was investigated under controlled and field conditions. Severity of *Fusarium* wilt was not modified by coinfection of chickpeas by *P. thornei* and *F. oxysporum* f. sp. *ciceris*, in simultaneous or sequential inoculations with the pathogens. Root infection with five nematodes per cm³ of soil and 5,000 chlamydo-spores per g of soil of the fungus resulted in significantly higher numbers of propagules of *F. oxysporum* f. sp. *ciceris* with the wilt-susceptible cultivar CPS 1, but not with the wilt-resistant one.

However, infection with 10 nematodes per cm³ of soil significantly increased root infection by *F. oxysporum* f. sp. *ciceris* in both cultivars, irrespective of fungal inoculum densities (250 to 2,000 chlamydo-spores per g of soil). Plant growth was significantly reduced by *P. thornei* infection on wilt-susceptible and wilt-resistant chickpeas in controlled and field conditions, except when shorter periods of incubation (45 days after inoculation) were used under controlled conditions. Severity of root necrosis was greater in wilt-susceptible and wilt-resistant cultivars when nematodes were present in the root, irrespective of length of incubation time (45 to 90 days), densities of nematodes (5 and 10 nematodes per cm³ of soil), fungal inocula, and experimental conditions. Nematode reproduction on the wilt-susceptible cultivars, but not on the wilt-resistant one, was significantly increased by *F. oxysporum* f. sp. *ciceris* infections under controlled and field conditions.

Additional keywords: *Cicer arietinum*, pathogenicity, resistance.

Fusarium wilt, caused by *Fusarium oxysporum* Schlechtend.:Fr. f. sp. *ciceris* (Padwick) Matuo & K. Sato, is a major constraint to production wherever chickpea (*Cicer arietinum* L.) is grown, particularly in the Mediterranean region and the Indian subcontinent (15,30). Yield losses of up to 10% have been attributed to *Fusarium* wilt in India (37) and Spain (40) and up to 40% in Tunisia (4). *F. oxysporum* f. sp. *ciceris* can survive in soil several years by means of chlamydo-spores (16), which makes crop rotation unsuitable as a management strategy for *Fusarium* wilt of chickpea. Therefore, the most practical and economical method for controlling *Fusarium* wilt of chickpea worldwide is the use of resistant cultivars (15,20,21). Root infection by plant-parasitic nematodes has been shown to affect host resistance to some formae speciales of *F. oxysporum* (27). Thus, the occurrence of plant-parasitic nematodes along with *F. oxysporum* f. sp. *ciceris* in soil used for chickpea production is of concern for the durability of *Fusarium* wilt-resistant chickpea cultivars (34).

The cereal and legume root-lesion nematode, *Pratylenchus thornei* Sher & Allen, damages chickpea crops in the Mediterranean region, including southern Spain, and in the Indian subcontinent (7-9,11,12,42). Populations of *P. thornei* in field soils range from 0.02 to 0.48 per cm³ in southern Spain (8). In addition, some root-knot nematodes infecting chickpea are known to interact with root-infecting fungi (26,35,36), thus increasing crop damage.

Both *F. oxysporum* f. sp. *ciceris* and *P. thornei* are widely distributed throughout the major chickpea production areas of the world (8,11,12,30). Thus, the possibility exists that these pathogens, when present together in chickpea fields, may interact synergistically, causing more damage and higher severity of *Fusarium* wilt than would result from infection with either pathogen alone.

However, to our knowledge, no information is available about the interaction between *F. oxysporum* f. sp. *ciceris* and *P. thornei*. The objective of this study was to determine the influence of root infections by *P. thornei* on the reaction of susceptible and resistant chickpea cultivars to *F. oxysporum* f. sp. *ciceris*, including effects on final nematode population densities in soil and roots and root colonization by the fungus.

MATERIALS AND METHODS

Pathogen inoculum and plant material. The most virulent race of *F. oxysporum* f. sp. *ciceris* (race 5) infecting chickpea in southern Spain and a population of *P. thornei* (Cañete population) representative of those found in the Spanish chickpea-growing areas were chosen for the studies (9,19,22). Monoconidial isolate Foc 8012 of *F. oxysporum* f. sp. *ciceris*, representative of race 5 of the pathogen, was used in all studies. This isolate was collected from a chickpea field at Cañete de las Torres (Córdoba) in 1980 and has been used in previous studies (20,40). The isolate was stored in sterile soil at 4°C. Active cultures of the fungus were obtained by placing small aliquots of a soil culture on potato-dextrose agar (PDA) plates that were then incubated at 25°C with a 12-h photoperiod of fluorescent and near-UV light at 36 µE m⁻² s⁻¹ for 7 days. Inoculum consisted of chlamydo-spores obtained as described by Alexander et al. (2). A suspension of autoclaved sandy loam soil in sterile distilled water (1 kg:1 liter) was filtered through filter paper, and the extract was sterilized through a 0.22-µm Millipore filter. The sterile soil extract collected in sterile 250-ml Erlenmeyer flasks (100 ml per flask) was infested with 1 g of fungal mycelial mass grown on Czapeck's medium and incubated on a rotary shaker at 120 rpm for 7 days under the same conditions as the PDA cultures. To obtain mycelial mass, cultures were filtered aseptically through sterile filter paper and washed with sterile distilled water to remove conidia as well as traces of nutrients. The infested soil extract was incubated on a rotary shaker under the

Corresponding author: P. Castillo; E-mail address: ag1cascp@lucano.uco.es

Publication no. P-1998-0619-01R

© 1998 The American Phytopathological Society

same conditions as above. After 2 months of incubation, cultures were examined by light microscopy to assure that they consisted mostly of chlamydospores. Chlamydospores then were collected by centrifugation and inoculum concentration was adjusted using a hemacytometer. Soil was infested by mixing 2 ml of chlamydospore suspension with 5 g of sterile talcum powder and then thoroughly mixing with sterile soil at the appropriate proportion. The inoculum density of *F. oxysporum* f. sp. *ciceris* race 5 in the infested soil was determined before use by dilution-plating on V8 juice-oxgall-PCNB agar (VOPA) (3).

The nematode population used in the studies was obtained from roots of chickpeas (cv. Blanco Lechoso) collected in fields at Cañete de las Torres (Córdoba). Starting from a single female, inoculum was increased several times on carrot disks that were incubated at 24°C each time for 6 weeks (7). Nematodes were extracted from carrot disks using a Baermann funnel, surface-disinfested with 0.02% ethoxyethyl mercury chloride and 0.1% streptomycin solutions for 2 and 24 h, respectively, and then thoroughly rinsed several times in sterile distilled water. Nematode population densities were determined from 1-ml aliquots, and then 10 ml of suspension was placed around the radicle of pregerminated seeds at sowing. Control plants were treated similarly with sterile distilled water.

Fusarium wilt-resistant and -susceptible chickpea cultivars were chosen for the studies on the basis of their known reaction to *F. oxysporum* f. sp. *ciceris* race 5 (19,22; R. M. Jiménez-Díaz, unpublished data). Moderately susceptible cultivars CPS 1 and PV 61 and the resistant cultivar UC 27 (5) were used. All cultivars were susceptible to *P. thornei* (7,8). Seeds of 'CPS 1' and 'UC 27' were kindly provided by H. A. van Rheeën (International Crops Research Institute for Semi-arid Tropics, Hyderabad, India) and I. W. Buddenhagen (University of California, Davis), respectively. Seeds of each cultivar were surface-disinfested with 2% NaOCl solution for 3 min and then germinated on autoclaved layers of moistened filter paper in petri plates at 25°C in the dark for 48 h ('CPS 1' and 'PV 61') or 72 h ('UC 27'). Germinated seeds, selected for uniformity (length of radicle = 1 to 2 cm), were sown into 15-cm-diameter clay pots (one per pot) containing 0.5 liters of an autoclaved potting mixture (sand/clay loam, 2:1, vol/vol).

Growth chamber studies. Three experiments were carried out under a controlled environment in a growth chamber. Cultivars CPS 1 and UC 27 were used in these studies. Growth chambers were adjusted to $24 \pm 1^\circ\text{C}$, 60 to 90% relative humidity, and a 14-h photoperiod of fluorescent light at $360.5 \pm 24.7 \mu\text{E m}^{-2} \text{s}^{-1}$. These environmental conditions are optimum for the development of Fusarium wilt in chickpea (19; J. A. Navas-Cortés and R. M. Jiménez-Díaz, unpublished data). Plants were watered daily with 100 ml of tap water and fertilized weekly with 100 ml of a nutrient solution (17).

In the first experiment, the effects of *F. oxysporum* f. sp. *ciceris* race 5 alone or in simultaneous inoculation with *P. thornei* on the development of Fusarium wilt of chickpea were investigated. Germinated seeds were sown in sterile soil or in soil infested with 5,000 chlamydospores of *F. oxysporum* f. sp. *ciceris* per g of soil. For the joint inoculation, 10 ml of sterile distilled water with nematode inoculum (five nematodes per cm^3 of soil) were added to the soil at sowing. There were four inoculation treatments arranged in a completely randomized design and replicated 10 times, each replicate consisting of a single potted plant. The experiment ended 45 days after inoculation.

The second experiment tested the effects of several inoculum densities of *F. oxysporum* f. sp. *ciceris* race 5 alone or in simultaneous inoculation with *P. thornei*. Because of high disease ratings on Fusarium wilt-susceptible cultivars in the first experiment, a series of lower inoculum densities of *F. oxysporum* f. sp. *ciceris* race 5 were used, and the duration of the experiment was extended to 90 days. Five inoculum densities of *F. oxysporum* f. sp. *ciceris* (0, 250, 500, 1,000, and 2,000 chlamydospores per g of soil) and two of *P. thornei* (0 and 10 nematodes per cm^3 of soil) were included in a complete factorial design comprising 10 inoculation

treatment combinations per cultivar. Inoculations were carried out as for the first experiment. Sterile distilled water (10 ml) was added in treatments without the nematode. Treatments were arranged in a randomized complete block design and replicated five times, each replicate consisting of four pots with a single plant.

The third experiment tested the effects of several inoculum densities of *F. oxysporum* f. sp. *ciceris* race 5 alone or in simultaneous or sequential inoculations with *P. thornei*. Four inoculum densities of *F. oxysporum* f. sp. *ciceris* race 5 (0, 1,000, 2,000, and 5,000 chlamydospores per g soil), two inoculum densities of *P. thornei* (0 and 10 nematodes per cm^3 of soil), and three inoculum timing treatments were included in a 4 by 2 by 3 factorial for a total of 24 inoculation treatments per cultivar. For each combination of fungal and nematode inoculum density, the inocula were introduced simultaneously (one treatment) or sequentially (two treatments). In the two sequential inoculation treatments, seeds were inoculated with one of the pathogens at sowing, and seedlings were grown in the absence of the other pathogen in soil for 10 days. Seedlings were then transplanted with the root ball intact into 15-cm-diameter clay pots containing the same soil mixture that had been infested just with the fungus or the nematode as described above. The duration of the experiment was 90 days. Treatments were arranged in a completely randomized design replicated 10 times, each replicate consisting of a single potted plant.

Field study. To validate data from growth chamber studies under field conditions, an experiment was conducted in a sandy loam soil (pH 8.5, 1.4% organic matter) at the Alameda del Obispo Research Station near Córdoba, Spain (latitude 38° north, longitude 5° east). This field had not been sown to chickpea before. Chickpea cultivars CPS 1, PV 61, and UC 27 were used. In this experiment, plastic bags (10 cm wide by 40 cm long) containing 3 liters of an autoclaved soil mixture (sand/clay loam, 2:1, vol/vol) were used. The experiment was established on 27 March 1996. The bags were placed in rows 1 m apart. Within rows, bags were 40 cm apart and were buried only 30 cm deep. The experiment was arranged in a split-plot design within four randomized complete blocks, with cultivars as the main plot and a factorial of 4 by 2 inoculum density treatment combinations (0, 1,000, 2,000, and 5,000 chlamydospores of *F. oxysporum* f. sp. *ciceris* race 5 per g of soil and 0 and 10 nematodes of *P. thornei* per cm^3 of soil) as subplots. Nematode and fungus inocula were applied as previously described for simultaneous inoculations in growth chamber experiments. There were five replications per treatment combination; each replication was a bag with a single plant. Each cultivar was inoculated with all eight inoculum density treatment combinations. The experiment ended 70 days after inoculation.

Data collection and analysis. Severity of Fusarium wilt was assessed at 5- and 7-day intervals in growth chamber and field experiments, respectively, up to the end of the experiments. Disease severity was rated on a 0 to 4 scale according to the percentage of foliage with yellowing or necrosis (0 = 0%, 1 = 1 to 33%, 2 = 34 to 66%, 3 = 67 to 100%, and 4 = dead plant) (20). Incidence of foliar symptoms (*I*, 0 or 1) and severity data (*S*, rated from 0 to 4), were used to calculate a disease intensity index (*DII*) (1) using the equation $DII = (I \times S)/4$. Additionally, for each treatment, the area under disease progress curve was estimated by trapezoidal integration (6) and standardized by duration time in days of the epidemic (SAUDPC).

When experiments were ended, isolations on VOPA were made from stem sections excised from second and eighth nodes of symptomless plants to determine the occurrence of vascular infections by *F. oxysporum* f. sp. *ciceris* race 5. Stem sections (1 cm) were surface-disinfested in 0.4% NaOCl for 1 min, blotted dry on sterile filter paper, plated on VOPA, and incubated for 3 to 5 days under conditions described previously for fungal isolates. At the end of the experiment, roots were washed free of soil, weighed, and assessed for severity of root necrosis, nematode numbers in roots and soil, and nematode reproduction index (R_f = final population/

initial population). Severity of root necrosis was assessed on a 0 to 10 scale according to the percentage of necrotic tissue visible, in which 0 = 0%, 1 = 1 to 10%, and 10 = 91 to 100%. Nematodes from 100 cm³ of soil and from 5-g root samples were extracted by centrifugation (10) and used for estimating final nematode population densities. CFU per g of fresh root tissue were quantified at the end of the experiments by dilution-plating on VOPA and incubating as before for 3 to 5 days (29).

The first growth chamber experiment was repeated once. Homogeneity of variance between experimental runs, tested by Barlett's test of equal variances, allowed the data to be combined for analyses of variance. Because no symptoms developed on uninocu-

lated control treatments, data obtained from those treatments were not included in the analysis of *Fusarium* wilt severity, to avoid use of zero in analysis of variance. All data on nematode population density were transformed into log₁₀ ($X + 1$). Orthogonal single-degree of freedom contrasts were computed to test the effect of selected experimental treatment combinations of interest (14).

RESULTS

In the first growth chamber experiment, no wilt symptoms developed in either the *Fusarium*-uninfested control plants or the *Fusarium* wilt-resistant cultivar UC 27 (Table 1). Severity of

TABLE 1. Effect of *Fusarium oxysporum* f. sp. *ciceris* race 5 and *Pratylenchus thornei* in simultaneous inoculation of *Fusarium* wilt-susceptible (CPS 1) and wilt-resistant (UC 27) chickpea cultivars in the first growth chamber experiment^a

Inoculation treatment	Inoculum density ^b		<i>DII</i> ^c	SAUDPC ^d	CFU g ⁻¹ root × 10 ⁴	Fresh root weight (g)	Dry shoot weight (g)	Root necrotic severity ^e	Rf ^f
	Pt	Foc							
Cultivar CPS 1									
1	0	0	0.00	0.00	0	11.5	2.4	3.8	... ^g
2	0	5,000	0.83	0.56	181	3.7	2.0	6.5	...
3	5	0	0.00	0.00	0	9.2	2.2	6.9	3.1
4	5	5,000	0.66	0.45	582	5.6	1.7	6.3	5.8
Contrasts (<i>P</i>) ^h									
1 versus 2			...	...	...	<0.01	ns	<0.01	...
1 versus 3			...	...	...	ns	ns	<0.01	...
2 versus 4			ns	ns	<0.01	ns	ns	ns	...
3 versus 4			...	...	...	0.01	0.01	ns	<0.01
Cultivar UC 27									
1	0	0	0.00	0.00	0	18.8	3.4	0.0	...
2	0	5,000	0.00	0.00	5	21.2	3.7	1.2	...
3	5	0	0.00	0.00	0	17.2	3.4	2.3	3.2
4	5	5,000	0.00	0.00	12	17.6	3.3	2.4	2.8
Contrasts (<i>P</i>) ^h									
1 versus 2			...	...	...	ns	ns	<0.01	...
1 versus 3			...	...	...	ns	ns	<0.01	...
2 versus 4			...	...	ns	0.04	ns	<0.01	...
3 versus 4			...	...	...	ns	ns	ns	ns

^a Data are the mean of two experiments with 10 replicated plants per treatment combination in each experiment.

^b Pt = *P. thornei* inoculum expressed as nematodes per cm³ of soil. Foc = *F. oxysporum* f. sp. *ciceris* inoculum expressed as chlamydospores per g of soil.

^c DII = disease intensity index, calculated at the last day of assessment 45 days after inoculation.

^d SAUDPC = area under disease progress curve standardized by duration of the epidemic.

^e Assessed on a scale of 0 (0% necrotic tissue) to 10 (91 to 100% necrotic tissue) 45 days after inoculation. Data were transformed to log($X + 1$) for analysis.

^f Rf (nematode reproduction factor) = final nematode density per plant/initial nematode population density per plant. Data were transformed to log($X + 1$) for analysis.

^g ... = not tested.

^h Orthogonal contrast of inoculation treatments. Probability for the *t* statistic of linear single-degree of freedom contrasts; ns = not significant ($P > 0.05$).

TABLE 2. Influence of different combinations of inocula of *Fusarium oxysporum* f. sp. *ciceris* race 5 and *Pratylenchus thornei* on disease reaction and growth of *Fusarium* wilt-susceptible chickpea cultivar CPS 1 with simultaneous inoculation in the second growth chamber experiment^a

Inoculation treatment	Inoculum density ^b		DII ^c	SAUDPC ^d	CFU g ⁻¹ root × 10 ⁴	Fresh root weight (g)	Dry shoot weight (g)	Root necrotic severity ^e	Rf ^f
	Pt	Foc							
1	0	0	0.00	0.00	0	15.3	3.4	5.9	... ^g
2	0	250	0.66	0.52	77	8.0	1.9	6.4	...
3	0	500	0.62	0.48	63	9.2	2.3	6.3	...
4	0	1,000	0.60	0.46	93	9.7	2.5	6.8	...
5	0	2,000	0.62	0.49	76	8.4	2.3	6.6	...
6	10	0	0.00	0.00	0	10.4	2.7	6.3	29.6
7	10	250	0.63	0.48	122	7.3	1.9	6.9	42.4
8	10	500	0.63	0.48	117	9.1	2.3	6.7	46.9
9	10	1,000	0.63	0.51	119	4.8	1.3	6.8	38.9
10	10	2,000	0.63	0.46	138	5.8	2.0	7.0	45.4
Contrasts (P) ^h									
1 to 5 versus 6 to 10			ns	ns	<0.01	<0.01	<0.01	<0.01	...
1 versus 2 to 5			...	...	...	<0.01	<0.01	<0.01	...
6 versus 7 to 10			...	...	...	<0.01	0.01	<0.01	<0.01

^a Data are the mean of five replicates of four plants each per treatment combination.

^b Pt = *P. thornei* inoculum expressed as nematodes per cm³ of soil. Foc = *F. oxysporum* f. sp. *ciceris* inoculum expressed as chlamydospores per g of soil.

^c DII = disease intensity index, calculated at the last day of assessment 90 days after inoculation.

^d SAUDPC = area under disease progress curve standardized by duration time in days of the epidemic.

^e Assessed on a scale of 0 (0% necrotic tissue) to 10 (91 to 100% necrotic tissue) 90 days after inoculation. Data were transformed to log($X + 1$) for analysis.

^f Rf (nematode reproduction factor) = final nematode density per plant/initial nematode population density per plant. Data were transformed to log($X + 1$) for analysis.

^g ... = not tested.

^h Orthogonal contrast of inoculation treatments. Probability for the *t* statistic of linear single-degree of freedom contrasts; ns = not significant ($P > 0.05$).

Fusarium wilt, determined by the *DII* and SAUDPC, did not differ significantly for the Fusarium wilt-susceptible cultivar CPS 1, whether inoculated with *P. thornei* or not (Table 1). However, coinfection of 'CPS 1' with the fungus and the nematode significantly increased root infection by *F. oxysporum* f. sp. *ciceris* race 5 as determined by the number of CFU/g of root. Inoculation with 5,000 chlamydospores of *F. oxysporum* f. sp. *ciceris* race 5 per g of soil significantly reduced plant growth, determined by fresh root and dry shoot weights in the Fusarium wilt-susceptible, but not wilt-resistant, cultivar. Inoculation with five nematodes per cm³ of soil did not affect plant growth in either cultivar, except for a reduction of root growth in the Fusarium wilt-resistant plants simultaneously inoculated with *F. oxysporum* f. sp. *ciceris* race 5 (Table 1). At the end of the experiment, infection with the fungus or the nematode gave rise to root necrosis that was significantly ($F = 362.3$, $P < 0.001$) more severe in 'CPS 1' than in 'UC 27'. Infection by *P. thornei* significantly increased root necrosis associated with fungus infection in the Fusarium wilt-resistant 'UC 27', but not in the wilt-susceptible 'CPS 1'. However, coinfection by *F. oxysporum* f. sp. *ciceris* race 5 did not increase necrosis caused by *P. thornei* alone (Table 1). Nematode reproduction was significantly higher in plants inoculated with the two pathogens compared with plants inoculated with the nematode alone in the Fusarium wilt-susceptible cultivar only (Table 1).

No wilt symptoms developed in the Fusarium-uninfested control or in the Fusarium wilt-resistant 'UC 27' plants inoculated with 250 to 2,000 chlamydospores of *F. oxysporum* f. sp. *ciceris* race 5 per g of soil in the second experiment. However, regardless of nematode treatment, a distinct yellowing affected the lowermost leaves of inoculated plants 50 days after inoculation. Infection with 10 nematodes per cm³ of soil did not significantly influence the severity of Fusarium wilt, estimated by the *DII* and SAUDPC, on the Fusarium wilt-susceptible cultivar CPS 1 (Table 2). However, root infection by *F. oxysporum* f. sp. *ciceris* race 5, determined by the number of CFU/g of root, was significantly higher in nematode-infected plants of both cultivars, irrespective of the inoculum density of *F. oxysporum* f. sp. *ciceris* race 5 (Tables 2 and 3). Contrary to the first experiment, inoculation with 10 nematodes per cm³ of soil significantly reduced fresh root and dry shoot weights of Fusarium wilt-susceptible and wilt-resistant cultivars 90 days after inoculation. Infection with less than 5,000 chlamydospores of *F. oxysporum* f. sp. *ciceris* race 5 per g of soil reduced plant growth in Fusarium wilt-susceptible 'CPS 1', irrespective of nematode infection. However, in Fusarium wilt-resistant 'UC 27', only

fresh root weight was significantly reduced and that occurred following coinfection with the two pathogens (Table 3). Plants infected by either *P. thornei* or *F. oxysporum* f. sp. *ciceris* race 5 showed increased root necrosis compared with controls at the end of the experiment, and root necrosis was significantly ($F = 2913.3$, $P < 0.001$) more severe in CPS 1 than in UC 27 cultivars (Tables 2 and 3). The severity of root necrosis was increased significantly when plants were simultaneously inoculated with the two pathogens in the Fusarium wilt-susceptible, but not in the wilt-resistant, cultivars (Tables 2 and 3). Nematode reproduction was significantly higher in the wilt-susceptible 'CPS 1' infected with *F. oxysporum* f. sp. *ciceris* race 5, but not in the Fusarium wilt-resistant 'UC 27' (Tables 2 and 3). Actual values of Rf were about 10-fold greater than those from the first experiment for which nematode inoculum density was half of that used in the second experiment.

When the experiment was terminated, *F. oxysporum* f. sp. *ciceris* race 5 was recovered from vascular tissues of 15 out of 20 symptomless 'CPS 1' plants exposed to several inoculum densities of the pathogen, but could be isolated from only 1 of 20 plants of the Fusarium wilt-resistant cultivar UC 27.

In the third growth chamber experiment, no symptoms of Fusarium wilt were found in the Fusarium-uninfested control and the inoculated Fusarium wilt-resistant 'UC 27' plants, irrespective of the inoculum densities (1,000 to 5,000 chlamydospores per g of soil) of *F. oxysporum* f. sp. *ciceris* race 5 or the time interval between inoculations (Tables 4 and 5). As in the second experiment, yellowing of basal leaves was observed in both nematode-inoculated and nematode-free plants by 50 days after inoculation. Severity of Fusarium wilt in the Fusarium wilt-susceptible cultivar CPS 1, determined by the *DII* and SAUDPC, was not significantly influenced by inoculation with 10 nematodes per cm³ of soil, irrespective of *F. oxysporum* f. sp. *ciceris* race 5 inoculum density and inoculation treatment (Table 4). In this cultivar, higher Fusarium wilt severity resulted from the simultaneous inoculation with *F. oxysporum* f. sp. *ciceris* race 5 and *P. thornei* as compared with that in the two sequential inoculations with each of the pathogens (Table 4). Severity of Fusarium wilt was significantly higher when seedlings were inoculated with the nematode prior to that with the fungus as compared with the sequence of the fungus followed by the nematode (Table 4). As for the second growth chamber experiment, growth of 'CPS 1' plants was significantly less in both *P. thornei* and *F. oxysporum* f. sp. *ciceris* race 5-infected plants, regardless of the inoculation treatment, with one exception. When infection by *F. oxysporum* f. sp. *ciceris* race 5 preceded nematode infection,

TABLE 3. Influence of different combinations of inocula of *Fusarium oxysporum* f. sp. *ciceris* race 5 and *Pratylenchus thornei* on disease reaction and growth of Fusarium wilt-resistant chickpea cultivar UC 27 with simultaneous inoculation in the second growth chamber experiment^a

Inoculation treatment	Inoculum density ^b		CFU g ⁻¹ root × 10 ⁴	Fresh root weight (g)	Dry shoot weight (g)	Root necrotic severity ^c	Rf ^d
	Pt	Foc					
1	0	0	0	37.0	5.1	0.0	...
2	0	250	1	35.5	5.3	1.1	...
3	0	500	1	30.8	5.0	1.4	...
4	0	1,000	1	37.3	5.2	0.9	...
5	0	2,000	3	31.2	5.0	1.4	...
6	10	0	0	30.8	4.4	1.9	31.4
7	10	250	4	27.9	4.7	1.6	29.7
8	10	500	3	21.7	4.6	1.6	33.0
9	10	1,000	1	23.6	4.3	2.7	38.2
10	10	2,000	5	21.1	4.3	2.6	31.5
Contrasts (P) ^f							
1 to 5 versus 6 to 10			0.01	<0.01	<0.01	<0.01	...
1 versus 2 to 5			...	ns	ns	<0.01	...
6 versus 7 to 10			...	<0.01	ns	ns	ns

^a Data are the mean of five replicates of four plants each per treatment combination.

^b Pt = *P. thornei* inoculum expressed as nematodes per cm³ of soil. Foc = *F. oxysporum* f. sp. *ciceris* inoculum expressed as chlamydospores per g of soil.

^c Assessed on a scale of 0 (0% necrotic tissue) to 10 (91 to 100% necrotic tissue) 90 days after inoculation. Data were transformed to log(X + 1) for analysis.

^d Rf (nematode reproduction factor) = final nematode density per plant/initial nematode population density per plant. Data were transformed to log(X + 1) for analysis.

^e ... = not tested.

^f Orthogonal contrast of inoculation treatments. Probability for the *t* statistic of linear single-degree of freedom contrasts; ns = not significant ($P > 0.05$).

shoot dry weight was comparable to that in plants infected by only *P. thornei* (Table 4). Also, plant growth was significantly higher in those plants inoculated with *F. oxysporum* f. sp. *ciceris* race 5 and then transplanted 10 days later to either sterile soil or soil infested with *P. thornei* compared with growth in plants inoculated first with *P. thornei* or simultaneously with the two pathogens (Table 4).

In the Fusarium wilt-resistant cultivar UC 27 (Table 5), nematode-infected plants had significantly lower root fresh weights when seedlings were inoculated with the fungus and the nematode simultaneously and when inoculation with the nematode preceded that with the fungus. No differences in root fresh weight were observed when seedlings were inoculated with the fungus first (Table 5). Shoot dry weight of *F. oxysporum* f. sp. *ciceris* race 5-infected plants was significantly lower when seedlings were inoculated simultaneously with both pathogens (Table 5). Fresh root and dry

shoot weights were significantly lower in plants simultaneously inoculated with both pathogens than in those inoculated sequentially. There were no significant differences between the two sequential inoculation treatments with respect to plant growth (Table 5).

As in the second growth chamber experiment, no necrosis developed in the roots of controls. However, the severity of root necrosis was significantly greater in plants infected with the nematode, the fungus, or both, irrespective of the inoculation treatment, at the end of the experiment. Root necrosis was significantly less severe in simultaneous as compared with sequential inoculations (Table 5). The nematode reproduction factor was of the same order of magnitude as in the second experiment. It was significantly greater in Fusarium wilt-susceptible 'CPS 1' plants inoculated with *F. oxysporum* f. sp. *ciceris* race 5 in simultaneous as well as in sequential inoculation with the fungus followed by the nematode;

TABLE 4. Effects of different concentrations of inocula of *Fusarium oxysporum* f. sp. *ciceris* race 5 and *Pratylenchus thornei* on the disease reaction and growth of Fusarium wilt-susceptible chickpea cultivar CPS-1 in simultaneous and sequential inoculations in the third growth chamber experiment^a

Inoculation treatment	Inoculum density ^b		<i>DII</i> ^c	SAUDPC ^d	Fresh root weight (g)	Dry shoot weight (g)	Root necrotic severity ^e	Rf ^f
	Pt	Foc						
Simultaneous inoculation								
1	0	0	0.00	0.00	15.16	3.09	4.28	... ^g
2	0	1,000	0.76	0.57	0.90	1.03	4.67	...
3	0	2,000	0.66	0.40	0.82	1.49	3.89	...
4	0	5,000	0.74	0.56	0.59	1.01	4.33	...
5	10	0	0.00	0.00	12.44	1.83	4.72	30.1
6	10	1,000	0.79	0.49	1.48	1.56	5.22	78.6
7	10	2,000	0.88	0.61	0.61	1.27	4.89	65.8
8	10	5,000	0.94	0.74	0.26	0.55	5.22	64.6
Contrasts (<i>P</i>) ^h								
1 to 4 versus 5 to 8			...	...	0.04	<0.01	<0.01	...
1 versus 2 to 4			...	...	<0.01	<0.01	ns	...
5 versus 6 to 8			...	...	<0.01	<0.01	ns	<0.01
2 to 4 versus 6 to 8			ns	ns	...	...	...	...
Sequential inoculation 1 (nematode prior fungus)								
9	0	0	0.00	0.00	16.53	3.57	4.89	...
10	0	1,000	0.51	0.34	1.67	1.79	6.22	...
11	0	2,000	0.76	0.54	2.62	1.73	6.33	...
12	0	5,000	0.61	0.38	1.36	1.42	6.11	...
13	10	0	0.00	0.00	9.16	2.89	6.22	...
14	10	1,000	0.64	0.43	0.84	1.45	6.22	...
15	10	2,000	0.74	0.55	0.78	1.51	6.11	...
16	10	5,000	0.84	0.52	0.91	1.08	6.33	...
Contrasts (<i>P</i>) ^h								
9 to 12 versus 13 to 16			...	...	<0.01	0.01	0.04	...
9 versus 10 to 12			...	...	<0.01	<0.01	<0.01	...
13 versus 14 to 16			...	...	<0.01	<0.01	ns	...
10 to 12 versus 14 to 16			ns	ns	...	...	...	...
Sequential inoculation 2 (fungus prior nematode)								
17	0	0	0.00	0.00	15.92	3.54	4.44	...
18	0	1,000	0.19	0.14	13.66	3.41	6.11	...
19	0	2,000	0.26	0.21	14.96	3.51	6.44	...
20	0	5,000	0.11	0.10	13.03	3.23	6.56	...
21	10	0	0.00	0.00	12.39	2.96	5.83	41.2
22	10	1,000	0.15	0.12	9.67	3.06	6.33	77.6
23	10	2,000	0.15	0.09	11.02	3.20	6.56	66.3
24	10	5,000	0.21	0.17	10.02	3.08	6.50	64.1
Contrasts (<i>P</i>) ^h								
17 to 20 versus 21 to 24			...	...	0.02	ns	ns	...
17 versus 18 to 20			...	...	ns	ns	<0.01	...
21 versus 22 to 24			...	...	ns	ns	0.04	<0.01
18 to 20 versus 22 to 24			ns	ns	...	...	...	...
General contrasts between inoculations (<i>P</i>) ^h								
1 to 8 versus 9 to 24			<0.01	<0.01	<0.01	<0.01	<0.01	...
9 to 16 versus 17 to 24			<0.01	<0.01	<0.01	<0.01	ns	...

^a Data are the mean of 10 replicates, each replicate consisting of one plant per treatment combination.

^b Pt = *P. thornei* inoculum expressed as nematodes per cm³ of soil. Foc = *F. oxysporum* f. sp. *ciceris* inoculum expressed as chlamydospores per g of soil.

^c *DII* = disease intensity index, calculated at the last day of assessment 90 days after inoculation.

^d SAUDPC = area under disease progress curve standardized by duration of the epidemic.

^e Assessed on a scale of 0 (0% necrotic tissue) to 10 (91 to 100% necrotic tissue) 90 days after inoculation. Data were transformed to log(*X* + 1) for analysis.

^f Rf (nematode reproduction factor) = final nematode density per plant/initial nematode population density per plant. Data were transformed to log(*X* + 1) for analysis.

^g ... = not tested.

^h Orthogonal contrast of inoculation treatments. Probability for the *t* statistic of linear single-degree of freedom contrasts; ns = not significant (*P* > 0.05).

but it was unaffected by *F. oxysporum* f. sp. *ciceris* race 5 infection in the Fusarium wilt-resistant cultivar 'UC 27' (Tables 4 and 5). Nematode reproduction could not be determined in the sequential inoculation with the nematode followed by the fungus, because the number of nematodes penetrating into roots was not estimated at transplanting (Tables 4 and 5).

No symptoms of Fusarium wilt developed in the field study in the *Fusarium*-uninfested control or the Fusarium wilt-resistant 'UC 27' plants inoculated with 1,000 to 5,000 chlamydospores of *F. oxysporum* f. sp. *ciceris* race 5 per g of soil. Inoculation with 10 nematodes per cm³ of soil did not significantly affect the severity of Fusarium wilt, as determined by the *DII* and SAUDPC, of either Fusarium wilt-susceptible cultivars (CPS 1 and PV 61) (Table 6). Nevertheless, nematode infection significantly reduced fresh root and dry shoot weights of Fusarium wilt-susceptible and wilt-resistant cultivars. Similarly, growth of Fusarium wilt-susceptible cultivars was significantly reduced by infection with *F. oxysporum* f. sp. *ciceris* race 5 either alone or together with *P. thornei*. However, infection of Fusarium wilt-resistant 'UC 27' by *F. oxysporum*

f. sp. *ciceris* race 5 reduced fresh root weight only, and the joint infection by the fungus and the nematode did not increase damage caused by *P. thornei* alone (Table 6). Little root necrosis developed in control plants by the end of the experiment. However, infection by either pathogen alone significantly increased root necrosis in the Fusarium wilt-susceptible and, to a lesser extent, in the wilt-resistant cultivars (Table 6). Infection with *F. oxysporum* f. sp. *ciceris* race 5 significantly increased root necrosis caused by *P. thornei* in the two Fusarium wilt-susceptible cultivars, but not in the wilt-resistant one (Table 6). Nematode reproduction was significantly increased by infection with *F. oxysporum* f. sp. *ciceris* race 5 in Fusarium wilt-susceptible plants, but not in Fusarium wilt-resistant plants (Table 6). The nematode reproduced to a much lesser extent under field conditions than in controlled conditions.

DISCUSSION

This study was designed to evaluate whether or not infection of chickpea roots by *P. thornei* would influence the plant reaction to

TABLE 5. Effects of different concentrations of inocula of *Fusarium oxysporum* f. sp. *ciceris* race 5 and *Pratylenchus thornei* on the disease reaction and growth of Fusarium wilt-resistant chickpea cultivar UC 27 in simultaneous and sequential inoculations in the third growth chamber experiment^a

Inoculation treatment	Inoculum density ^b		Fresh root weight (g)	Dry shoot weight (g)	Root necrotic severity ^c	Rf ^d
	Pt	Foc				
Simultaneous inoculation						
1	0	0	34.32	5.08	0.00	... ^e
2	0	1,000	23.41	4.06	1.78	...
3	0	2,000	29.26	4.67	2.17	...
4	0	5,000	23.50	3.86	2.28	...
5	10	0	24.51	4.59	2.68	36.7
6	10	1,000	27.28	4.71	2.68	34.4
7	10	2,000	26.19	4.37	2.39	44.5
8	10	5,000	18.77	3.46	2.61	31.0
Contrasts (<i>P</i>) ^f						
1 to 4 versus 5 to 8			0.01	ns	<0.01	...
1 versus 2 to 4			<0.01	<0.01	<0.01	...
5 versus 6 to 8			ns	ns	ns	ns
Sequential inoculation 1 (nematode prior fungus)						
9	0	0	35.99	5.39	0.00	...
10	0	1,000	35.30	5.66	2.28	...
11	0	2,000	28.72	5.18	2.44	...
12	0	5,000	34.31	5.34	2.28	...
13	10	0	30.67	5.52	3.56	...
14	10	1,000	23.94	5.38	4.33	...
15	10	2,000	29.82	5.01	3.94	...
16	10	5,000	27.44	5.43	2.78	...
Contrasts (<i>P</i>) ^f						
9 to 12 versus 13 to 16			0.01	ns	<0.01	...
9 versus 10 to 12			ns	ns	<0.01	...
13 versus 14 to 16			ns	ns	ns	...
Sequential inoculation 2 (fungus prior nematode)						
17	0	0	35.88	5.24	0.00	...
18	0	1,000	35.56	4.67	2.11	...
19	0	2,000	31.10	5.74	2.17	...
20	0	5,000	35.50	5.68	1.94	...
21	10	0	31.48	5.54	3.39	38.7
22	10	1,000	33.94	5.52	3.33	36.6
23	10	2,000	30.61	5.60	3.39	51.4
24	10	5,000	29.58	5.51	3.28	32.8
Contrasts (<i>P</i>) ^f						
17 to 20 versus 21 to 24			ns	ns	<0.01	...
17 versus 18 to 20			ns	ns	<0.01	...
21 versus 22 to 24			ns	ns	ns	ns
General contrasts between inoculations (<i>P</i>) ^f						
1 to 8 versus 9 to 24			<0.01	<0.01	<0.01	...
9 to 16 versus 17 to 24			ns	ns	<0.01	...

^a Data are the mean of 10 replicates, each replicate consisting of one plant per treatment combination.

^b Pt = *P. thornei* inoculum expressed as nematodes per cm³ of soil. Foc = *F. oxysporum* f. sp. *ciceris* inoculum expressed as chlamydospores per g of soil.

^c Assessed on a scale of 0 (0% necrotic tissue) to 10 (91 to 100% necrotic tissue) 90 days after inoculation. Data were transformed to log(*X* + 1) for analysis.

^d Rf (nematode reproduction factor) = final nematode density per plant/initial nematode population density per plant. Data were transformed to log(*X* + 1) for analysis.

^e ... = not tested.

^f Orthogonal contrast of inoculation treatments. Probability for the *t* statistic of linear single-degree of freedom contrasts; ns = not significant (*P* > 0.05).

the highly virulent race 5 of *F. oxysporum* f. sp. *ciceris*. Both growth chamber and field experiments clearly demonstrated that neither wilt reaction of Fusarium wilt-susceptible or wilt-resistant chickpea cultivars nor severity of Fusarium wilt in susceptible cultivars were modified by coinfection of plants with *F. oxysporum* f. sp. *ciceris* race 5 and *P. thornei*. Furthermore, Fusarium wilt severity was similar whether chickpea seedlings were inoculated simultaneously or sequentially with both pathogens. That occurred irrespective of whether the plant-fungus-nematode interactions took place under continuous, optimum conditions for Fusarium wilt development in the growth chamber or under a fluctuating environment in the field. No symptoms characteristic of Fusarium wilt were found in the Fusarium wilt-resistant 'UC 27' in growth chamber or field experiments. However, when 'UC 27' plants were maintained for longer than 50 days following inoculation, chlorosis of basal leaves developed. These symptoms were observed in nematode- and fungal-infected plants as well as in uninfected controls and, therefore, can be attributed to natural senescence.

Interactions between *Pratylenchus* spp. and several formae speciales of *F. oxysporum* have been described on several crops (28,

31,33,38). These studies demonstrated that the infection by root-lesion nematodes increased the incidence or severity of Fusarium wilt on susceptible cultivars. However, those previously published results were not confirmed in other compatible plant-nematode-fungus combinations (18). Thus, it appears that modification of Fusarium wilt incidence or severity may be related to the specific nematode-fungus combination. Furthermore, these controversial results indicate that interactions between soilborne fungi and root-lesion nematodes are biological and physiological rather than physical in nature (27). In our study, we used inoculum densities of *F. oxysporum* f. sp. *ciceris* race 5 and *P. thornei* higher than those found in soil in southern Spain (8; R. M. Jiménez-Díaz, unpublished data) or than that optimum for Fusarium wilt development (19; J. A. Navas-Cortés and R. M. Jiménez-Díaz, unpublished data) in the simultaneous and sequential inoculations of chickpeas. While none of the inoculation treatments influenced the plant reaction to the fungus, the nematode inoculum density influenced the root colonization by *F. oxysporum* f. sp. *ciceris*. Thus, root infections with five nematodes per cm³ of soil and 5,000 chlamydospores per g of soil resulted in a higher number of propagules of *F. oxysporum* f.

TABLE 6. Effects of different combinations of inocula of *Fusarium oxysporum* f. sp. *ciceris* race 5 and *Pratylenchus thornei* on the disease reaction and growth of Fusarium wilt-susceptible (CPS 1 and PV 61) and wilt-resistant (UC 27) chickpea cultivars in simultaneous inoculation under field conditions^a

Inoculation treatment	Inoculum density ^b		DIF ^c	SAUDPC ^d	Fresh root weight (g)	Dry shoot weight (g)	Root necrotic severity ^e	Rf ^f
	Pt	Foc						
Cultivar CPS 1								
1	0	0	0.00	0.00	5.11	2.20	3.65	... ^g
2	0	1,000	0.89	0.57	1.38	0.65	7.08	...
3	0	2,000	0.93	0.56	0.81	0.65	6.80	...
4	0	5,000	0.98	0.67	0.55	0.51	7.05	...
5	10	0	0.00	0.00	3.49	1.71	4.20	2.53
6	10	1,000	0.88	0.56	0.91	0.55	7.60	3.57
7	10	2,000	0.91	0.64	0.65	0.56	7.25	3.37
8	10	5,000	0.94	0.65	0.38	0.36	7.38	2.88
Contrasts (P) ^h								
1 to 4 versus 5 to 8			ns	ns	<0.01	0.03	0.01	...
1 versus 2 to 4			...	...	<0.01	<0.01	<0.01	...
5 versus 6 to 8			...	...	<0.01	<0.01	<0.01	<0.01
Cultivar UC 27								
1	0	0	0.00	0.00	9.82	4.82	1.28	...
2	0	1,000	0.00	0.00	8.53	3.84	2.55	...
3	0	2,000	0.00	0.00	8.31	5.74	2.15	...
4	0	5,000	0.00	0.00	8.73	6.14	2.25	...
5	10	0	0.00	0.00	7.01	4.68	3.23	4.30
6	10	1,000	0.00	0.00	7.14	4.55	2.63	5.03
7	10	2,000	0.00	0.00	8.01	4.70	2.88	5.31
8	10	5,000	0.00	0.00	6.93	3.75	2.60	4.41
Contrasts (P) ^h								
1 to 4 versus 5 to 8			...	...	<0.01	0.03	<0.01	...
1 versus 2 to 4			...	...	0.02	ns	<0.01	...
5 versus 6 to 8			...	...	ns	ns	ns	ns
Cultivar PV 61								
1	0	0	0.00	0.00	12.35	4.01	1.45	...
2	0	1,000	0.87	0.56	2.75	1.08	4.95	...
3	0	2,000	0.93	0.65	1.48	0.95	5.90	...
4	0	5,000	1.00	0.73	0.54	0.45	4.35	...
5	10	0	0.00	0.00	8.45	2.60	2.60	2.07
6	10	1,000	0.91	0.59	2.03	0.77	4.95	3.47
7	10	2,000	0.88	0.57	2.39	1.28	5.50	4.15
8	10	5,000	0.98	0.65	1.17	0.74	5.20	5.44
Contrasts (P) ^h								
1 to 4 versus 5 to 8			ns	ns	0.03	0.03	<0.01	...
1 versus 2 to 4			...	...	<0.01	<0.01	<0.01	...
5 versus 6 to 8			...	...	<0.01	<0.01	<0.01	<0.01

^a Data are the mean of 10 replicates, each replicate consisting of one plant per treatment combination.

^b Pt = *P. thornei* inoculum expressed as nematodes per cm³ of soil. Foc = *F. oxysporum* f. sp. *ciceris* inoculum expressed as chlamydospores per g of soil.

^c DII = disease intensity index, calculated at the last day of assessment 90 days after inoculation.

^d SAUDPC = area under disease progress curve standardized by duration of the epidemic.

^e Assessed on a scale of 0 (0% necrotic tissue) to 10 (91 to 100% necrotic tissue) 70 days after inoculation. Data were transformed to log(X + 1) for analysis.

^f Rf (nematode reproduction factor) = final nematode density per plant/initial nematode population density per plant. Data were transformed to log(X + 1) for analysis.

^g ... = not tested.

^h Orthogonal contrast of inoculation treatments. Probability for the *t* statistics of linear single-degree of freedom contrasts; ns = not significant (*P* > 0.05).

sp. *ciceris* race 5 in roots on Fusarium-susceptible cultivars, but not on the Fusarium wilt-resistant cultivar. However, infections with 10 nematodes per cm³ of soil significantly increased root infection by *F. oxysporum* f. sp. *ciceris* race 5 in both cultivars, irrespective of fungal inoculum densities. These results differ from observations made with other crops, for which numbers of fungal propagules were not affected by coinfection with root-lesion nematodes (28,33). This suggests that *P. thornei* might modify the physiology of cortical cells of chickpea to facilitate more extensive colonization by *F. oxysporum* f. sp. *ciceris* race 5. Also, it appears that the nature of such modifications may be influenced by the extent of infection by the nematode as related with the nematode inoculum density.

Chickpea growth was reduced by *P. thornei* infection on both the Fusarium wilt-susceptible and wilt-resistant cultivars tested under growth chamber and field conditions. However, in growth chamber studies, reduction of plant growth was observed only when inoculated plants were incubated for a long period. This result is similar to that found in other studies (9,12,39,42) and indicates that results in some of our previous studies may have been influenced by differences in the length of the observation period during experiments (7,8). Similarly, reduced plant growth was observed in Fusarium wilt-susceptible plants infected with *F. oxysporum* f. sp. *ciceris* race 5, but that did not occur with the Fusarium wilt-resistant cultivar used, irrespective of fungal inoculum density, incubation period, or time interval between inoculations.

Infection of chickpea by *P. thornei* increased the severity of root necrosis on both the Fusarium wilt-susceptible cultivars CPS 1 and PV 61 and the wilt-resistant cultivar UC 27, irrespective of incubation period, nematode and fungal inoculum densities, and environmental conditions. These results support other researchers who found a positive correlation between severity of root necrosis and nematode population density in roots of chickpea (12). However, as some necrosis occurred in uninfected control plants, the extended necrosis in root systems of Fusarium wilt-susceptible cultivars CPS 1 and PV 61 might be related to the occurrence of phenolic compounds in the root (25,32).

Reproduction of *P. thornei* was higher in roots of *F. oxysporum* f. sp. *ciceris* race 5-infected susceptible cultivars, but not in the resistant UC 27 cultivar, irrespective of experimental conditions, incubation period, nematode and fungal inoculum densities, or time interval between inoculations. This supports other authors who reported that reproduction of *Pratylenchus* spp. was stimulated in some crops by simultaneous root infections of *Fusarium* spp., causing cortical necrosis (18,23,24) and vascular infections of *Verticillium dahliae* Kleb (13,41). Infection of wilt-susceptible chickpea by *F. oxysporum* f. sp. *ciceris* race 5 may have made the root cortex of chickpea more suitable as host tissue for *P. thornei* as compared with healthy roots. That infection of the wilt-resistant cultivar by *F. oxysporum* f. sp. *ciceris* race 5 did not enhance the nematode reproduction might be due either to the limited extent of root infection by the fungus, the defense mechanisms involved in the resistant reaction, or both.

No modifications of the severity of Fusarium wilt was found in these experiments in spite of the enhancement of root colonization by the highly virulent race 5 of *F. oxysporum* f. sp. *ciceris* in the wilt-susceptible cultivars. Thus, it appears that the durability of Fusarium wilt-resistant chickpea cultivars is unlikely to be compromised by infection with root-lesion nematodes. Nevertheless, the subsequent enhancement of nematode reproduction and fungus colonization in Fusarium wilt-susceptible chickpea is a significant observation, because of higher inoculum production of both pathogens in soil and roots, and warrants further investigations.

ACKNOWLEDGMENTS

Research was supported by grant AGF97-1479 from Comisión Interministerial de Ciencia y Tecnología (CICYT), Spain. We thank R. Rowe,

Ohio State University, and B. Hau, University of Hannover, for reviewing the manuscript prior to submission.

LITERATURE CITED

- Acimovic, M. 1979. Evaluation procedures for the intensity of disease occurrence in sunflower. *Helia* 2:55-57.
- Alexander, J. V., Bourret, J. A., Gold, A. H., and Snyder, W. C. 1966. Induction of chlamydospore formation by *Fusarium solani* in sterile soil extracts. *Phytopathology* 56:353-354.
- Bouhot, D., and Rouxel, F. 1971. Technique sélective et quantitative d'analyse des *Fusarium oxysporum* et *Fusarium solani* dans le sol. Mode d'emploi. *Ann. Phytopathol.* 3:251-254.
- Bouslama, M. 1980. Chickpea improvement in Tunisia. Pages 277-280 in: Proc. Int. Workshop Chickpea Improv. ICRISAT (Int. Crops Res. Inst. Semi-Arid Tropics), Hyderabad, India.
- Buddenhagen, I. W., and Workneh, F. 1988. Fusarium wilt of chickpea in California. (Abstr.) *Phytopathology* 78:1563.
- Campbell, C. L., and Madden, L. V. 1990. Introduction to Plant Disease Epidemiology. John Wiley & Sons, New York.
- Castillo, P., Gómez Barcina, A., and Jiménez-Díaz, R. M. 1996. Plant-parasitic nematodes associated with chickpea in southern Spain and effects of soil temperature on reproduction of *Pratylenchus thornei*. *Nematologica* 42:211-219.
- Castillo, P., Jiménez-Díaz, R. M., Gómez Barcina, A., and Vovlas, N. 1995. Parasitism of the root-lesion nematode *Pratylenchus thornei* on chickpea. *Plant Pathol.* 44:728-733.
- Castillo, P., Vovlas, N., and Jiménez-Díaz, R. M. 1998. Pathogenicity and histopathology of *Pratylenchus thornei* populations on selected chickpea genotypes. *Plant Pathol.* 47:370-376.
- Coolen, W. A. 1979. Methods for the extraction of *Meloidogyne* spp. and other nematodes from roots and soil. Pages 317-329 in: Root-Knot Nematodes (*Meloidogyne* Species) Systematics, Biology and Control. F. Lamberti and C. E. Taylor, eds. Academic Press, London.
- Di Vito, M., Greco, N., Halila, H. M., Mabsoute, L., Labdi, M., Beniwal, S. P. S., Saxena, M. C., Singh, K. B., and Solh, M. B. 1994. Nematodes of cool-season food legumes in North Africa. *Nematol. Mediterr.* 22:3-10.
- Di Vito, M., Greco, N., and Saxena, M. C. 1992. Pathogenicity of *Pratylenchus thornei* on chickpea in Syria. *Nematol. Mediterr.* 20:71-73.
- Faulkner, L. R., and Skotland, C. B. 1965. Interaction of *Verticillium dahliae* and *Pratylenchus minyus* in *Verticillium* wilt of peppermint. *Phytopathology* 55:583-586.
- Gomez, K. A., and Gomez, A. A. 1984. Statistical Procedures for Agricultural Research. 2nd ed. John Wiley & Sons, New York.
- Haware, M. P. 1990. Fusarium wilt and other important diseases of chickpea in the Mediterranean area. *Options Mediterr. Ser. Semin.* 9:61-64.
- Haware, M. P., Nene, Y. L., and Natarayan, M. 1966. The survival of *Fusarium oxysporum* f. sp. *ciceris* in the soil in the absence of chickpea. *Phytopathol. Mediterr.* 35:9-12.
- Hoagland, D. R., and Arnon, D. I. 1950. The water culture method for growing plants without soil. *Calif. Agric. Exp. Stn. Circ.* No. 347.
- Hutton, D. G., Wilkinson, R. E., and Mai, W. F. 1973. Effect of two plant-parasitic nematodes on Fusarium dry root rot of beans. *Phytopathology* 63:749-751.
- Jiménez-Díaz, R. M., Crinó, P., Halila, M. H., Mosconi, C., and Trapero-Casas, A. 1993. Screening for resistance to Fusarium wilt and Ascochyta blight in chickpea. Pages 77-96 in: Breeding for Stress Tolerance in Cool-Season Food Legumes. K. B. Singh and M. C. Saxena, eds. John Wiley & Sons, Chichester, England.
- Jiménez-Díaz, R. M., Singh, K. B., Trapero-Casas, A., and Trapero-Casas, J. L. 1991. Resistance in kabuli chickpeas to Fusarium wilt. *Plant Dis.* 75:914-918.
- Jiménez-Díaz, R. M., and Trapero-Casas, A. 1990. Improvement of chickpea resistance to wilt and root rot diseases. *Options Mediterr. Ser. Semin.* 9:65-72.
- Jiménez-Díaz, R. M., Trapero-Casas, A., and Cabrera de la Colina, J. 1989. Races of *Fusarium oxysporum* f. sp. *ciceris* infecting chickpea in southern Spain. Pages 515-520 in: Vascular Wilt Diseases of Plants. NATO ASI Ser. H, vol. 28. E. C. Tjamas and C. H. Beckman, eds. Springer-Verlag, Berlin.
- Jin, X., Kotcon, J. B., and Morton, J. B. 1991. Interactions between *Pratylenchus penetrans* and *Fusarium avenaceum* in red clover. *Nematropica* 21:105-109.
- Jordan, E. M., Loots, G. C., Jooste, W. J., and De Waele, D. 1987. Effects of root-lesion nematodes (*Pratylenchus brachyurus* Godfrey and *P. zeae* Graham) and *Fusarium moniliforme* Sheldon alone or in combination, on maize. *Nematologica* 33:213-219.
- Kumar, A., and Jalih, B. L. 1985. Differential effects of seed leachates on seed-borne pathogens associated with chickpea (*Cicer arietinum* L.). *Indian Phytopathol.* 38:99-103.
- Kumar, R., Ahmad, S., and Saxena, S. K. 1988. Disease complex in chick-

- pea involving *Meloidogyne javanica* with *Fusarium oxysporum*. Int. Nematol. Network Newsl. 5:12-14.
27. Mai, W. F., and Abawi, G. S. 1987. Interaction among root-knot nematodes and *Fusarium* wilt fungi on host plants. Annu. Rev. Phytopathol. 25:317-338.
 28. Mauza, B. E., and Webster, J. M. 1982. Suppression of alfalfa growth by concomitant populations of *Pratylenchus penetrans* and two *Fusarium* species. J. Nematol. 14:364-367.
 29. Nelson, P. E., Toussoun, T. A., Burgess, L. W., Marasas, W. F. O., and Liddell, C. M. 1986. Isolating, identifying, and producing inoculum of pathogenic species of *Fusarium*. Pages 54-59 in: Methods for Evaluating Pesticides for Control of Plant Pathogens. K. D. Hickey, ed. The American Phytopathological Society, St. Paul, MN.
 30. Nene, Y. L., and Reddy, M. V. 1987. Chickpea diseases and their control. Pages 233-270 in: The Chickpea. M. C. Saxena and K. B. Singh, eds. Commonw. Agric. Bur. Int., Wallingford, England.
 31. Oyekan, P. O., and Mitchell, J. E. 1971. Effect of *Pratylenchus penetrans* on the resistance of a pea variety to *Fusarium* wilt. Plant Dis. Rep. 55: 1032-1035.
 32. Sankuning, K., Thamizheran, N., and Govindarajaln, T. 1986. Release of phenols from chickpea seeds. Int. Chickpea Newsl. 14:24-25.
 33. Seinhorst, J. W., and Kuniyasu, K. 1971. Interaction on *Pratylenchus penetrans* and *Fusarium oxysporum* f. sp. *pisi* race 2 and *Rotylenchus uniformis* and *F. oxysporum* f. sp. *pisi* race 1 on peas. Nematologica 17:444-452.
 34. Sharma, S. B., and Cerkauskas, R. F. 1985. Interação entre *Meloidogyne javanica* e *Fusarium oxysporum* f. sp. *ciceris* sobre o grão-de bico. Nematol. Brasil. 9:113-121.
 35. Sharma, S. B., and McDonald, D. 1990. Global status of nematode problems of groundnut, pigeonpea, chickpea, sorghum and pearl millet, and suggestions for future work. Crop Prot. 9:453-458.
 36. Sharma, S. B., Smith, D. H., and McDonald, D. 1992. Nematode constraints of chickpea and pigeonpea production in the semiarid tropics. Plant Dis. 76:868-874.
 37. Singh, K. B., and Dahiya, B. S. 1973. Breeding for wilt resistance in chickpea. Pages 13-14 in: Symposium on Wilt Problem and Breeding for Wilt Resistance in Bengal Gram. Indian Agric. Res. Inst., New Dehli, India.
 38. Sumner, D. R., and Minton, N. A. 1987. Interaction of *Fusarium* wilt and nematodes in Cobb soybean. Plant Dis. 71:20-23.
 39. Tiyyagi, S. A., and Parveen, M. 1992. Pathogenic effect of root-lesion nematode *Pratylenchus thornei* on plant growth, water absorption capability, and chlorophyll content of chickpea. Int. Chickpea Newsl. 26:18-20.
 40. Trapero-Casas, A., and Jiménez-Díaz, R. M. 1985. Fungal wilt and root rot diseases of chickpea in southern Spain. Phytopathology 75:1146-1151.
 41. Vrain, T. C. 1987. Effect of *Ditylenchus dipsaci* and *Pratylenchus penetrans* on Verticillium wilt of alfalfa. J. Nematol. 19:379-383.
 42. Walia, R. K., and Seshadri, A. R. 1985. Pathogenicity of the root-lesion nematode *Pratylenchus thornei* on chickpea. Int. Chickpea Newsl. 12:31.